



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Liu, M;Lycett, K;Wong, TY;Grobler, A;Juonala, M;He, M;Dwyer, T;Burgner, D;Wake, M

Title:

Associations of retinal microvascular caliber with large arterial function and structure: A population-based study of 11 to 12 year-olds and midlife adults

Date:

2020-08-01

Citation:

Liu, M., Lycett, K., Wong, T. Y., Grobler, A., Juonala, M., He, M., Dwyer, T., Burgner, D. & Wake, M. (2020). Associations of retinal microvascular caliber with large arterial function and structure: A population-based study of 11 to 12 year-olds and midlife adults. *Microcirculation*, 27 (6), <https://doi.org/10.1111/micc.12642>.

Persistent Link:

<https://hdl.handle.net/11343/275946>

MS. MENGJIAO LIU (Orcid ID : 0000-0002-8337-7633)

Article type : Original Research

Associations of retinal microvascular caliber with large arterial function and structure: A population-based study of 11-12 year-olds and mid-life adults

Running title: Co-variation of micro and macrovascular phenotypes

Authors:

Mengjiao Liu,^{1,2} Kate Lycett,^{1,2,3} Tien Yin Wong,^{1,4,5} Anneke Grobler,^{1,2} Markus Juonala,^{2,6,7} Mingguang He,^{1,4,8} Terry Dwyer,⁹ David Burgner,^{1,2,10,11} Melissa Wake.^{1,2,12}

Affiliations:

¹ The University of Melbourne, Melbourne, VIC, Australia

² Murdoch Children's Research Institute, Melbourne, VIC, Australia

³ Centre for Social & Early Emotional Development, Deakin University, Melbourne, VIC, Australia

⁴ Department of Ophthalmic Epidemiology, Centre for Eye Research Australia, The University of Melbourne, Melbourne, Australia

⁵ Singapore Eye Research Institute, Singapore National Eye Center, Singapore

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/MICC.12642](https://doi.org/10.1111/MICC.12642)

This article is protected by copyright. All rights reserved

⁶ Department of Medicine, University of Turku, Turku, Finland

⁷ Division of Medicine, Turku University Hospital, Turku, Finland

⁸ State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-Sen University, Guangzhou, China

⁹ The George Institute for Global Health, University of Oxford, Oxford, United Kingdom

¹⁰ Department of Paediatrics, Monash University, Melbourne, Australia

¹¹ Infectious Diseases, Royal Children's Hospital, Melbourne, Australia

¹² Department of Paediatrics & The Liggins Institute, The University of Auckland, Auckland, New Zealand

Grant numbers and source(s) of support: This work has been supported to date by the National Health and Medical Research Council of Australia (NHMRC; 1041352, 1109355), The Royal Children's Hospital Foundation (2014-241), Murdoch Children's Research Institute, The University of Melbourne, National Heart Foundation of Australia (100660) and Financial Markets Foundation for Children (2014-055; 2016-310). ML is supported by a Melbourne Research Scholarship. KL is supported by Australian National Health & Medical Research Council (NHMRC) Early Career Fellowship 1091124 and National Heart Foundation Postdoctoral Fellowship 101239. DB is supported by NHMRC Senior Research Fellowship 1064629 and is an Honorary Future Leader Fellowship of the National Heart Foundation of Australia (100369). MW was supported by NHMRC Senior Research Fellowship 1046518, Principal Research Fellowship 1160906 and Cure Kids New Zealand. Research at the Murdoch Children's Research Institute is supported by the Victorian Government's Operational Infrastructure Program. The funding bodies did not play any role in the study other than the generous provision of funds.

Corresponding author:

Prof Melissa Wake

E-mail: melissa.wake@mcri.edu.au

Tel: +61 3 9345 5761

This article is protected by copyright. All rights reserved

Author Manuscript

This article is protected by copyright. All rights reserved

1 **Abstract**

2 **Objective:** We examined associations between retinal microvascular and large arterial
3 phenotypes to explore relationships between the micro- and macro-vasculature in childhood
4 and mid-life.

5 **Methods:** Participants were 1288 children (11-12 years, 50.9% female) and 1264 adults
6 (mean age 44 years, 87.6% female) in a cross-sectional population-based study. Exposures
7 were retinal arteriolar and venular caliber quantified from retinal images. Outcomes included
8 arterial function (pulse wave velocity, PWV; carotid arterial elasticity) and structure (carotid
9 intima-media thickness, CIMT). Multivariable regression models were performed adjusting
10 for age, sex and family socioeconomic position.

11 **Results:** In children, one standard deviation (SD) wider arteriolar caliber was associated with
12 slower PWV (-0.09SD, 95%CI -0.16, -0.02) and higher elasticity (0.13SD, 95%CI 0.06,
13 0.20); per SD wider venular caliber was associated with faster PWV (0.06SD, 95%CI -0.01,
14 0.12) and lower elasticity (-0.08SD, 95%CI -0.14, -0.01). The size of adult associations was
15 approximately double. Wider arteriolar caliber was associated with smaller CIMT (-0.09SD,
16 95%CI -0.16, -0.03) in adults but not children. Venular caliber and CIMT showed little
17 evidence of association.

18 **Conclusions:** Narrower retinal arterioles and wider venules are associated with large arterial
19 function as early as mid-childhood. Associations strengthen by mid-life and also extend to
20 arterial structure, although effect sizes remain small.

21 **Keywords:** Retinal arteriolar caliber; retinal venular caliber; carotid intima-media thickness;
22 pulse wave velocity; arterial elasticity; population-based study.

1 **1. Introduction**

2 Identifying individuals at increased risk for cardiovascular disease (CVD) early in life could
3 help target interventions to reduce the global disease burden.^{1,2} Decades of epidemiological
4 and clinical evidence show that large artery measures consistently predict atherosclerosis
5 throughout the lifespan, and future CVD outcomes in adults.^{3,4} Commonly-used indicators of
6 poorer large artery cardiovascular phenotypes include faster pulse wave velocity (PWV),
7 reduced arterial elasticity and increased carotid intima-media thickness (CIMT).⁵⁻⁷

8 While generally receiving less public health attention, the microcirculation is also associated
9 with CVD.^{8,9} The retina is a unique site from which the microcirculation can be imaged
10 directly and non-invasively, with repeated measurement possible over time.¹⁰ Structural
11 vascular changes in the retina – in particular, arteriolar narrowing and venular dilation – have
12 been associated with systemic CVD, including coronary heart disease and heart failure.¹¹⁻¹³

13 The retinal blood vessels and large arteries are inherently interconnected elements of the
14 vascular tree and are influenced by the same intrinsic and environmental features with
15 systemic circulation.^{14,15} However, the extent to which microvascular and macrovascular
16 phenotypes co-develop, and if so from what age and from what state of health, are unclear.

17 We previously conducted a systematic review of 26 studies, including a meta-analysis of five
18 of these studies, to examine the association between retinal vascular caliber and intermediate
19 phenotypes of large arteries (ie, CIMT, PWV).¹⁶ Most studies involved adults in later life,
20 with only one study in children.¹⁷ Retinal arteriolar caliber showed small associations with
21 large arterial function (pooled estimates $r = -0.17$) and arterial structure ($r = -0.05$), with
22 stronger evidence in cardiometabolic patients than in the general population. Thus, large
23 arteries may share some pathophysiologic processes with the microvasculature, at least by
24 later life and in those already expressing disease endpoints.

25 However, there is a paucity of data from children and younger adults, and it remains unclear
26 when in the life course interrelationships between the small and large vascular phenotypes

1 emerge and how they change with age. If they already co-vary prior to emergent disease, then
2 this would suggest that the relationship is not solely the result of dysregulation, potentially
3 advancing understanding of CVD pathogenesis. With technology and artificial intelligence
4 putting retinal imaging into clinical and population research, microvascular status could
5 potentially enter precision prediction models for future CVD and other diseases.¹⁸ Before this
6 could take place, it is important to understand the small vessels' unique and shared
7 contributions with more traditional large artery measures. Besides, most previous studies
8 have only assessed one aspect of large arterial health (ie, functional *or* structural intermediate
9 phenotypes), but not both in the same individuals.¹⁶ Assuming that large arterial dysfunction
10 precedes structural changes, retinal microvascular associations with large artery function
11 might emerge earlier than with large artery structure.¹⁹

12 The national, population-based Longitudinal Study of Australian Children (LSAC) provided
13 an opportunity to address these evidence gaps in two generations taking part in a large cross-
14 sectional wave explicitly focusing on phenotypic health measurement. We aimed to examine
15 associations of retinal microvascular caliber with large arterial intermediate phenotypes
16 (measured by PWV, carotid arterial elasticity and CIMT) in Australian children aged 11-12
17 years and mid-life adults (the children's parents).

18 **2. Materials and methods**

19 **2.1 Study design and participants**

20 The Child Health CheckPoint (CheckPoint) is a cross-sectional population-derived wave
21 nested within the national LSAC study.²⁰ Details of the initial study design and recruitment
22 are outlined elsewhere.^{21, 22} Briefly, LSAC recruited a nationally representative birth cohort
23 (B cohort) of 5107 infants using a 2-stage clustered design and has since followed them in
24 biennial 'waves' of data collection up to 2016. Most of families retained to wave 6 in 2014
25 (73.7%, n=3764).

26 At the wave 6 visit, all participating families (n=3513) were invited to consent to their

1 contact details being shared with the CheckPoint team. CheckPoint took place between
2 February 2015 and March 2016, nested between LSAC waves 6 and 7 at child age 11-12
3 years. In total, 1874 children and their parents participated in this detailed cross-sectional
4 biophysical assessment.²³ Most attended the CheckPoint Main Assessment Centre in seven
5 large cities around Australia, where all measures were performed. A small minority of
6 families who attended more limited assessments in regional cities or home visits are not
7 included in the analyses as the retinal photograph was not collected at these visits, and home
8 visits did not include CIMT and arterial elasticity because the large and delicate equipment
9 could not be readily transported.

10 The CheckPoint study was approved by the Royal Children's Hospital Melbourne Human
11 Research Ethics Committee (33225D) and the Australian Institute of Family Studies Ethics
12 Committee (14-26). The attending parents/caregivers provided written informed consent for
13 themselves and their children to participate in the study.

14 **2.2 Retinal vascular caliber assessment**

15 Two optic disc-centered digital photographs from each eye were taken by a fundus camera
16 (EOS 60D SLR) in a darkened room. Details of scoring procedures are described elsewhere.²⁴
17 Briefly, right eye images were selected as the first choice for scoring. Each image was graded
18 by one of four experienced graders (two from the Zhongshan Ophthalmic Centre, China and
19 two from the Centre for Eye Research Australia, Australia) masked to participant age and sex
20 using the software program Integrative Vessel Analysis (IVAN, University of Wisconsin,
21 Madison, USA). Retinal vessels were identified as arterioles or venules from a specific area
22 (one-half to one-disc diameter from the margin of the optic disc). Each grader selected a
23 segment of each vessel within this area for measurement. Diameters of all the selected
24 segments were measured automatically by the IVAN software. Summary estimates of the
25 average retinal vascular caliber were calculated combining measurements of the six largest
26 arterioles or venules according to the revised Knudston-Parr formula.²⁵ Reproducibility of
27 measurement by IVAN was high, with inter- and intra-grader intraclass correlation

1 coefficients ranging of 0.76 to 0.99.²⁴

2 **2.3 Large arterial intermediate phenotypes**

3 Details of each measure are described elsewhere.^{26, 27} Carotid-femoral PWV measures arterial
4 function and was assessed using the Sphygmocor XCEL (AtCor Medical Pty Ltd., West
5 Ryde, Australia) with participants in the supine position. Briefly, the arterial waveforms of
6 the right carotid and femoral artery were measured by blood pressure cuffs on the arm and
7 leg. Data were collected one to three times for each participant and the mean was calculated.
8 PWV was calculated as the ratio of the distance traveled by the pulse wave and the time delay
9 between the waveforms, as expressed in meters per second (m/s). The agreement of PWV
10 waveforms between two analysts was high (0.99).²⁶ A quicker pulse wave indicates higher
11 levels of arterial stiffness.

12 CIMT measures large arterial structure, an indicator of subclinical atherosclerosis.²⁸ It was
13 assessed using standardized protocols via B-mode ultrasound (GE Healthcare Vivid-I BT06
14 with 10MHz linear array probe, Chicago, IL, USA).^{29, 30} To image the right carotid artery, the
15 participants lay in the supine position with the neck slightly extended and the head turned 45°
16 towards the left. Measurements were made at the distal 10mm of the right common carotid
17 artery, and images were taken across five sites within a minimum of three longitudinal loops
18 of five to ten cardiac cycles. CIMT is defined as the distance from the lumen-intima interface
19 to the media-adventitia interface. Images were analyzed using semi-automated edge-detection
20 software (Carotid Analyser, Medical Imaging Applications, Coralville, IA, USA) to calculate
21 CIMT far wall mean, where higher scores represent a thicker carotid artery wall. The inter-
22 and intra-grader intraclass correlation coefficients of CIMT analyses were moderate, ranging
23 of 0.59 to 0.71.²⁶

24 Carotid arterial elasticity (%/mmHg) measures the ability of the arteries to expand as a
25 response to pulse pressure caused by cardiac contraction and relaxation.³¹ We calculated
26 elasticity from carotid artery images using intima-intima lumen diameter (LD) and the
27 different value of systolic blood pressure and diastolic blood pressure (ΔP), according to

This article is protected by copyright. All rights reserved

1 previously published work from the Cardiovascular Risk in Young Finns Study:³²

$$\frac{\left(\frac{LD_{max} - LD_{min}}{LD_{min}}\right)}{\Delta P} \times 100\%$$

2
3 Lower arterial elasticity is considered to represent lower arterial compliance.

4 **2.4 Assessment of covariates**

5 We considered several covariates when examining associations between retinal
6 microvascular caliber and large artery phenotypes. The technical details of these measures are
7 summarized in Table 1. For example, age, sex and family socioeconomic position (SEP) were
8 considered as covariates, given that both large arterial and microvascular phenotypes have
9 been shown to vary by age (worse with age), have distinct sex differences, and to be worse in
10 those from poorer socioeconomic groups.^{12, 26} Body mass index (BMI, z-score for children),³³
11 low-density lipoprotein (LDL) cholesterol, smoking exposure (passive smoking for children
12 and ever smoked for adults) and early life risk factors for children (birth weight and
13 gestational age) were also included as covariates. These are commonly reported risk factors
14 for cardiovascular disease that have been reported to influence both large arterial and
15 microvascular variations.^{12, 34-36} Additionally, we considered mean arterial blood pressure as a
16 covariate as it might lie on the causal pathway linking microvascular variations to large
17 arterial changes.¹⁵

18 **2.5 Statistical analyses**

19 Linear regression assumptions were tested when fitting models to assess associations of
20 retinal vascular caliber with large arterial intermediate phenotypes in children and adults. We
21 considered retinal arteriolar or venular caliber as the independent variable and large arterial
22 intermediate phenotypes as the dependent variable,^{37, 38} although the reverse would be equally
23 plausible since they were measured cross-sectionally. It has been proposed that modeling
24 arteriolar and venular caliber simultaneously may give less biased results.³⁹ We thus
25 performed all analyses with both arteriolar and venular caliber in the same model.

This article is protected by copyright. All rights reserved

1 We fitted two main models and ran one sensitivity analysis. Model 1 estimates were adjusted
2 for age, sex, SEP and jointly for arteriolar or venular caliber. Model 2 estimates for children
3 additionally adjusted for BMI z-score, LDL cholesterol, passive smoking, birth weight and
4 gestational age; for adults, we further adjusted estimates for BMI, LDL cholesterol and ever
5 smoking. This fully adjusted model could inform whether the association is independent of
6 these common potential risk factors. We also conducted a sensitivity analysis further
7 adjusting estimates for mean arterial blood pressure to examine whether it could lie on the
8 causal pathway. For all analyses, we internally standardized exposures and outcomes (ie,
9 mean of zero and a standard deviation [SD] of 1) to present standardized regression
10 coefficients. All analyses were conducted in Stata 14.2 (StataCorp LP, TX, USA).

Author Manuscript

1 3. Results

2 3.1 Sample characteristics

3 Supplementary Figure presents the study flow from wave 1 of LSAC onwards. Of the 1874
4 parent-child dyads that took part in CheckPoint, 1288 children and 1264 adults had retinal
5 caliber data available. The CheckPoint sample mean family SEP (0.26) was above the mean
6 SEP of all families enrolled in LSAC at wave 1 (mean 0, SD 1), indicating more advantage in
7 participating families. Participants with retinal data available were similar to those without
8 these images, except for family SEP, which was around a quarter of a SD lower (ie, less
9 advantaged) in those missing retinal data (Table 2). In addition, children without retinal
10 images were slightly older (11.7 vs. 11.4 years of age) and more likely to be male (51% vs.
11 45%).

12 3.2 Main results

13 Table 3 shows the associations of retinal vascular caliber with large arterial phenotypes in
14 children and adults. In Model 1 (estimates adjusted for age, sex, SEP and jointly considering
15 arteriolar and venular caliber), both worse arteriolar and venular caliber were found to
16 consistently associate with adverse large arterial function. For example, in children, a one SD
17 higher arteriolar caliber was associated with slower PWV (-0.15SD, 95% CI -0.21 to -0.09)
18 and higher arterial elasticity (0.14SD, 95% CI 0.06 to 0.20), while a one SD higher venular
19 caliber was associated with faster PWV (0.09SD, 95% CI 0.03 to 0.15) and lower arterial
20 elasticity (-0.07SD, 95% CI -0.13 to -0.01). Similar patterns were mirrored in adults, but
21 associations were about twice the size seen in children (Figure).

22 In Model 2, when estimates were additionally adjusted for common risk factors (ie BMI, LDL
23 and smoking exposure), the associations of arteriolar and venular caliber with PWV were
24 attenuated in both children and adults by 30-40%. For example, in children per SD higher
25 arteriolar caliber was associated with slower PWV (-0.09SD, 95% CI -0.16 to -0.02), while
26 higher venular caliber was associated with faster PWV (0.06SD, 95% CI -0.01 to 0.12). The
27 associations with arteriolar elasticity changed little after the further adjustment in Model 2.
28 The model R^2 for the association of retinal vascular caliber and functional phenotypes was
29 about 6% to 13% in children and about 30% in adults.

30 Results were mixed for associations of retinal microvascular caliber and large arterial
31 structure. There was little evidence of an association between arteriolar caliber and CIMT in
32 children, while in adults, per SD higher arteriolar caliber was associated with smaller CIMT

1 scores (-0.09SD, 95% CI -0.16 to -0.03, Model 1), and this association attenuated by 44%
2 after common risk factor adjustment (-0.05SD, 95% CI -0.12 to 0.01, Model 2).
3 Unexpectedly, higher retinal venular caliber showed small associations with smaller CIMT in
4 children (-0.08SD, 95% CI -0.15 to -0.01), but there was little evidence of an association in
5 adults. The proportion of variance explained in model 2 was about 4% for children and 20%
6 for adults.

7 **3.3 Sensitivity analysis**

8 Further adjustment for mean arterial blood pressure attenuated, but did not eliminate,
9 associations of arteriolar and venular caliber with arterial elasticity in children and adults.
10 However, it fully attenuated associations with PWV (Table 3). Adding mean arterial blood
11 pressure to the models did not change the lack of adjusted associations with CIMT.

1 4. Discussion

2 *Principal findings*

3 In this study, we found that the microvasculature is associated with functional artery
4 phenotypes (ie, PWV, arterial elasticity) from early in life, and this association strengthens
5 with age. By midlife, there is evidence of an association between the microvasculature and
6 large artery structural phenotypes (ie, greater CIMT). It is likely that these associations will
7 increase with age and with the development of manifest cardiometabolic disease.¹⁶

8 *Strengths and limitations*

9 Our study has several strengths. It adds important new knowledge to the only other study (to
10 our knowledge) to consider the association of the microvascular and large arterial
11 intermediate phenotypes in generally healthy children.¹⁷ Our study drew on a large,
12 randomly-selected national sample of Australian children and their parents, spanning diverse
13 geographical areas across a very large country. All measures were objectively measured and
14 processed under standard operating procedures, with moderate to high inter- and intra-rater
15 reliability, and were conducted at the same time using the same equipment and processes for
16 the two generations. This cross-generational design enabled us to efficiently infer temporal
17 strengthening of associations with age, which would otherwise take decades to observe. Our
18 well-phenotyped sample allowed us to consider the effects of important covariates. Unlike
19 other studies, we were able to examine relationships of retinal microvascular caliber with
20 both arterial function and structure simultaneously in the same individuals, across the two
21 generations.

22 There are also limitations. Firstly, as our analytic sample was on average somewhat more
23 advantaged than the national average, our findings may not generalize well to the most
24 disadvantaged families (some of whom were nonetheless included). Secondly, limited data
25 were available for adult men as most adults in the CheckPoint sample were mothers, but
26 similar patterns of associations were nonetheless seen in adult women and men
27 (Supplementary Table). Thirdly, the cross-sectional nature of the study precludes inference
28 regarding the temporal sequence of the associations.

29 *Interpretation in light of other studies*

30 To date, only one study has examined similar associations in healthy children.¹⁷ The
31 Generation R Study found that in children (median age six years) each SD increase in retinal
32 arteriolar and venular caliber was associated with 0.02SD (95% CI -0.01 to 0.05) and 0.04SD
This article is protected by copyright. All rights reserved

1 (95% CI 0.01 to 0.07) higher carotid-femoral PWV (SD 0.9 cm/s).¹⁷ In our study with
2 children aged 11-12 years, we found a much stronger association with PWV and replicated
3 this finding with another arterial function measure, arterial elasticity. In turn, we now show
4 that the magnitude of associations between retinal vascular caliber and arterial functional
5 phenotypes roughly doubles between mid-childhood and mid-adulthood. The explanatory
6 power (ie, R^2 values) was small at age 11-12 years and much stronger in adults. These results
7 are consistent with our meta-analysis of slightly older adults (mean age 54 years), which
8 showed narrower retinal arteriolar caliber was associated with faster PWV ($r = -0.17$, 95%CI
9 -0.25 to -0.10).¹⁶

10 We found little evidence that retinal arteriolar caliber was associated with CIMT in children,
11 but a small association was observed in adults. In children of similar age (mean age 13.6
12 years) with type 1 diabetes, retinal vascular caliber has been associated with CIMT.⁴⁰ In our
13 meta-analysis with older adults (mean age 63 years), narrower retinal arteriolar caliber was
14 weakly associated with higher CIMT scores ($r = -0.05$, 95%CI -0.09 to -0.02),¹⁶ and effect
15 sizes were greater in patients with cardiometabolic disease. Taken together, both age and
16 cardiometabolic disease conditions may contribute to the degree of co-variation between
17 small and large artery structure.

18 Higher CIMT is associated with adverse changes in arterial structure, which has been shown
19 to be linked with cardiovascular risk factors such as obesity and hypertension in children.⁴¹
20 The association we observed between wider (worse) venular caliber and lower (better) CIMT
21 in children may be a chance finding, given its very small magnitude, unexpected direction
22 and full attenuation on adjustment and in the sensitivity analysis.

23 *Implications*

24 The potential association between the macro- or microcirculation and CVD risk are usually
25 investigated separately. While it is plausible that their genetic determinants could be quite
26 separate, biologically the two vascular systems are functionally related to the pulsatile
27 transmission of pressure and flow.¹⁵ We have provided population-based evidence that the
28 phenotypes of macro- and microcirculation co-vary from mid-childhood, and other studies
29 suggest that this relationship is emerging from as young as six years onwards.¹⁷ The strength
30 of the association increases with both age and with manifest cardiometabolic disease.¹⁶

31 Some of the observed associations between the macro- and microcirculation may arise from
32 shared cardiometabolic risk factors.⁴² For example, we found that cardiovascular risk factors

1 such as BMI, LDL cholesterol, and smoking exposures attenuated the association of retinal
2 vascular caliber with functional and structural large artery phenotypes to a varying extent.
3 However, these factors only contributed to a small amount of co-variations between small
4 and large vascular phenotypes. It may be that small and large vessels are differentially
5 susceptible to risk factors,⁸ or that they are mainly determined by distinct factors. In addition,
6 the fully attenuated relationships between retinal caliber and PWV after adjusting for mean
7 arterial blood pressure suggests the association may be mediated via blood pressure. This
8 further emphasizes the likely importance of public health interventions to reduce blood
9 pressure throughout life in order to protect all elements of the vascular tree.

10 There is strong evidence that microvascular dysfunction plays an important role in the
11 pathogenesis of CVD.⁹ The intermediate phenotype of large arteries represents a snapshot of
12 preclinical cardiovascular health. In the current study, we found that phenotypic measures of
13 microcirculation were associated with large arterial phenotypes, which in keeping with
14 experiment models showing that abnormality of small and large vessels often coexist in the
15 pathogenesis of CVD.⁴³ Given the association between microvascular and large arterial
16 phenotypes was relatively small, each phenotype may contribute to CVD through largely
17 separate causal pathways. Further studies with clinical CVD outcomes would determine
18 whether combining phenotypes of the microvasculature and large arteries improves risk
19 prediction and inform the optimal use of these measures.

20 *Conclusion*

21 Retinal microvascular caliber is associated with large arterial function by mid-childhood,
22 with stronger (though still small) associations observed in mid-life. Cumulative exposure to
23 shared risk factors may underlie the age-dependent co-variation between small and large
24 arterial phenotypes. Longitudinal studies with clinical outcomes would inform the optimal
25 use of these parameters in early cardiovascular risk management.

27 **Perspectives**

28 Variations in the micro- and macro-vasculature underlie CVD pathogenesis, but how they
29 may influence each other at different life stages is not well understood. We showed a small
30 amount of co-variation between the two in childhood, which doubled in strength by mid-life.
31 This suggests it is important to understand the unique contribution of both the micro- and
32 macro-vasculature to CVD across the life course, and whether the assessment of both the

1 small and large vessels could improve risk prediction for CVD.

Author Manuscript

Conflicts of interest

The authors report no relationships that could be construed as a conflict of interest.

Acknowledgments

This study uses data from the Longitudinal Study of Australian Children (LSAC) and Child Health CheckPoint. We thank the LSAC and CheckPoint study participants and families. We also thank the CheckPoint team and the Murdoch Children's Research Institute. LSAC is conducted in partnership between the Department of Social Services (DSS), the Australian Institute of Family Studies (AIFS) and the Australian Bureau of Statistics (ABS). The findings and views reported in this paper are those of the authors and should not be attributed to DSS, AIFS or the ABS.

Author contributions

MW conceived and led the CheckPoint study with the CheckPoint team, and was LSAC's Health Design Leader. KL was the primary student supervisor, along with MW, DB, TW, MH, and oversaw all aspects of the study and the manuscript preparation. ML performed data analysis and wrote the main paper. All authors provided expert advice and critical review of this manuscript, and had final approval of the submitted and published version of this paper.

References

1. Zieske AW, Malcom GT, Strong JP. Natural history and risk factors of atherosclerosis in children and youth: the PDAY study. *Pediatr Pathol Mol Med*. 2002;21:213-37.
2. World Health Organization (WHO), Prevention of cardiovascular disease, in *Guideline for assessment and management of cardiovascular risk*, WHO; 2007.
3. Vlachopoulos C, Aznaouridis K, Stefanadis C. Prediction of cardiovascular events and all-cause mortality with arterial stiffness: a systematic review and meta-analysis. *J Am Coll Cardiol*. 2010;55:1318-27.
4. Seidemann SB, Claggett B, Bravo PE, Gupta A, Farhad H, Klein BE, Klein R, Di Carli MSolomon SD. Retinal vessel calibers in predicting long-term cardiovascular outcomes: The Atherosclerosis Risk in Communities Study. *Circulation*. 2016;134:1328-1338.
5. Ohkuma T, Ninomiya T, Tomiyama H, Kario K, Hoshide S, Kita Y, Inoguchi T, Maeda Y, Kohara K, Tabara Y, Nakamura M, Ohkubo T, Watada H, Munakata M, Ohishi M, Ito N, Nakamura M, Shoji T, Vlachopoulos C, Yamashina A Collaborative Group for JB. Brachial-ankle pulse wave velocity and the risk prediction of cardiovascular disease: an individual participant data meta-analysis. *Hypertension*. 2017;69:1045-1052.
6. Boesen ME, Singh D, Menon BK Frayne R. A systematic literature review of the effect of carotid atherosclerosis on local vessel stiffness and elasticity. *Atherosclerosis*. 2015;243:211-22.
7. Carpenter M, Sinclair H, Kunadian V. Carotid intima media thickness and its utility as a predictor of cardiovascular disease: A review of evidence. *Cardiol Rev*. 2016;24:70-5.
8. Bosetti F, Galis ZS, Bynoe MS, Charette M, Cipolla MJ, Del Zoppo GJ, Gould D, Hatsukami TS, Jones TL, Koenig JI, Luty GA, Maric-Bilkan C, Stevens T, Tolunay HE, Koroshetz W. Small Blood Vessels: Big Health Problems" Workshop P. "Small Blood Vessels: Big Health Problems?": Scientific Recommendations of the National Institutes of Health Workshop. *J Am Heart Assoc*. 2016;5.
9. Musa MG, Torrens C, Clough GF. The microvasculature: a target for nutritional programming and later risk of cardio-metabolic disease. *Acta Physiol (Oxf)*. 2014;210:31-45.
10. Kipli K, Hoque ME, Lim LT, Mahmood MH, Sahari SK, Sapawi R, Rajae N, Joseph

-
- A. A review on the extraction of quantitative retinal microvascular image feature. *Comput Math Methods Med.* 2018;2018:4019538.
11. Li LJ, Ikram MK, Wong TY. Retinal vascular imaging in early life: insights into processes and risk of cardiovascular disease. *J Physiol.* 2016;594:2175-203.
 12. Ikram MK, Ong YT, Cheung CY, Wong TY. Retinal vascular caliber measurements: clinical significance, current knowledge and future perspectives. *Ophthalmologica.* 2013;229:125-36.
 13. Grassi G, Mancia G. 'Keep an eye' on the heart: retinal microcirculation disarray in congestive heart failure. *Eur Heart J.* 2018;39:57-59.
 14. Flammer J, Konieczka K, Bruno RM, Virdis A, Flammer AJ, Taddei S. The eye and the heart. *Eur Heart J.* 2013;34:1270-8.
 15. Safar ME, Struijker-Boudier HA. Cross-talk between macro- and microcirculation. *Acta Physiol (Oxf).* 2010;198:417-30.
 16. Liu M, Wake M, Wong TY, He M, Xiao Y, Burgner D, Lycett K. Associations of retinal microvascular caliber with intermediate phenotypes of large arterial function and structure: A systematic review and meta-analysis. *Microcirculation.* 2019:e12557.
 17. Gishti O, Jaddoe VW, Felix JF, Klaver CC, Hofman A, Wong TY, Ikram MK, Gaillard R. Retinal microvasculature and cardiovascular health in childhood. *Pediatrics.* 2015;135:678-85.
 18. Schmidt-Erfurth U, Sadeghipour A, Gerendas BS, Waldstein SM, Bogunovic H. Artificial intelligence in retina. *Prog Retin Eye Res.* 2018;67:1-29.
 19. Balletshofer BM, Rittig K, Stock J, Lehn-Stefan A, Overkamp D, Dietz K, Haring HU. Insulin resistant young subjects at risk of accelerated atherosclerosis exhibit a marked reduction in peripheral endothelial function early in life but not differences in intima-media thickness. *Atherosclerosis.* 2003;171:303-9.
 20. Wake M, Clifford S, York E, Mensah F, Burgner D, Davies S. Introducing growing up in Australia's child health check point: A physical health and biomarkers module for the longitudinal study of Australian children. *Family Matters.* 2014:15.
 21. Edwards B. Growing Up in Australia: The Longitudinal Study of Australian children entering adolescence and becoming a young adults. *Family Matters.* 2014:5-14.
 22. Sanson A, Robert J. Growing Up in Australia takes its first steps. *Family Matters.* 2004:46-53.
 23. Clifford S, Davies S, Gillespie A, Lange K, Wake M, Longitudinal Study of

Australian Children's Child Health CheckPoint Data User Guide - December 2018, Murdoch Children's Research Institute: Melbourne; 2018.

24. Dascalu J, Liu M, Lycett K, Grobler AC, He M, Burgner DP, Wong TY, Wake M. Retinal microvasculature: population epidemiology, concordance in Australian children aged 11–12 years and their parents. *BMJ Open*. 2019;9:44-52.
25. Knudtson MD LK, Hubbard LD, Wong TY, Klein R, Klein BEK. Revised formulas for summarizing retinal vessel diameters. *Curr Eye Res*. 2003;27:143-149.
26. Liu RS, Dunn S, Grobler AC, Lange K, Becker D, Goldsmith G, Carlin JB, Juonala M, Wake M, Burgner DP. Carotid artery intima–media thickness, distensibility and elasticity: population epidemiology and concordance in Australian children aged 11–12 years old and their parents. *BMJ Open*. 2019;9:23-33.
27. Clifford SA, Davies S, Wake M. Child Health CheckPoint Team. Child Health CheckPoint: cohort summary and methodology of a physical health and biospecimen module for the Longitudinal Study of Australian Children. *BMJ Open*. 2019;9:3-22.
28. Iwakiri T, Yano Y, Sato Y, Hatakeyama K, Marutsuka K, Fujimoto S, Kitamura K, Kario K, Asada Y. Usefulness of carotid intima-media thickness measurement as an indicator of generalized atherosclerosis: findings from autopsy analysis. *Atherosclerosis*. 2012;225:359-62.
29. Nezu T, Hosomi N, Aoki S, Matsumoto M. Carotid intima-media thickness for atherosclerosis. *J Atheroscler Thromb*. 2016;23:18-31.
30. Touboul PJ, Hennerici MG, Meairs S, Adams H, Amarenco P, Bornstein N, Csiba L, Desvarieux M, Ebrahim S, Hernandez Hernandez R, Jaff M, Kownator S, Naqvi T, Prati P, Rundek T, Sitzer M, Schminke U, Tardif JC, Taylor A, Vicaut E, Woo KS. Mannheim carotid intima-media thickness and plaque consensus (2004-2006-2011). An update on behalf of the advisory board of the 3rd, 4th and 5th watching the risk symposia, at the 13th, 15th and 20th European Stroke Conferences, Mannheim, Germany, 2004, Brussels, Belgium, 2006, and Hamburg, Germany, 2011. *Cerebrovasc Dis*. 2012;34:290-6.
31. Zuo G, Zhang M, Jia X, Zheng L, Li Y, Zhao H, Wang C, Liang C, Du X. Correlation between brachial-ankle pulse wave velocity, carotid artery intima-media thickness, ankle-brachial index, and the severity of coronary lesions. *Cell Biochem Biophys*. 2014;70:1205-11.
32. Juonala M, Jarvisalo MJ, Maki-Torkko N, Kahonen M, Viikari JS, Raitakari OT. Risk factors identified in childhood and decreased carotid artery elasticity in adulthood: the

-
- Cardiovascular Risk in Young Finns Study. *Circulation*. 2005;112:1486-93.
33. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, Flegal KM, Guo SS, Wei R, Mei Z, Curtin LR, Roche AF, Johnson CL. CDC growth charts: United States. *Adv Data*. 2000;314:1-27.
34. Cote AT, Phillips AA, Harris KC, Sandor GG, Panagiotopoulos C, Devlin AM. Obesity and arterial stiffness in children: systematic review and meta-analysis. *Arterioscler Thromb Vasc Biol*. 2015;35:1038-44.
35. Santos IS, Alencar AP, Rundek T, Goulart AC, Barreto SM, Pereira AC, Bensenor IM, Lotufo PA. Low impact of traditional risk factors on carotid intima-media thickness: The ELSA-Brasil Cohort. *Arterioscler Thromb Vasc Biol*. 2015;35:2054-9.
36. Gopinath B, Baur LA, Wang JJ, Teber E, Liew G, Cheung N, Wong TY, Mitchell P. Smaller birth size is associated with narrower retinal arterioles in early adolescence. *Microcirculation*. 2010;17:660-8.
37. Garcia-Ortiz L, Recio-Rodriguez JI, Agudo-Conde C, Patino-Alonso MC, Rodriguez-Sanchez E, Maderuelo-Fernandez JA, Gomez-Marcos MA. The role of retinal vessels caliber as a marker of vascular aging in large arteries. *J Hypertens*. 2015;33:818-26.
38. Triantafyllou A, Anyfanti P, Gavriilaki E, Zabulis X, Gkaliagkousi E, Petidis K, Triantafyllou G, Gkollas V, Pырpasopoulou A, Douma S. Association between retinal vessel caliber and arterial stiffness in a population comprised of normotensive to early-stage hypertensive individuals. *Am J Hypertens*. 2014;27:1472-8.
39. Liew G, Sharrett AR, Kronmal R, Klein R, Wong TY, Mitchell P, Kifley A, Wang JJ. Measurement of retinal vascular caliber: issues and alternatives to using the arteriole to venule ratio. *Invest Ophthalmol Vis Sci*. 2007;48:52-7.
40. Pena AS, Liew G, Anderson J, Giles LC, Gent R, Wong TY, Couper JJ. Early atherosclerosis is associated with retinal microvascular changes in adolescents with type 1 diabetes. *Pediatr Diabetes*. 2018;19:1467-1470.
41. Urbina EM, Williams RV, Alpert BS, Collins RT, Daniels SR, Hayman L, Jacobson M, Mahoney L, Mietus-Snyder M, Rocchini A, Steinberger J, McCrindle B, American Heart Association Atherosclerosis, Hypertension, Obesity in Youth Committee of the Council on Cardiovascular Disease in the Young. Noninvasive assessment of subclinical atherosclerosis in children and adolescents: recommendations for standard assessment for clinical research: a scientific statement from the American Heart Association. *Hypertension*. 2009;54:919-50.
42. Nguyen TT, Islam FM, Farouque HM, Klein R, Klein BE, Cotch MF, Herrington

DMWong TY. Retinal vascular caliber and brachial flow-mediated dilation: the Multi-Ethnic Study of Atherosclerosis. *Stroke*. 2010;41:1343-8.

43. Chantler PD. and Frisbee JC. Arterial function in cardio-metabolic diseases: from the microcirculation to the large conduits. *Prog Cardiovasc Dis*. 2015;57:489-96.

Author Manuscript

Legends

TABLES

Table 1. Measures of covariates

Table 2. Sample characteristics

Table 3. Associations of retinal microvascular caliber with large arterial intermediate phenotypes

FIGURE: Associations of retinal microvascular caliber and large arterial intermediate phenotypes. Models included arteriolar and venular caliber simultaneously, adjusting for age, sex and socioeconomic position (Model 1)

Footnotes: Effect size = standardized regression coefficients β , representing per standard deviation increase in arteriolar/venular caliber for a standard deviation change in large arterial phenotypes. Abbreviations: PWV, pulse wave velocity; CIMT, carotid intima-media thickness

Author Manuscript

Table 1. Measures of covariates

Measure	Equipment	Brief protocol
<i>Model 1</i>		
Age	Parental questionnaire	Age was calculated to nearest week using date of birth, either imported from Medicare Australia's database at the time of LSAC enrolment (child) or self-reported (parent), and date of assessment.
Sex	Parental questionnaire	Sex was self-reported in the CheckPoint questionnaire by children and parents respectively.
Family socioeconomic position (SEP)	Parental questionnaire	SEP was measured in LSAC wave 6 (child age 10-11), approximately 12 months prior to measurement of biophysical measures in CheckPoint. This SEP variable summarized parent-reported combined household income, current or most recent occupation of each parent and highest achieved educational qualification of each parent. Each component of the score was scaled and an unweighted average was calculated over three values in a single parent household, or over five values in a dual parent household. The unweighted average variable at LSAC was then standardized within the wave to have a mean 0, and standard deviation (SD) of 1.
<i>Model 2</i>		
Body mass index (BMI)	Height: Invicta Stadiometer Weight: 4-limb segmental body composition scales	Height (cm): Participants were measured in bare feet. Mean of two measurements used. If two measures differ by >0.5cm, a third was taken and mean of all measurements was used. Weight (kg): Bare feet and in light clothing, participants' weight were measured to the nearest 50g. BMI was calculated as kg/m ² . For children, an age- and sex-adjusted BMI z-score was calculated using the US Centres for Disease Control and Prevention growth reference charts.[33]
Low-density lipoprotein cholesterol	Proton nuclear magnetic resonance (NMR)	Venous blood samples of 30 ml were collected from each participant at the main and mini assessment centers. Participants were at a minimum semi-fasted (ie, no food/drink for at least 2 hours). A high-throughput NMR metabolomics platform was used to quantify the lipid and metabolite measures from stored EDTA plasma samples. The platform applies a single experimental setup, which allows for the simultaneous quantification of routine lipids

Measure	Equipment	Brief protocol
Smoking exposure	Parental questionnaire	At each LSAC wave, parents reported the number of smokers at home. We created a binary variable of “ever exposed to passive smoke” from this data; it was positive for the child if the answer to the smokers at home question in any LSAC wave (age 0 to 11) was greater than zero.
Birth weight	Parental questionnaire	At LSAC Wave 1, the parent was asked “How much did the child weigh at birth?”
Gestational age	Parent Health Record Parental questionnaire	At LSAC Wave 1, the interviewer copied gestational age from the Parent Health Record recorded prospectively by the birthing hospital (the great majority). If the record was not available, interviewers asked parents the question “After how many weeks of pregnancy was [child’s name] born?”
<i>Sensitivity analysis</i>		
Mean arterial blood pressure	SphygmoCor XCEL	Following seven minutes in the supine position at rest, systolic and diastolic blood pressures were measured at the brachial artery up to three times, with mean values reported. Mean arterial blood pressure was calculated as 1/3 systolic and 2/3 diastolic blood pressure.

Table 2. Sample characteristics

Characteristics ^a	Children				Adults			
	With data on retinal image		Without data on retinal image		With data on retinal image		Without data on retinal image	
	N	Mean (SD ^b) or %	N	Mean (SD ^b) or %	N	Mean (SD ^b) or %	N	Mean (SD ^b) or %
Age (years)	1288	11.4 (0.5)	586	11.7 (0.5)	1264	43.8 (5.1)	610	43.6 (5.6)
Gender (% female)	1288	50.9	586	45.1	1264	87.6	610	90.0
Birth weight (kg)	1285	3.44 (0.55)	585	3.43 (0.62)				
Gestational age (weeks)	1279	39.2 (2.0)	586	38.9 (4.0)				
Socioeconomic position	1283	0.26 (0.97)	584	-0.02 (1.01)	1260	0.26 (0.97)	607	-0.01 (1.01)
Smoking exposure	1287	20.7	586	27.8	1263	8.4	610	11.2
Body mass index z-score ^c	1288	0.30 (0.96)	584	0.31 (1.04)				
Body mass index (kg/m ²)					1259	27.86 (6.15)	602	27.74 (6.11)
Mean arterial blood pressure (mmHg)	1213	78.07 (5.66)	564	76.65 (6.03)	1169	89.05 (9.70)	580	87.20 (8.98)
Retinal arteriolar calibre (µm)	1288	159.1 (11.9)			1264	151.0 (14.0)		
Retinal venular calibre (µm)	1288	230.7 (16.6)			1264	218.9 (18.5)		
Pulse wave velocity (m/s)	1242	4.44 (0.54)	564	4.49 (0.63)	1134	6.96 (1.14)	541	6.66 (1.08)
Arterial elasticity (%/10mmHg)	1151	4.83 (0.87)	193	4.66 (1.00)	1053	2.44 (0.06)	199	2.50 (0.66)
Carotid intima-media thickness (µm)	1273	580.03 (46.40)	212	589.01 (47.59)	1236	663.55 (97.01)	232	667.04 (99.37)

^a. Most characteristics measured at the Child Health CheckPoint; birth weight, gestation were collected at LSAC wave 1 in 2004, SEP was measured at LSAC Wave 6 in 2015, and smoking exposure was generated combing the questionnaire from LSAC Wave 1 to 6; ^b. Standard deviation; ^c. Body mass index was transformed to z-score with widely used Centres for Disease Control and Prevention (US)-growth charts.


Table 3. Associations of retinal microvascular caliber with large arterial intermediate phenotypes

Retinal caliber (per SD increase)	Pulse wave velocity			Carotid arterial elasticity			Carotid intima-media thickness		
	Standardized β (95% CI)	p-value	R ²	Standardized β (95% CI)	p-value	R ²	Standardized β (95% CI)	p-value	R ²
Children									
Model 1: adjusted for age, sex, socioeconomic position and jointly considering arteriolar and venular caliber									
Arteriolar	-0.15 (-0.21, -0.09)	<0.001	7.3	0.13 (0.06, 0.20)	<0.001	3.3	-0.03 (-0.10, 0.03)	0.30	2.2
Venular	0.09 (0.03, 0.15)	<0.01		-0.07 (-0.13, -0.01)	0.03		-0.07 (-0.13, -0.01)	0.03	
Model 2: Model 1 plus BMI z-score, low-density lipoprotein cholesterol, passive smoking, birth weight and gestational age									
Arteriolar	-0.09 (-0.16, -0.02)	<0.01	12.8	0.13 (0.06, 0.20)	<0.01	6.1	-0.02 (-0.09, 0.05)	0.51	3.6
Venular	0.06 (-0.01, 0.12)	0.10		-0.08 (-0.14, -0.01)	0.03		-0.08 (-0.15, -0.01)	0.02	
Sensitivity analysis: Model 2 plus mean arterial blood pressure									
Arteriolar	-0.02 (-0.09, 0.04)	0.51	20.1	0.08 (0.00, 0.15)	0.04	10.0	-0.03 (-0.10, 0.04)	0.42	3.3
Venular	0.02 (-0.04, 0.08)	0.47		-0.05 (-0.12, 0.02)	0.15		-0.07 (-0.14, 0.00)	0.06	
Adults									
Model 1: adjusted for age, sex, socioeconomic position and jointly considering arteriolar and venular caliber									
Arteriolar	-0.31 (-0.37, -0.24)	<0.001	16.3	0.31 (0.24, 0.37)	<0.001	18.0	-0.09 (-0.16, -0.03)	<0.01	16.8
Venular	0.12 (0.05, 0.18)	<0.001		-0.14 (-0.21, -0.08)	<0.001		0.05 (-0.01, 0.11)	0.12	
Model 2: Model 1 plus BMI, low-density lipoprotein cholesterol and ever smoking									
Arteriolar	-0.20 (-0.27, -0.14)	<0.001	26.7	0.24 (0.17, 0.30)	<0.001	29.5	-0.05 (-0.12, 0.01)	0.12	20.9
Venular	0.06 (-0.00, 0.13)	0.06		-0.10 (-0.16, -0.04)	<0.01		0.05 (-0.02, 0.11)	0.17	
Sensitivity analysis: Model 2 plus mean arterial blood pressure									
Arteriolar	0.01 (-0.06, 0.07)	0.82	43.7	0.12 (0.06, 0.19)	<0.001	36.0	-0.01 (-0.08, 0.06)	0.72	21.9
Venular	-0.02 (-0.08, 0.04)	0.57		-0.06 (-0.12, -0.00)	0.05		0.01 (-0.05, 0.08)	0.72	

All models included arteriolar and venular caliber, plus other covariates. SDs of arteriolar and venular caliber for children were 11.9 and 16.6 μm , and for adults

were 14.0 and 18.5 μm , respectively. Abbreviations: SD, standard deviation; CI, confidence interval; BMI, body mass index; R^2 , model R square.

Author Manuscript



micc_12642_f1.tif