



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Biswas, T;Azzopardi, P;Anwar, SN;de Vries, TD;Encarnacion-Cruz, LM;Hasan, MM;Huda, MM;Pervin, S;Das Gupta, R;Mitra, DK;Rawal, LB;Al Mamun, A

Title:

Assuring Bangladesh's future: non-communicable disease risk factors among the adolescents and the existing policy responses.

Date:

2022-05-16

Citation:

Biswas, T., Azzopardi, P., Anwar, S. N., de Vries, T. D., Encarnacion-Cruz, L. M., Hasan, M. M., Huda, M. M., Pervin, S., Das Gupta, R., Mitra, D. K., Rawal, L. B. & Al Mamun, A. (2022). Assuring Bangladesh's future: non-communicable disease risk factors among the adolescents and the existing policy responses.. *J Health Popul Nutr*, 41 (1), pp.22-. <https://doi.org/10.1186/s41043-022-00294-x>.

Persistent Link:

<https://hdl.handle.net/11343/316253>

License:

[CC BY](#)

RESEARCH ARTICLE

Open Access



Assuring Bangladesh's future: non-communicable disease risk factors among the adolescents and the existing policy responses

Tuhin Biswas^{1,2*}, Peter Azzopardi³, Syeda Novera Anwar⁴, Tim David de Vries⁵, Luis Manuel Encarnacion-Cruz⁶, Md. Mehedi Hasan^{1,2}, M. Mamun Huda^{1,2}, Sonia Pervin^{1,2,7}, Rajat Das Gupta⁸, Dipak Kumar Mitra⁹, Lal B. Rawal¹⁰ and Abdullah Al Mamun^{1,2}

Abstract

Background: The aim of this study is to assess the current status of non-communicable disease (NCD) risk factors amongst adolescents in Bangladesh. We also critically reviewed the existing policy responses to NCD risk among adolescents in Bangladesh.

Methods: This study used a mixed method approach. To quantify the NCD risk burden, we used data from the Global School-based Student Health Survey conducted in Bangladesh. To understand policy response, we reviewed NCD-related policy documents introduced by the Government of Bangladesh between 1971 and 2018 using the WHO recommended NCD Action Plan 2013–2020 as study framework. Information from the policy documents was extracted using a matrix, mapping each document against the six objectives of the WHO 2013–2020 Action Plan.

Results: Almost all adolescents in Bangladesh had at least one NCD risk factor, and there was a high prevalence of concurrent multiple NCD risk factors; 14% had one NCD risk factor while 22% had two, 29% had three, 34% had four or more NCD risk factors. Out of 38 policy documents, eight (21.1%) were related to research and/or surveys, eight (21.1%) were on established policies, and eleven (29%) were on legislation acts. Three policy documents (7.9%) were related to NCD guidelines and eight (21.1%) were strategic planning which were introduced by the government and non-government agencies/institutes in Bangladesh.

Conclusions: The findings emphasize the needs for strengthening NCD risk factors surveillance and introducing appropriate intervention strategies targeted to adolescents. Despite the Government of Bangladesh introducing several NCD-related policies and programs, the government also needs more focus on clear planning, implementation and monitoring and evaluation approaches to preventing NCD risk factors among the adolescents in Bangladesh.

Keywords: Non-communicable disease, Adolescents, Health policy, Bangladesh

Introduction

Adolescents represent almost a quarter of the world's population [1], and adolescence is now recognised as a pivotal developmental stage where investments can result in healthier young people, healthier adults and a healthier next generation [2, 3]. Nearly 35% of the global burden

*Correspondence: t.biswas@uqconnect.edu.au

¹ Institute for Social Science Research, The University of Queensland, QLD, Australia

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

of disease has its origins in adolescence, and more than 3000 adolescents die every day, mostly due to non-communicable diseases (NCDs), intentional and unintentional injuries and other preventable causes [4, 5].

NCD-related deaths are increasing, especially in low- and middle-income countries (LMICs) [6] and over half of these deaths are associated with risk that emerge during the adolescence. These include, but not limited to tobacco and alcohol use, poor diet, and insufficient physical activity [6]. According to the Lancet Commission on Adolescent Health (2016), tobacco use, alcohol consumption, overweight, obesity, and mental health problems were identified as the major health risks for adolescents around the world [7, 9]. Early tobacco use is a major risk factor for NCDs throughout the life. In addition, smoking at a young age also increases the risk of many diseases among adolescents such as respiratory illness, asthma, and reduced pulmonary function [6]. Alcohol consumption is linked to development of different types of cancers, hypertension, hemorrhagic stroke and other NCDs [8, 10–12]. Numerous studies have reported that overweight and obesity are increasing markedly across adolescence and young adulthood [13]. To combat the burden of NCDs worldwide, the Sustainable Development Goals (SDGs) include a specific target for reducing premature death from NCDs by one-third, through prevention and treatment and promotion of mental health and well-being by 2030. However, the importance of NCD prevention among adolescents in global NCD declarations and action plans is yet to be fully realised.

Like many LMICs, Bangladesh is undergoing rapid urbanization, which has triggered changes in population dynamics and their disease patterns. The country is at an advanced phase of the third stage of the epidemiological transition, which means that deaths from NCDs are expected to increase rapidly in the coming years [14, 15]. Adolescents account for one third of the country's total population. A recent study in Bangladesh reported that 18% of the adolescents had three or more risk factors, with males reporting higher prevalence than females [4, 5]. Another study by Urmy et al. reported that younger age, non-slum urban and slum residence, higher paternal education, and depression were associated with the coexistence of multiple risk factors among the adolescent [6]. However, no study has been conducted in Bangladesh which provides comprehensive details on NCD risk factors among adolescents and its policy response.

Bangladesh is categorized as a 'multi-burden country'—i.e., high rate of Communicable/maternal/nutritional, high rate of injury, and a high rate of NCDs for adolescents [7, 9]. The government of Bangladesh has duly recognized this complex profile of need and has formulated a number of policy documents and strategies

to address the NCD burden and associated risk factors. However, the evidence of the effective implementation of these policies is limited. A comprehensive analysis of the existing policies is warranted to ascertain the adequacy of national strategies to tackle the adolescent NCDs,

In order to determine the current status of NCD risk factors among the adolescents in Bangladesh, we analyzed the Global School-based Health Survey (GSHS). We also examined the existing national policies and strategies through the lens of the World Health Organization's 2013–2020 Action Plan for the Global Strategy for the Prevention and Control of NCDs. This policy analysis determined the extent to which the objectives of the WHO Action Plan have been met at national policy level.

Methods

Data source

Quantitative data

The quantitative data presented in this study come from the Global School-based Health Survey (GSHS), which was conducted for the first time in Bangladesh in 2014 [16]. The 2014 Bangladesh GSHS was administered among adolescents aged 12–17 years to capture information on a wide range of health indicators. The 2014 Bangladesh GSHS employed a two-stage cluster sampling technique. At the first stage, the schools were selected randomly from a list of schools. Classes that provide a representative sample of the general population aged 12–17 years were selected within the selected schools at the second stage of sampling.

Policy documents

We also reviewed different policy documents on adolescent health issue. We considered if these policy documents were aligned directly or indirectly with the prevention and control of NCD risk factors among the adolescents in Bangladesh. Different search engines such as PubMed and Google Scholar were used to identify relevant documents. The key terms used in the website searches were 'adolescent health,' 'health education,' 'mental health,' 'nutrition for adolescents,' 'violence against adolescents,' combined with 'policy,' 'action plan,' 'strategy,' 'guideline and "Bangladesh"'. The search strategy employed with PubMed is as follows:

```
(((("adolescent health"[MeSH Terms] OR
("adolescent"[All Fields] AND "health"[All Fields])
OR "adolescent health"[All Fields]) OR ("mental
health"[MeSH Terms] OR ("mental"[All Fields]
AND "health"[All Fields]) OR "mental health"[All
Fields])) OR (("nutritional status"[MeSH Terms]
OR ("nutritional"[All Fields] AND "status"[All
Fields]) OR "nutritional status"[All Fields])
```

OR "nutrition"[All Fields] OR "nutritional sciences"[MeSH Terms] OR ("nutritional"[All Fields] AND "sciences"[All Fields]) OR "nutritional sciences"[All Fields] AND ("adolescent"[MeSH Terms] OR "adolescent"[All Fields] OR "adolescents"[All Fields])) OR (("violence"[MeSH Terms] OR "violence"[All Fields]) AND against[All Fields] AND ("adolescent"[MeSH Terms] OR "adolescent"[All Fields] OR "adolescents"[All Fields])) AND (((("guideline"[Publication Type] OR "guidelines as topic"[MeSH Terms] OR "guideline"[All Fields]) OR strategy[All Fields]) OR ("United Evangelical Action"[Journal] OR "Action Natl"[Journal] OR "action"[All Fields]) AND plan[All Fields]) AND ("policy"[MeSH Terms] OR "policy"[All Fields])) AND ("Bangladesh"[MeSH Terms] OR "Bangladesh"[All Fields]).

We also searched the gray literature in the Demography and Health Survey (DHS) database, WHO regional databases and Global Burden of Diseases (GBD) database and Bangladesh government and ministries home pages. Response rate was 91%.

Measurement

Quantitative data

GSHS capture information on a wide range of health indicators, including insufficient physical activity, alcohol consumption, any form of tobacco, sedentary behavior, insufficient fruits and vegetables consumption, overweight/obesity and psychological distress. Details of each measurement are described in Additional file 1.

For policy analysis, a pre-structured case methodology was adopted, using an existing conceptual NCD framework to define the structure for data collection and analysis [17]. The framework used the six objectives of the WHO Action Plan as listed in Additional file 2. The scope of the analysis is limited to the policies relevant to adolescent NCDs.

Data analysis

The data analysis was performed on a total of 2,989 students. Data were analyzed using the IBM SPSS (version 20) software. All analyses were adjusted for sample weight. Policy documents that are available in public domain and were published between 1971 and 2018 were identified through online searches in PubMed and Google Scholar. Extensive review of existing policies relevant to prevention and control of NCDs in Bangladesh was done in an earlier publication by using data display matrix. Furthermore, we will now focus on

the policy documents that may have direct or indirect impact on the health systems preparedness to tackle NCDs among Bangladeshi adolescents.

Ethical consideration

In each of the participating countries, the GSHS received ethics approval from the Ministry of Education or a relevant Institutional Ethics Review Committee, or both. Only adolescents and their parents who provided written or verbal consent participated. As the current study used retrospective publicly available data, we did not need ethics approval from any Institutional Ethics Review Committee.

Patient and public involvement

Patients were not involved in the study. The survey covered school-based young adolescents aged between 12 and 17 years old.


Results

Findings from the secondary analyses of GSHS data

Prevalence of NCD risk factors among the adolescents

Of the 2,972 adolescents aged 12–17 years, 1191 (40.07%) were males and 1781 (59.93%) were females. Figure 1 shows the prevalence of different NCD risk factors among the adolescents by gender. Insufficient vegetables consumption (boys: 69%, 95% CI 65–71%; girls: 68%, 95% CI 65–70%; $p = 0.46$), insufficient fruits consumption (boys: 77%, 95% CI 74–79% vs girls: 79%, 95% CI 76–80%; $p < 0.005$), ate fast food (boys: 56%, 95% CI: 52–59% vs girls: 47%, 95% CI 44–50%; $p < 0.005$), insufficient physically activity (boys: 44%, 95% CI 40–47% vs girls: 48%, 95% CI 45–51%; $p < 0.005$) and psychological distress (boys: 4%, 95% CI 2–5% vs girls: 6%, 95% CI 4–7%; $p < 0.005$) were found to be similar for adolescent boys and girls. Prevalence of alcohol consumption, used any form of tobacco and overweight/obesity were higher among the boys compared to the girls. Use of any form of tobacco was more than seven times higher among the boys compared to the girls.

We examined the clustering (presence of multiple risk factors in an individual) of NCD risk factors in the sample and have analyzed by gender. According to Fig. 2, only 1% adolescent did not have any NCD risk factor. Overall 14% had at least one risk factor, 22% had two risk factors, 29% had three risk factors and 34% had four or more risk factors.




Findings from the review of policy documents related to adolescents NCD risk

Policy response

A total of 38 documents were identified [16, 18–54]. Eight (21.1%) were research and/or surveys, eight were on established policies (21.1%), while eleven (29%) were on acts. Three (7.9%) were related to guidelines and eight

(21.1%) were strategic planning documents from government and non-government agencies/institutes (Fig. 3). Table 1 presents the summary of the abovementioned policy documents against the objectives of the WHO Action Plan.



Objective 1: prioritize NCDs prevention and control This objective was addressed by eight documents, among which three were strategy papers [18, 19, 25] and five were policy papers (Table 1) [26, 27, 30, 32, 33]. Government of Bangladesh already implemented a 10-year long adolescent reproductive health strategy from (ARHS) 2006–2016 [18]. After completing the period of ARHS, government has recently formulated and started to implement the National Strategy for Adolescent Health 2017–2030 with the alignment of sustainable development goals. This is supposed to be a comprehensive strategy covering all dimensions to ensure the proper wellbeing of adolescents. Adolescent sexual and reproductive health, violence against adolescents, adolescent nutrition and mental health of adolescents are identified as four strategic thematic areas for this strategy, including two cross-cutting issues of social and behavioral change communication and health systems strengthening. But in this strategy, most of the NCD risk factors among the adolescent overlooked.

Objective 2: strengthen national capacity and multi-sectoral action This objective was addressed by thirty documents. Of whom, eight were strategy papers [18–25], eleven were policy acts [34–44], three were guidelines [45–47] and eight were policy papers (Table 1) [26–33]. Both national and international agencies extend their financial and technical support to various health promotion programs and researches targeted toward particular age groups, e.g., the adolescents.

Objective 3: reduce modifiable risk factors for NCDs This objective was addressed by twenty seven documents, among which eight were strategy papers [18–25], eight were policy papers [26–33] and eleven were acts [34–44] (Table 1). The “Smoking and Tobacco Products Usage (Control) Act” of 2005 had addressed many issues regarding this objective. The act was amended in 2013 and had strictly restricted the harmful use of tobacco products by all age groups, especially the adolescents as the purchase and sale of tobacco products to the underage (below the age of 18 years) was banned [39].

Objective 4: strengthen health systems This objective was addressed by twenty documents, among which eight were strategy papers [15–22], eight were policy papers [23–30], one survey [42–44] and three were guideline [42–44] documents (Table 1). The National NCD Risk Factor Survey 2011 had guided several initiatives to tackle NCDs. Non-Communicable Diseases Centre (NCDC) has been established within the Ministry of Health and Family Welfare. NCD corners in 300 Upazila Health Complexes have facilitated the access to health care to the grass root level. However, in the context of health systems structure in Bangladesh, still there is no specific counselling corner for adolescent, especially adolescent’s mental health wellbeing [20].

Objective 5: national capacity for high-quality research This objective was addressed by sixteen docu-




Table 1 Summary of policy documents for each objective of the Action Plan

List of policy documents and published year	Objective 1	Objective 2	Objective 3	Objective 4	Objective 5	Objective 6
<i>Strategies</i>						
Bangladesh adolescent reproductive health strategy (2006) [18]	✓	✓	✓	✓	✓	✓
Strategic plan for surveillance and prevention of NCDs in Bangladesh 2011–2015 (2011) [19]	✓	✓	✓	✓	✓	✓
NCD corner initiative (2013) [20]	×	✓	✓	✓	✓	✓
National communication strategy and action plan for reduction of NCD high risk behaviours in Bangladesh 2014–2016 (2014) [21]	×	✓	✓	✓	✓	✓
The seventh 5-year plan 2016–2020 (2016) [22]	×	✓	✓	✓	✓	✓
Health, Nutrition and Population Strategic Investment Plan, HNPSIP 2016–2021 (2016) [23]	×	✓	✓	✓	✓	✓
The multi-sectoral action plan for the NCD control and prevention 2018–2025 (2018) [24]	×	✓	✓	✓	✓	✓
National Strategy for Adolescent Health 2017–2030 (2017) [25]	✓	✓	✓	✓	✓	✓
<i>Policies</i>						
National sports policy (1998) [26]	✓	✓	✓	✓	✓	×
National Food Policy Plan of Action 2008–2015 (2008) [27]	✓	✓	✓	✓	✓	×
Education Policy (2010) [28]	×	✓	✓	✓	✓	×
Child Labour Elimination Policy (2010) [29]	×	✓	✓	✓	✓	×
The National Health Policy (2011) [30]	✓	✓	✓	✓	✓	×
Bangladesh Population Policy (2012) [31]	×	✓	✓	✓	✓	×
Bangladesh National Children Policy (2011) [32]	✓	✓	✓	✓	✓	×
Nutrition Policy (2015) [33]	✓	✓	✓	✓	✓	×
<i>Legislation Acts</i>						
The Child Marriage Restraint Act 1929 (amended in 1983) (1929) [34]	×	✓	✓	×	×	×
Cruelty to Women (Deterrent Punishment) Act (1983) [35]	×	✓	✓	×	×	×
Narcotics control act (1990) [36]	×	✓	✓	×	×	×
Playground, open space, park and natural water reservoir conservation act (2000) [37]	×	✓	✓	×	×	×
Prevention of Oppression against Women and Children Act 2000 (amended in 2003) [38]	×	✓	✓	×	×	×
Immoral Traffic Act (2010) [39]	×	✓	✓	×	×	×
The Human Trafficking Prevention and Deterrence Act (2012) [40]	×	✓	✓	×	×	×
Safe food act (2013) [41]	×	✓	✓	×	×	×
Smoking and tobacco products usage (control) (amendment) Act (2013) [42]	×	✓	✓	×	×	×
Children Act (2013) [43]	×	✓	✓	×	×	×
Mental health act (2014) [44]	×	✓	✓	×	×	×
<i>Guideline</i>						
A field guide for detection management surveillance of arsenicosis (2005) [45]	×	✓	×	✓	×	×
Acute chest pain management protocol (2012) [46]	×	✓	×	✓	×	×
Dietary Guideline Bangladesh (2013) [47]	×	✓	×	✓	×	×
<i>Surveys and researches</i>						
Strategies to improve reproductive health services for adolescents in Bangladesh: a community-based study (2005) [48]	×	×	×	×	✓	✓
National cancer registry survey (2007) [49]	×	×	×	×	✓	✓
Global Youth Tobacco Survey Bangladesh (2013) [50]	×	×	×	×	✓	✓
Bangladesh demographic and health survey (2011) [51]	×	×	×	×	✓	✓
NCD risk factor survey (2011) [52]	×	×	×	✓	✓	✓
National Micronutrients Status Survey 2011–12 (2011) [53]	×	×	×	×	✓	✓

Table 1 (continued)

List of policy documents and published year	Objective 1	Objective 2	Objective 3	Objective 4	Objective 5	Objective 6
Bangladesh national health facility survey (2014) [54]	×	×	×	×	✓	✓
Global school health survey (2014) [16]	×	×	×	×	✓	✓
Total	8	30	27	20	24	16

ments, among which eight were strategy papers [18–25], eight were policy papers [26–33] and eight were surveys or research [16, 48–54] (Table 1). The Directorate General of Health Services (DGHS) [52] invests in NCD-related researches. In 2010, the DGHS has conducted the NCD Risk Factor Survey. Later in 2013, the National Center for Control of Rheumatic Fever and Heart Disease had conducted the GATS [50], and the Global School-based Student Health Survey 2014 [16]. NIPORT conducts Bangladesh Demographic and Health Survey at every 4 years interval [51]. International agencies such as WHO and Japan International Cooperation Agency (JICA) also facilitate several surveys with their financial and technical support.

Objective 6: monitor the trends and evaluate progress This objective was addressed by sixteen documents, among which eight were strategy papers [18–25] and eight were surveys or research (Table 1) [16, 48–54]. This objective has been addressed by NCD Risk Factor Survey, GATS, Global School Health survey, and the mental health survey [16, 48–54]. The Global School Health survey had important findings on the nutritional and mental health status of school-going students in Bangladesh [16]. BDHS 2011 had incorporated a few components of NCD. But BDHS 2011 completely ignored adolescents NCDs [51]. Bangladesh health facility survey also described current situation of the respective areas [54].

Discussion

This study provides the contemporary evidence on NCD risk factors among the adolescent in Bangladesh, and we have presented the policy provisions to combating the NCD risk factors among the adolescents in Bangladesh. The study reported high prevalence of NCD risk factors among the adolescent. More than thirty percent of adolescents had four or more NCD risk factors. Moreover, analysis of the policy documents suggests that, over the years, the government of Bangladesh has undertaken many endeavors to tackle NCDs. Some of the initiatives demonstrate innovative approach and effectiveness, such as the NCD corners initiative at Upazila Health Complexes. The ban on tobacco product advertisements and plain packaging of tobacco products were also considered to be bold steps

toward prevention of a major NCD risk factor. The National Safe Food Act of 2013 promoted food safety for all.

In our study, we found that insufficient fruits, insufficient vegetables and fats food consumption are the three common risk factors among the adolescent. This is similar to an analysis in Vietnam that reported low fruit/vegetable intake and unhealthy diet were common among school-going adolescents aged 13–17 years [7, 9]. Similar to some other Asian countries, we found that smoking and alcohol prevalence is comparatively low in Bangladesh [8, 10]. Although in our study we did not explore any school health promotion programs for improving adolescent health wellbeing, a recent study in Vietnam reported high quality of health promotion programs associated with lower the odds of lifestyle risk behaviors [7, 9].

According to our study, nearly fifty percent of adolescents are physically inactive. With a vision of NCD prevention from early age, WHO has outlined guidelines for games and physical activities within educational institutes [8]. Bangladesh has incorporated this vision in the National Children's Policy, educational policy and the National Sports Policy. Unfortunately, these policies do not reflect on a large number of educational institutions, especially in urban areas, where there is not enough open space for physical activities or sports.

Appropriate nutrition is important for adolescence to secure current and future generation health [32]. In our study, we found that around 10% adolescents were overweight/obese, and 78% and 68% adolescent consumed insufficient fruits and vegetables, respectively. More than fifty percent adolescent ate fast food. It demonstrated that prevalence of unhealthy food habit is increasing, with the trend continuing into early adulthood [33]. This is an impending public health problem that needs further action as adolescent obesity strongly predicts adult obesity and associated morbidity. Addressing this issue at a population level is a critical starting point to avert potential long-term impacts of adolescent obesity [34].

Bangladesh has an adolescent and youth population of approximately 52 million, amounting to 1/3rd of the country's total population. [41]. According our study, NCD risk factors among the adolescent comparatively high. At the same time, the strict enforcement of these

policies is still very challenging in country context of Bangladesh. Like many other countries [55] it is challenging to design and implement appropriate actions for the wellbeing of adolescent health. Greater emphasis must also be placed on further enhancing the data collection system with a view to strengthening the Health Management Information System so as to better serve the adolescents.

A recent study in Bangladesh entitled “A scorecard for tracking actions to reduce the burden of non-communicable diseases” reported that among the four domains of governance, risk factor surveillance, research, and health system response, the country’s performance score was low in three domains, except for the governance domain (moderate performance) [56]. In addition to that, the country lacks any integrated community public health program focused on monitoring NCDs amongst adolescents on a regular basis. Bangladesh also lacks any national surveillance program focused on NCDs. Only a few tertiary hospitals maintain such NCD surveillance systems [57] but most of them targeting adult population. Another study identified a total of 11 NCD programs in Bangladesh focusing on tobacco-related illnesses, diabetes or cardiovascular diseases [58]. Unfortunately, to date, there are no such programs being developed to target the NCDs prevention for adolescents in Bangladesh.

Strength and limitations

The strengths of this study include nationally representative sample taken through standardized questionnaire. To the best of our knowledge, this is a first large population-based study ever been conducted to examine the NCD risk factors among adolescents in Bangladesh. The present study has some limitations. Firstly, data were collected from adolescents enrolled in schools. Secondly, as the questionnaire was self-reported, it is possible that some respondents might have misreported the questions asked. Thirdly, the scope of this policy analysis was limited to publicly available documents in relation to the objectives of the WHO Action Plan.

Conclusion

Adolescents in Bangladesh report a high prevalence of concurrent multiple risk factors for NCDs. The government of Bangladesh has demonstrated political commitment toward adhering the global action plans to reduce the burden of NCDs in all age groups. However, more planning and coordination of existing programs are warranted with a focus on adolescents. Given the growing burden of NCD risk factors in this age group, it is likely to result in excessive NCDs burden in near future, if no appropriate and immediate actions are

taken. Priority should be given on the prevention of modifiable risk factors from an early age. Programs and interventions targeted at educational institutions could provide an integrated platform for NCDs prevention and control. Further research and monitoring of the trends of NCDs among adolescents would guide us toward appropriate strategies to address the NCD burden.

Abbreviations

NCD: Non-communicable disease; LMICs: Low and middle-income countries; SDGs: Sustainable development goals; GSHS: Global School-based Health Survey; WHO: World Health Organization; CDC: Centre for Disease Control and Prevention; UNAIDS: United Nations Programme on HIV and AIDS.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s41043-022-00294-x>.

Additional file 1. Measurement of NCD risk factors (insufficient physical activity, alcohol consumption, any form of tobacco, sedentary behavior, insufficient fruits and vegetables consumption, overweight/obesity and psychological distress).

Additional file 2. Six objectives of the WHO Action Plan.

Acknowledgements

The data used in this study were obtained from the open access dataset of Global School-based Student Health Survey (Bangladesh). The authors acknowledge the contributions of the GSHS team for their efforts in providing open access to the dataset. We thank George Patton for his initial feedback. The authors gratefully acknowledge support from the Australian Research Council (ARC) Centre of Excellence for Children and Families over the Life Course (CE200100025).

Author contributions

All authors read and approved the final manuscript.

Funding

The author(s) received no specific funding for this work.

Data availability

We thank the US Centers for Disease Control and WHO for making Global School-based Student Health Survey (GSHS) data publicly available for analysis.

Declarations

Ethics approval and consent to participate

In each of the participating countries, the GSHS received ethics approval from the Ministry of Education or a relevant Institutional Ethics Review Committee, or both. Only adolescents and their parents who provided written or verbal consent participated. As the current study used retrospective publicly available data, we did not need ethics approval from any Institutional Ethics Review Committee.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Institute for Social Science Research, The University of Queensland, QLD, Australia. ²ARC Centre of Excellence for Children and Families Over the Life Course, The University of Queensland, QLD, Australia. ³Burnet Institute, Melbourne, VIC, Australia. ⁴Institute of Health Economics, University of Dhaka, Dhaka, Bangladesh. ⁵NCD Alliance, Geneva, Switzerland. ⁶Utrecht University, Utrecht, Netherlands. ⁷Maternal and Child Health Division (MCHD), icddr, Dhaka, Bangladesh. ⁸Department of Epidemiology and Biostatistics, Arnold School of Public Health, University of South Carolina, Columbia, SC, USA. ⁹Department of Public Health, North South University, Dhaka, Bangladesh. ¹⁰School of Health, Medical and Applied Sciences, CQ University, Sydney, Australia.

Received: 27 July 2021 Accepted: 14 April 2022

Published online: 16 May 2022

References

- Magadi MA. Multilevel determinants of teenage childbearing in sub-Saharan Africa in the context of HIV/AIDS. *Health Place*. 2017;46:37–48.
- Steinberg L, Morris AS. Adolescent development. *Annu Rev Psychol*. 2001;52:83–110.
- Patton GC, Olsson CA, Skirbekk V, et al. Adolescence and the next generation. *Nature*. 2018;554(7693):458.
- Plan UK. Noncommunicable disease prevention and adolescents, 2017. Retrieved from <https://plan-uk.org/file/ncd-prevention-and-adolescents-report/download?token=tkG9kOxg>.
- Khan A, Uddin R, Islam SMS. Clustering patterns of behavioural risk factors for cardiovascular diseases in Bangladeshi adolescents: a population-based study. *Health Policy Technol*. 2019;8(4):386–92.
- Urmy NJ, et al. Noncommunicable disease risk factors among adolescent boys and girls in Bangladesh: evidence from a national survey. *Osong Public Health Res Perspect*. 2020;11(6):351.
- Long KQ, et al. Clustering lifestyle risk behaviors among Vietnamese adolescents and roles of school: a Bayesian multilevel analysis of global school-based student health survey 2019. *Lancet Reg Health West Pac*. 2021;15: 100225.
- Teh CH, et al. Clustering of lifestyle risk behaviours and its determinants among school-going adolescents in a middle-income country: a cross-sectional study. *BMC Public Health*. 2019;19(1):1–10.
- Patton GC, Sawyer SM, Santelli JS, et al. Our future: a Lancet commission on adolescent health and wellbeing. *Lancet*. 2016;387:423–78.
- Bonomo YA, Patton GC, Bowes G. What are the longer term outcomes of adolescent alcohol consumption in young adulthood? Results from a 10-year cohort study. *Alcohol Clin Exp Res*. 2006;30:117A.
- Viner RM, Taylor B. Adult outcomes of binge drinking in adolescence: findings from a UK national birth cohort. *J Epidemiol Community Health*. 2007;61:902–7.
- Riley EP, McGee CL. Fetal alcohol spectrum disorders: an overview with emphasis on changes in brain and behavior. *Exp Biol Med (Maywood)*. 2005;230:357–65.
- Patton GC, Coffey C, Carlin JB, et al. Overweight and obesity between adolescence and young adulthood: a 10-year prospective cohort study. *J Adolesc Health*. 2011;48:275–80.
- Ahsan KZ, Alam MN, Streatfield PK, et al. Has Bangladesh entered the fourth stage of the epidemiologic transition. In: Proceedings of the international seminar on mortality: past, present and future, The University of Campinas, Brazil; 2017.
- Biswas T, Townsend N, Islam MS, et al. Association between socioeconomic status and prevalence of non-communicable diseases risk factors and comorbidities in Bangladesh: findings from a nationwide cross-sectional survey. *BMJ Open*. 2019;9(3): e025538.
- World Health Organization. Global school-based student health survey, Bangladesh 2014. 2014. Retrieved from <http://www.who.int/chp/gshs/bangladesh/en/>. Accessed 7 Apr 2019.
- Biswas T, Pervin S, Tanim MIA, et al. Bangladesh policy on prevention and control of non-communicable diseases: a policy analysis. *BMC Public Health*. 2017;17(1):582.
- Ministry of Health and Family Welfare. Bangladesh adolescent reproductive health strategy 2006. Retrieved from: [http://www.dgfp.gov.bd/site/page/dd79f17e-0ee5-4aad-8619-b3b8ebdb37f0/National-ARH-Strategy-Bangladesh-\(2006\)](http://www.dgfp.gov.bd/site/page/dd79f17e-0ee5-4aad-8619-b3b8ebdb37f0/National-ARH-Strategy-Bangladesh-(2006)). Accessed 7 Apr 2019.
- Ministry of Health and Family Welfare, Directorate general of health services. Strategic plan for surveillance and prevention of non-communicable diseases in Bangladesh 2011–2015. Retrieved from https://www.ghdonline.org/uploads/BGD_SP_Surveillance_NCD_2011.pdf. Accessed 7 Apr 2019.
- Shahriar TH. Move to set up NCD corners in public hospitals. *The Daily Sun*; 2017. Retrieved from <http://www.daily-sun.com/printversion/details/202375/Move-to-set-up-NCD-corners-in-public-hospitals->. Accessed 7 Apr 2019.
- World Health Organization. National communication strategy & action plan for reduction of NCD high risk behaviors in Bangladesh, 2014–2016; 2015.
- General Economics Division, Planning Commission, Government of the People's Republic of Bangladesh. The 7th five year plan FY2016–FY2020: accelerating growth, empowering citizens. Dhaka; 2015.
- Ministry of Health and Family Welfare. Health, Nutrition and Population Strategic Investment Plan (HNPSIP) 2016–2021. Dhaka; 2016. Retrieved from https://www.bma.org.bd/pdf/strategic_Plan_HPNSDP_2016-21.pdf. Accessed 7 Apr 2019.
- Non-communicable Disease Control Programme, Directorate General of Health Services, Health Services Division, Ministry of Health and Family Welfare. Multisectoral Action Plan for Prevention and Control of Non-Communicable Diseases 2018–2025 with a Three-Year Operational Plan. Dhaka; 2018.
- Ministry of Health and Family Welfare. National Strategy for Adolescent Health 2017–2030. Dhaka; 2016. Retrieved from <http://coastbd.net/wp-content/uploads/2017/07/National-Strategy-for-Adolescent-Health-2017-2030-Final-Full-Book-21-06-17.pdf>. Accessed 7 Apr 2019.
- Ministry of Youth and Sports. National Sports Policy 1998; 1998. https://www.mof.gov.bd/en/budget/14_15/gender_budget/en/10_36_Youth-English.pdf. Accessed 7 Apr 2019.
- Food Planning and Monitoring Unit (FPMU), Ministry of Food and Disaster Management Government of the People's Republic of Bangladesh. National Food Policy: Plan of Action 2008–2015. Dhaka; 2008.
- Ministry of Education, Government of the People's Republic of Bangladesh. National Education Policy 2010. Dhaka; 2010. Retrieved from http://old.moedu.gov.bd/index.php?option=com_content&task=view&id=338&Itemid=416. Accessed April 7, 2019.39. Ministry of Labour and Employment, Government of the People's Republic of Bangladesh. National Child Labour Elimination Policy. Dhaka; 2010.
- Ministry of Labour and Employment. Government of the People's Republic of Bangladesh. Dhaka: National Child Labour Elimination Policy; 2010.
- Ministry of Health and Family Welfare. National Health Policy 2011. Dhaka; 2011. Retrieved from http://bdhealth.com/App_pages/Main/NationalHPB.aspx. Accessed 7 Apr 2019.
- Ministry of Health and Family Welfare. Bangladesh Population Policy 2012. Dhaka; 2012. Retrieved from http://bangladesh.gov.bd/sites/default/files/files/bangladesh.gov.bd/policy/98896a22_df81_4a82_b70c_24125dec56d7/Bangladesh-Population-Policy-2012.pdf. Accessed 7 Apr 2019.
- Ministry of Women and Children Affairs. National Children Policy 2011. Retrieved from <http://ecd-bangladesh.net/document/documents/National-Children-Policy-2011-English-04.12.2012.pdf>. Accessed 17 May 2019.
- Ministry of Health and Family Welfare. National Nutrition Policy 2015. Retrieved from <http://extwprlegs1.fao.org/docs/pdf/bgd152517.pdf>. Accessed 17 May 2019.
- Government of Bangladesh. The Child Marriage Restraint Act 1929 (Amended in 1983). Retrieved from http://bdlaws.minlaw.gov.bd/print_sections_all.php?id=149. Accessed 17 May 2019.
- Government of Bangladesh. Cruelty to women (deterrent punishment) ordinance; 1983. Retrieved from <http://legislib.com/mediaOpen?id=52>. Accessed 17 May 2019.
- Ministry of Home Affairs. Narcotic Control Act, 1990. Dhaka; 1990. Retrieved from <http://www.dnc.gov.bd/rulsacts.html>. Accessed 7 Apr 2019.
- Government of Bangladesh. Mega City, Divisional Town and District Town's Municipal Areas Including Country's All the Municipal Areas' Playground, Open Space, Park and Natural Water Reservoir Conservation Act,

2000. Retrieved from <http://bwdb.gov.bd/archive/pdf/198.pdf>. Accessed 7 Apr 2019.
38. Government of Bangladesh. Prevention of Oppression against Women and Children Act 2000 (Amended in 2003). Retrieved from http://iknow.politics.org/sites/default/files/prevention_act_bangladesh.pdf. Accessed 17 May 2019.
39. Legislative and Parliamentary Affairs Division, Ministry of Law, Justice and Parliamentary Affairs. The Suppression of Immoral Traffic Act, 1933 (Act No. VI of 1933). Dhaka; 2010. Retrieved from http://bdlaws.minlaw.gov.bd/print_sections_all.php?id=159. Accessed 23 May 2019.
40. Ministry of Law, Justice and Parliamentary affairs. The Human Trafficking Prevention and Deterrence Act 2012. Retrieved from <https://www.refworld.org/pdfid/543f75664.pdf>. Accessed 17 May 2019.
41. Ministry of Food. Safe Food Act, 2013. Dhaka; 2013. Retrieved from <http://www.thedailystar.net/safe-food-act-to-be-effective-from-feb-1-61978>. Accessed 7 Apr 2019.
42. Ministry of Health and Family Welfare. An Act for the Amendment of Smoking and Tobacco Products Usage (Control) Act. Dhaka; 2013. Retrieved from <https://www.tobaccocontrollaws.org/files/live/Bangladesh/Bangladesh%20-%20TC%20Amdt.%20Act%202013.pdf>. Accessed 7 Apr 2019.
43. Legislative and Parliamentary Affairs Division, Ministry of Law, Justice and Parliamentary Affairs. Children's Act, 2013 (Act No. 24 of 2013), Dhaka; 2010. Retrieved from http://bdlaws.minlaw.gov.bd/bangla_pdf_part.php?id=1119. Accessed 17 May 2019.
44. Mental Health Act 2014. Retrieved from <http://www.thedailystar.net/draft-bangladesh-mental-health-act-2014-rights-perspective-51534>. Accessed 7 Apr 2019.
45. World Health Organization. A field guide: detection, management and surveillance of arsenicosis. New Delhi; 2006. Retrieved from: http://www.searo.who.int/entity/water_sanitation/documents/arsenicosis2006/en/. Accessed 7 Apr 2019.
46. Management of Acute Chest Pain. Retrieved from <http://hsmgdghs-bd.org/Documents/CMP/Management%20of%20Acute%20chest%20pain.pdf>. Accessed 1 Mar 2017.
47. Bangladesh Institute of Research and Rehabilitation in Diabetes, Endocrine and Metabolic Disorders (BIRDEM). Retrieved from: Dietary Guidelines for Bangladesh. Dhaka; 2013. <http://www.fao.org/3/a-as880e.pdf>. Accessed 17 May 2019.
48. Huq NL, Nahar Q, Larson CP, et al. Strategies to improve reproductive health services for adolescents in Bangladesh: a community-based study. *Icddr Work Pap No 164*. 2005.
49. Ministry of Health and Family Welfare. National Cancer Registry Survey, 2007. Dhaka; 2008. Retrieved from http://dghs.gov.bd/bn/licts_file/images/Strategy/2009_NationalCancerControlStrategy2009-15.pdf. Accessed 7 Apr 2019.
50. World Health Organization, Regional Office for South-East Asia. Global Youth Tobacco Survey (GYTS), Bangladesh Report, 2013; 2015. Retrieved from <http://www.who.int/iris/handle/10665/164335>. Accessed 17 May 2019.
51. National Institute of Population Research and Training. Bangladesh Demographic and Health Survey 2011. Dhaka; 2013. Retrieved from <https://dhsprogram.com/pubs/pdf/FR311/FR311.pdf>. Accessed 7 Apr 2019.
52. World Health Organization. Non-communicable disease risk factor survey Bangladesh 2010; 2011. Retrieved from http://www.who.int/chp/steps/2010_STEPS_Report_Bangladesh.pdf. Accessed 7 Apr 2019.
53. icddr, UNICEF Bangladesh, GAIN, Institute of Public Health and Nutrition. National Micronutrients Status Survey 2011–12. Retrieved from https://static1.squarespace.com/static/56424f6ce4b0552eb7fdc4e8/t/57490d3159827e39bd4d2314/1464405328062/Bangladesh_NMS_final_report_2011-12.pdf. Accessed 17 May 2019.
54. National Institute of Population Research and Training. Bangladesh National Health Facility Survey 2014. Dhaka; 2016. Retrieved from <https://dhsprogram.com/pubs/pdf/SPA23/SPA23.pdf>. Accessed 1 Mar 2017.
55. Islam, A. and T.J.C.D.I. Biswas, Chronic non-communicable diseases and the healthcare system in Bangladesh: current status and way forward, vol 1, no 2; 2014. p. 6.
56. Roman AV, Perez W, Smith RJTL. A scorecard for tracking actions to reduce the burden of non-communicable diseases. *Lancet*. 2015;386(9999):1131–2.
57. El-Saharty S, et al. Tackling noncommunicable diseases in Bangladesh: now is the time. The World Bank; 2013.
58. Bleich SN, et al. Noncommunicable chronic disease in Bangladesh: overview of existing programs and priorities going forward. *Health Policy*. 2011;100(2–3):282–9.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

