

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Drew, DA;Kim, AE;Lin, Y;Qu, C;Morrison, J;Lewinger, JP;Kawaguchi, E;Wang, J;Fu, Y;Zemlianskaia, N;Díez-Obrero, V;Bien, SA;Dimou, N;Albanes, D;Baurley, JW;Wu, AH;Buchanan, DD;Potter, JD;Prentice, RL;Harlid, S;Arndt, V;Barry, EL;Berndt, SI;Bouras, E;Brenner, H;Budiarto, A;Burnett-Hartman, A;Campbell, PT;Carreras-Torres, R;Casey, G;Chang-Claude, J;Conti, DV;Devall, MAM;Figueiredo, JC;Gruber, SB;Gsur, A;Gunter, MJ;Harrison, TA;Hidaka, A;Hoffmeister, M;Huyghe, JR;Jenkins, MA;Jordahl, KM;Kundaje, A;Le Marchand, L;Li, L;Lynch, BM;Murphy, N;Nassir, R;Newcomb, PA;Newton, CC;Obón-Santacana, M;Ogino, S;Ose, J;Pai, RK;Palmer, JR;Papadimitriou, N;Pardamean, B;Pellatt, AJ;Peoples, AR;Platz, EA;Rennert, G;Ruiz-Narvaez, E;Sakoda, LC;Scacheri, PC;Schmit, SL;Schoen, RE;Stern, MC;Su, YR;Thomas, DC;Tian, Y;Tsilidis, KK;Ulrich, CM;Um, CY;van Duijnhoven, FJB;Van Guelpen, B;White, E;Hsu, L;Moreno, V;Peters, U;Chan, AT;Gauderman, WJ

Title:

Two genome-wide interaction loci modify the association of nonsteroidal anti-inflammatory drugs with colorectal cancer

Date:

2024-05-31

Citation:

Drew, D. A., Kim, A. E., Lin, Y., Qu, C., Morrison, J., Lewinger, J. P., Kawaguchi, E., Wang, J., Fu, Y., Zemlianskaia, N., Díez-Obrero, V., Bien, S. A., Dimou, N., Albanes, D., Baurley, J. W., Wu, A. H., Buchanan, D. D., Potter, J. D., Prentice, R. L. ,... Gauderman, W. J. (2024). Two genome-wide interaction loci modify the association of nonsteroidal anti-inflammatory drugs with colorectal cancer. *Science Advances*, 10 (22), <https://doi.org/10.1126/sciadv.adk3121>.

Persistent Link:

<https://hdl.handle.net/11343/353746>

License:

CC BY-NC

CANCER

Two genome-wide interaction loci modify the association of nonsteroidal anti-inflammatory drugs with colorectal cancer

David A. Drew^{1,2,†*}, Andre E. Kim^{3,†}, Yi Lin⁴, Conghui Qu⁴, John Morrison³, Juan Pablo Lewinger³, Eric Kawaguchi³, Jun Wang⁵, Yubo Fu³, Natalia Zemlianskaia³, Virginia Díez-Obrero^{6,7,8}, Stephanie A. Bien⁴, Niki Dimou⁹, Demetrius Albanes¹⁰, James W. Baurley^{11,12}, Anna H. Wu⁵, Daniel D. Buchanan^{13,14,15}, John D. Potter^{4,16}, Ross L. Prentice⁴, Sophia Harlid¹⁷, Volker Arndt¹⁸, Elizabeth L. Barry¹⁹, Sonja I. Berndt¹⁰, Emmanouil Bouras²⁰, Hermann Brenner^{18,21,22}, Arif Budiarto¹¹, Andrea Burnett-Hartman²³, Peter T. Campbell²⁴, Robert Carreras-Torres^{6,25}, Graham Casey²⁶, Jenny Chang-Claude^{27,28}, David V. Conti³, Matthew A.M. Devall²⁹, Jane C. Figueiredo^{3,30}, Stephen B. Gruber^{31,32}, Andrea Gsur³³, Marc J. Gunter^{9,34}, Tabitha A. Harrison⁴, Akihisa Hidaka⁴, Michael Hoffmeister¹⁸, Jeroen R. Huyghe⁴, Mark A. Jenkins³⁵, Kristina M. Jordahl^{4,36}, Anshul Kundaje^{37,38}, Loic Le Marchand³⁹, Li Li^{29,40}, Brigid M. Lynch^{35,41}, Neil Murphy⁹, Rami Nassir⁴², Polly A. Newcomb^{4,43}, Christina C. Newton⁴⁴, Mireia Obón-Santacana^{6,45,46}, Shuji Ogino^{47,48,49,50}, Jennifer Ose^{51,52}, Rish K. Pai⁵³, Julie R. Palmer⁵⁴, Nikos Papadimitriou⁹, Bens Pardamean¹¹, Andrew J. Pellatt⁵⁵, Anita R. Peoples^{51,52}, Elizabeth A. Platz⁵⁶, Gad Rennert^{57,58,59}, Edward Ruiz-Narvaez⁶⁰, Lori C. Sakoda^{4,61}, Peter C. Scacheri⁶², Stephanie L. Schmit^{63,64}, Robert E. Schoen⁶⁵, Mariana C. Stern⁵, Yu-Ru Su⁶⁶, Duncan C. Thomas³, Yu Tian^{27,67}, Konstantinos K. Tsilidis^{34,68}, Cornelia M. Ulrich^{51,52}, Caroline Y. Um⁴⁴, Fränzel J.B. van Duijnhoven⁶⁹, Bethany Van Guelpen^{17,70}, Emily White^{4,36}, Li Hsu^{4,71}†*, Victor Moreno^{6,45,46,72}‡, Ulrike Peters^{4,36}‡*, Andrew T. Chan^{1,2}‡, W. James Gauderman^{3,*‡}

Copyright © 2024 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution NonCommercial License 4.0 (CC BY-NC).

Regular, long-term aspirin use may act synergistically with genetic variants, particularly those in mechanistically relevant pathways, to confer a protective effect on colorectal cancer (CRC) risk. We leveraged pooled data from 52 clinical trial, cohort, and case-control studies that included 30,806 CRC cases and 41,861 controls of European ancestry to conduct a genome-wide interaction scan between regular aspirin/nonsteroidal anti-inflammatory drug (NSAID) use and imputed genetic variants. After adjusting for multiple comparisons, we identified statistically significant interactions between regular aspirin/NSAID use and variants in 6q24.1 (top hit *rs72833769*), which has evidence of influencing expression of *TBC1D7* (a subunit of the *TSC1-TSC2* complex, a key regulator of *MTOR* activity), and variants in 5p13.1 (top hit *rs350047*), which is associated with expression of *PTGER4* (codes a cell surface receptor directly involved in the mode of action of aspirin). Genetic variants with functional impact may modulate the chemopreventive effect of regular aspirin use, and our study identifies putative previously unidentified targets for additional mechanistic interrogation.

INTRODUCTION

Aspirin, a nonsteroidal anti-inflammatory drug (NSAID), is inversely associated with colorectal cancer (CRC) risk. In meta-analyses and systematic reviews of large observational studies, regular long-term use of aspirin is associated with CRC risk reduction of 20 to 30% (1–3). Reduction of CRC risk was also observed in well-designed clinical trials of colorectal neoplasia outcomes among individuals with Lynch syndrome or prior colorectal adenoma or CRC (4–10). However, the precise mechanism of action has not yet been fully elucidated, although several modes of action have been suggested for aspirin's anticancer effects (3, 11). Despite a potential overlap in mechanism (i.e., inhibition of prostaglandin synthesis), the relationship of non-aspirin NSAIDs (henceforth simply termed “NSAIDs”) and CRC risk is less consistent, potentially owing to more heterogeneous use in published studies, contamination of non-aspirin NSAID categories with aspirin use, or confounding by indication for use (i.e.,

individuals with higher inflammatory states). Because not all studies specifically differentiate between aspirin and other NSAID use, additional study of the impact of this broader drug class on CRC risk is warranted. Genetic variation is a key individual factor that likely interacts with aspirin and NSAIDs to ultimately determine CRC risk. In general, gene-drug interaction studies aim to clarify these relationships and implicate regions involved in the mode of action (12), which may identify subpopulations of individuals that might most benefit from an aspirin preventive strategy, particularly in light of the potential harms.

In this analysis, we conducted a genome-wide interaction scan (GWIS) of regular aspirin/NSAID use and imputed genetic markers. A previous GWIS, conducted on a smaller subset of individuals, identified interaction loci in regions 12p12.3 and 15q25.2 (13). We expanded upon that analysis by greatly increasing the sample size and using additional statistical methods that improve power to detect interaction loci and infer functional impact.

RESULTS

Regular aspirin/NSAID use and CRC risk

Our combined study sample included 72,667 individuals (30,806 cases and 41,861 controls) with known aspirin/NSAID use status and CRC outcome status (table S1). We considered regular aspirin and/or NSAID use as a combined variable (aspirin/NSAID) because not all studies collected data on aspirin use separate from other NSAID use, and aspirin and NSAIDs likely have common anticancer mechanisms. Complete study inclusion and exclusion criteria and how regular aspirin/NSAID use is defined are described in Materials and Methods. Secondary analyses restricted to studies with information on aspirin use only are also presented ($N = 72,137$; 30,574 cases and 41,563 controls). Regular aspirin/NSAID use was less prevalent among CRC cases compared to controls (34% versus 40%, respectively) as was aspirin use alone (27% of cases versus 31% of controls). As expected, CRC cases tended to be older, had a higher body mass index (BMI) and energy intake, had a greater proportion of family history of CRC, were less educated, and were more likely to be exposed to other known risk factors including heavy alcohol intake and tobacco smoking compared to controls.

In meta-analyses of study-specific associations, regular aspirin/NSAID use [odds ratio (OR) = 0.76; 95% confidence interval (CI) 0.72 to 0.81] and aspirin use alone (OR = 0.80; 95% CI = 0.76 to 0.84) were associated with reduced CRC risk (Fig. 1). There was statistically significant cross-study heterogeneity in the aspirin/NSAIDs and aspirin-only associations, which appeared to be largely due to study design (fig. S1). The estimated reductions in CRC risk were more pronounced in case-control studies for aspirin/NSAID use ($OR_{\text{aspirin/NSAID}} = 0.67$, 95% CI = 0.62 to 0.72) than in cohort studies ($OR_{\text{aspirin/NSAID}} = 0.85$, 95% CI = 0.80 to 0.91) ($P_{\text{het}} < 0.001$). Similar trends in estimates were observed for regular use of aspirin-only (case-control studies: $OR_{\text{aspirin}} = 0.72$, 95% CI = 0.67 to 0.77; cohort studies: $OR_{\text{aspirin}} = 0.88$, 95% CI = 0.83 to 0.94) ($P_{\text{het}} < 0.001$). Further adjustment by established CRC risk factors—including BMI, alcohol intake, smoking, and red meat consumption—did not substantially change OR estimates of aspirin/NSAID use (table S2). Analyses stratified by sex show nominally stronger inverse associations with regular aspirin/NSAID use and CRC risk among women compared to men (multivariate model $P_{\text{interaction}} = 0.014$), but there were no statistically significant sex differences for aspirin use


¹Clinical & Translational Epidemiology Unit, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA. ²Division of Gastroenterology, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA. ³Division of Biostatistics, Department of Population and Public Health Sciences, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA. ⁴Public Health Sciences Division, Fred Hutchinson Cancer Center, Seattle, Washington, USA. ⁵Department of Population and Public Health Sciences, Keck School of Medicine, University of Southern California, Los Angeles, California, USA. ⁶Colorectal Cancer Group, ONCOBELL Program, Bellvitge Biomedical Research Institute (IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain. ⁷Consortium for Biomedical Research in Epidemiology and Public Health (CIBERESP), Madrid, Spain. ⁸Department of Clinical Sciences, Faculty of Medicine, University of Barcelona, Barcelona, Spain. ⁹Nutrition and Metabolism Branch, International Agency for Research on Cancer, World Health Organization, Lyon, France. ¹⁰Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA. ¹¹Bioinformatics and Data Science Research Center, Bina Nusantara University, Jakarta, Indonesia. ¹²BioRealm LLC, Walnut, CA, USA. ¹³Colorectal Oncogenomics Group, Department of Clinical Pathology, The University of Melbourne, Parkville, Victoria 3010 Australia. ¹⁴University of Melbourne Centre for Cancer Research, Victorian Comprehensive Cancer Centre, Parkville, Victoria 3010 Australia. ¹⁵Genomic Medicine and Family Cancer Clinic, The Royal Melbourne Hospital, Parkville, Victoria, Australia. ¹⁶Research Centre for Hauora and Health, Massey University, Wellington, New Zealand. ¹⁷Department of Radiation Sciences, Oncology Unit, Umeå University, Umeå, Sweden. ¹⁸Division of Clinical Epidemiology and Aging Research, German Cancer Research Center (DKFZ), Heidelberg, Germany. ¹⁹Department of Epidemiology, Geisel School of Medicine at Dartmouth, Hanover, NH, USA. ²⁰Laboratory of Hygiene, Social & Preventive Medicine and Medical Statistics, Department of Medicine, School of Health Sciences, Aristotle University of Thessaloniki, Thessaloniki, Greece. ²¹Division of Preventive Oncology, German Cancer Research Center (DKFZ) and National Center for Tumor Diseases (NCT), Heidelberg, Germany. ²²German Cancer Consortium (DKTK), German Cancer Research Center (DKFZ), Heidelberg, Germany. ²³Institute for Health Research, Kaiser Permanente Colorado, Denver, CO, USA. ²⁴Department of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, NY, USA. ²⁵Digestive Diseases and Microbiota Group, Girona Biomedical Research Institute (IDIBGI), Salt, 17190 Girona, Spain. ²⁶Center for Public Health Genomics, University of Virginia, Charlottesville, VA, USA. ²⁷Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany. ²⁸University Medical Centre Hamburg-Eppendorf, University Cancer Centre Hamburg (UCC), Hamburg, Germany. ²⁹Department of Family Medicine, University of Virginia, Charlottesville, VA, USA. ³⁰Department of Medicine, Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA. ³¹Department of Medical Oncology & Therapeutics Research, City of Hope National Medical Center, Duarte, CA, USA. ³²Center for Precision Medicine, City of Hope National Medical Center, Duarte, CA, USA. ³³Center for Cancer Research, Medical University Vienna, Vienna, Austria. ³⁴Department of Epidemiology and Biostatistics, Imperial College London, School of Public Health, London, UK. ³⁵Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, Victoria, Australia. ³⁶Department of Epidemiology, School of Public Health, University of Washington, Seattle, WA, USA. ³⁷Department of Genetics, Stanford University, Stanford, CA, USA. ³⁸Department of Computer Science, Stanford University, Stanford, CA, USA. ³⁹University of Hawaii Cancer Center, Honolulu, HI, USA. ⁴⁰UVA Comprehensive Cancer Center, Charlottesville, VA, USA. ⁴¹Cancer Epidemiology Division, Cancer Council Victoria, Melbourne, Victoria, Australia. ⁴²Department of Pathology, School of Medicine, Umm Al-Qura University, Mecca, Saudi Arabia. ⁴³School of Public Health, University of Washington, Seattle, WA, USA. ⁴⁴Department of Population Science, American Cancer Society, Atlanta, GA, USA. ⁴⁵Unit of Biomarkers and Susceptibility (UBS), Oncology Data Analytics Program (ODAP), Catalan Institute of Oncology (ICO), L'Hospitalet del Llobregat, 08908 Barcelona, Spain. ⁴⁶Consortium for Biomedical Research in Epidemiology and Public Health (CIBERESP), 28029 Madrid, Spain. ⁴⁷Department of Epidemiology, Harvard T.H. Chan School of Public Health, Harvard University, Boston, MA, USA. ⁴⁸Program in MPE Molecular Pathological Epidemiology, Department of Pathology, Brigham and Women's Hospital, Boston, MA, USA. ⁴⁹Harvard Medical School, Boston, MA, USA. ⁵⁰Broad Institute of MIT and Harvard, Cambridge, MA, USA. ⁵¹Huntsman Cancer Institute, Salt Lake City, UT, USA. ⁵²Department of Population Health Sciences, University of Utah, Salt Lake City, UT, USA. ⁵³Department of Laboratory Medicine and Pathology, Mayo Clinic Arizona, Scottsdale, AZ, USA. ⁵⁴Slope Epidemiology Center at Boston University, Boston, MA, USA. ⁵⁵Department of Cancer Medicine, University of Texas MD Anderson Cancer Center, Houston, TX, USA. ⁵⁶Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA. ⁵⁷Department of Community Medicine and Epidemiology, Lady Davis Carmel Medical Center, Haifa, Israel. ⁵⁸Ruth and Bruce Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel. ⁵⁹Calit National Cancer Control Center, Haifa, Israel. ⁶⁰Department of Nutritional Sciences, University of Michigan School of Public Health, Ann Arbor, MI, USA. ⁶¹Division of Research, Kaiser Permanente Northern California, Oakland, CA, USA. ⁶²Department of Genetics and Genome Sciences, Case Western Reserve University, Cleveland, OH, USA. ⁶³Genomic Medicine Institute, Cleveland Clinic, Cleveland, OH, USA. ⁶⁴Population and Cancer Prevention Program, Case Comprehensive Cancer Center, Cleveland, OH, USA. ⁶⁵Department of Medicine and Epidemiology, University of Pittsburgh Medical Center, Pittsburgh, PA, USA. ⁶⁶Biostatistics Division, Kaiser Permanente Washington Health Research Institute, Seattle, WA, USA. ⁶⁷School of Public Health, Capital Medical University, Beijing, China. ⁶⁸Department of Hygiene and Epidemiology, University of Ioannina School of Medicine, Ioannina, Greece. ⁶⁹Division of Human Nutrition and Health, Wageningen University & Research, Wageningen, Netherlands. ⁷⁰Wallenberg Centre for Molecular Medicine, Umeå University, Umeå, Sweden. ⁷¹Department of Biostatistics, University of Washington, Seattle, WA, USA. ⁷²Department of Clinical Sciences, Faculty of Medicine and Health Sciences and Universitat de Barcelona Institute of Complex Systems (UBICS), University of Barcelona (UB), L'Hospitalet de Llobregat, 08908 Barcelona, Spain.

*Corresponding author. Email: dadrew@mgh.harvard.edu (D.A.D.); lih@fredhutch.org (L.H.); upeters@fredhutch.org (U.P.); jimg@usc.edu (W.J.G.)

†These authors contributed equally to this work.

‡These authors contributed equally to this work.

A Aspirin/NSAIDs



B Aspirin only




Fig. 1. Association of regular aspirin/NSAID use with CRC according to sex and tumor location. Results from meta-analysis of association between regular use of (A) aspirin/NSAID or (B) aspirin-only and colorectal cancer, overall and stratified by sex and tumor site. Models adjusted for age and sex. Heterogeneity measures include Cochran's Q statistic p-value (P_{het}) and Higgin's statistic (I^2), which describes the proportion of observed variance due to heterogeneity and not attributed to sampling error.

only (table S2). There were no notable differences in associations when stratifying models by tumor site (table S2).

GWIS results

We found no evidence of genomic inflation or residual population stratification in the genome-wide scan one degree of freedom (df) Gene x Environment (GxE) test P values for aspirin/NSAID or aspirin-only exposure variables (fig. S2). We identified a statistically

significant interaction between regular aspirin/NSAID use and *rs72833769* (chr6:12577203 T/C, $P = 1.27 \times 10^{-8}$; Table 1), a marker in locus 6q26 upstream of gene *PHACTR1* (Fig. 2A). The overall minor allele frequency for this single-nucleotide polymorphism (SNP) was 0.067 (Table 1). This interaction for *rs72833769* was similarly significantly associated when restricted to aspirin use only ($P_{\text{aspirin}} = 4.02 \times 10^{-9}$; Fig. 2B). While *rs72833769* did not have a direct marginal association with CRC risk (OR = 1.00; 95% CI = 0.96 to 1.05),

Table 1. Significant results from genome-wide interaction scans of regular aspirin/NSAID use and CRC risk. SNP, single-nucleotide polymorphism; Chr, chromosome; BP Position, base pair position based on NCBI Build37. “Overall” is the *P* value for the 1-df GxE test (*rs72833769*) or two-step procedure (*rs350047*). Imputed SNPs were coded as expected gene dosage. Multiplicative interaction terms were modeled as the product of Aspirin/NSAIDs and each SNP of interest. All statistical tests were two-sided.

Method	SNP	Chr	BP Position	Locus	Gene	Ref	Alt	Alt allele freq (1000G)	Exposure	<i>P</i> value (overall)	EDGE <i>P</i> value (step 1)	EDGE <i>P</i> value (step 2)	<i>P</i> value (3 df)
1-degree-of-freedom (df) GxE													
	<i>rs72833769</i>	6	12577203	6p24.1	Upstream of <i>PHACTR1</i>	T	C	0.02	Aspirin/NSAID	1.27×10^{-8}	–	–	–
									Aspirin only	4.02×10^{-9}	–	–	–
Two-step (EDGE) and 3-df													
	<i>rs350047</i>	5	40252294	5p13.1	<i>LINC00603</i> (upstream of <i>PTGER4</i>)	C	T	0.41	Aspirin/NSAID	8.20×10^{-8}	5.22×10^{-6}	4.41×10^{-5}	6.50×10^{-9}
									Aspirin only	2.00×10^{-8}	9.40×10^{-6}	1.08×10^{-5}	3.12×10^{-9}




Fig. 2. Manhattan plots of genome-wide interaction scans. (A) traditional logistic regression interaction test (1-df) for aspirin/NSAID (B) traditional logistic regression interaction test (1-df) for aspirin-only, (C) 3-df joint test for aspirin/NSAID, and (D) 3-df joint test for aspirin-only. Red line represents a genome-wide significance threshold of $5 \times 10^{-8}/2.5$ after adjustment for multiple testing.

stratified analyses showed regular aspirin/NSAID use or aspirin-only use were significantly associated with lower CRC risk only among homozygous carriers of the common allele T (aspirin/NSAID: $OR_{TT} = 0.73$, 95% CI = 0.71 to 0.76; aspirin only: $OR_{TT} = 0.76$, 95% CI = 0.73 to 0.79) (Tables 2 and 3 and fig. S3, A and B). In contrast, regular aspirin/NSAID use was not significantly associated with risk of CRC among individuals carrying the CT or CC genotype (aspirin/NSAID: $OR_{CT} = 0.94$, 95% CI = 0.85 to 1.03; $OR_{CC} = 1.44$, 95% CI = 0.82 to 2.54; aspirin-only: $OR_{CT} = 1.00$, 95% CI = 0.90 to 1.11; $OR_{CC} = 1.40$, 95% CI = 0.75 to 2.59). Adjustment for additional CRC risk factors did not materially affect the estimate of interaction (Tables 2 and 3, model 2). In addition, interaction effects for this SNP did not differ substantially by sex or tumor subsite, although interaction estimates were modestly stronger for colon locations than rectal tumors (table S3A).

We identified a second locus *rs350047* (chr5:40252294 C/T, $MAF_{1000\text{ Genomes}} = 0.48$) of interest using both our two-step EDGE method (fig. S3) and our 3-df joint test (Table 1). On the basis of the two-step testing, this SNP shows significant evidence of interaction with aspirin-only use; interactions were marginally significant for aspirin/NSAID use. For aspirin/NSAID use, *rs350047* achieved a step 1 “EDGE” P value of 5.22×10^{-6} and step 2 “GxE” P value of 4.41×10^{-5} (overall two-step P value = 8.2×10^{-8}). The corresponding P values for aspirin use only were 9.40×10^{-6} for step 1 and 1.08×10^{-5} for step 2, which in combination achieve genome-wide significance (overall two-step P value = 2.0×10^{-8}) (Table 1). This locus was statistically significant for both exposure variables based on 3-df joint test (aspirin/NSAID $P = 6.5 \times 10^{-9}$; aspirin only $P = 3.12 \times 10^{-9}$). Combined, the consistency across methods supports the identification of this hit. This genetic variant lies in locus 5p13.1 upstream of *PTGER4* and *LINC00603*, within a genomic region identified in a previously published GWAS of CRC risk and tagged by deletion-insertion marker *rs58791712* (14). Stratified analyses showed that regular aspirin/NSAID use and aspirin only use was inversely associated with CRC risk across all genotype groups, but the association was of greater magnitude among homozygous carriers of the T allele (aspirin/NSAID: $OR_{TT} = 0.67$, 95% CI = 0.63 to 0.72; aspirin-only: $OR_{TT} = 0.69$, 95% CI = 0.64 to 0.74) compared to heterozygous or homozygous carriers of the C allele (aspirin/NSAID: $OR_{CT} = 0.78$, 95% CI = 0.74 to 0.81; $OR_{CC} = 0.81$, 95% CI = 0.76 to 0.87; aspirin-only: $OR_{CT} = 0.81$, 95% CI = 0.77 to 0.85; $OR_{CC} = 0.85$, 95% CI = 0.79 to 0.91) (Tables 2 and 3 and fig. S3, C and D). Adjustment for additional CRC risk factors similarly did not affect the estimates of this interaction (Tables 2 and 3, model 2). The interaction is somewhat stronger among men than women (table S3B), although the three-way interaction P value was not statistically significant ($P_{G \times E \times S_{ex}} = 0.07$). Similarly, interaction effects at this locus were similar in magnitude across tumor site (table S3B). In our genome-wide scan for rare variants, we did not find any significant interactions with aspirin/NSAID or aspirin-only use.

Interactions stratified by CRC molecular subtypes

Case counts with available tumor marker information on *BRAF* and *KRAS* mutation status and the presence of CpG island methylator phenotype (CIMP) or microsatellite instability (MSI) are summarized in table S4. Generally, when fitting traditional case-control logistic regression models, interactions between aspirin/NSAID use and *rs72833769* or *rs350047* are replicated within the subset of cases with available tumor-marker data, at a significance level $P < 0.05$

(Fig. 3; overall cases versus controls). In stratified analyses according to the presence or absence of molecular subtype markers, significant association for the interaction between *rs72833769* and aspirin/NSAIDs were limited to tumors absent of each of the available tumor markers (Fig. 3A; *BRAF* wild-type, CIMP-low/negative, non-MSI-high, and *KRAS* wild-type tumors versus controls: all $P < 0.05$); however, estimates for the interaction between cases absent for these molecular markers and those positive for the individual markers were not significantly heterogeneous (all $P_{het} > 0.05$). In contrast, significant association for the interaction between *rs350047* and aspirin/NSAID is observed only among cases positive for *BRAF* mutation, CIMP-high, or MSI-high compared to controls, but not among cases with *KRAS* mutation nor absent any individual marker (Fig. 3B). Similarly, no significant heterogeneity between cases with the molecular marker present versus absent was observed (all $P_{het} > 0.05$), although the estimates for *BRAF* mutant versus wild-type and CIMP-high versus CIMP-low/negative approached significance. Further restricting the analysis to aspirin use alone did not materially alter estimates, but the reduced sample size resulted in slight attenuation of observed statistical significance.

Functional follow-up

The regional plot for *rs72833769* (6p24.1) shows several genes in the vicinity of this locus, including *PHACTR1* and *END1* (fig. S5A). Several data sources provide evidence that the region tagged by *rs72833769* plays a regulatory role in the transcription of neighboring genes. In CRC tumor and normal tissue, and CRC cell lines derived from work by Cohen *et al.* (15), the lead SNP *rs72833769* showed little evidence of functional activity. However, several SNPs in linkage disequilibrium (LD) with *rs72833769* coincided with accessible chromatin regions based on H3K27ac markers primarily in CRC tumors (table S5A and fig. S6A). This region also contains overlaps with regulatory regions based on H3K4me1, H3K4me3, H3K9ac, and H3K27 histone modification signals in connective, gastrointestinal, and immune cell types (16). ENSEMBL queries also show several regulatory features in this locus, in addition to a transcript region for lncRNA RP11-125 M16.1 (table S5A).


Evidence for an expression quantitative trait locus (eQTL) was less pronounced. None of the lead or LD SNPs were significant eQTLs for any gene/tissue in the GTEx v.8 database. In the Barcelona and University of Virginia genotyping and RNA sequencing (BarcUVa-Seq) dataset, a single SNP *rs12194512* [LD SNP, coefficient of determination (R^2) = 0.36] was a significant eQTL with *GFOD1*, a gene approximately 800 kb upstream of the main finding. While *rs72833769* was not specifically identified as an eQTL for *PHACTR1*, it was significantly associated with predicted *PHACTR1* expression ($P = 8.4 \times 10^{-6}$). Last, in the eQTLGEN database, we found significant eQTLs with LD SNPs *rs499627* ($R^2 = 0.32$) and *rs538788* ($R^2 = 0.24$) for the expression of *TBC1D7* (table S5A).

The regional plot for *rs350047* (5p13.1) shows that the SNP lies within a long noncoding RNA region, *LINC00604*, and is in LD with a known GWAS region identified in 2018 by Schmit *et al.* (14) (fig. S5B). This locus appears to reside in a region with little histone modifications or deoxyribonuclease (DNase) accessible sites based on evidence from CRC normal and tumor tissues and CRC cell lines (table S5B and fig. S6B). However, SNPs in LD with *rs350047* overlap with functionally active sites in connective and immune cells (17). Several LD SNPs are significant eQTLs for *PTGER4* in GTEx v.8 suprapubic epithelial cells. This finding is

Table 2. Associations between aspirin/NSAID intake and CRC risk stratified by rs72833769. Odds ratios (ORs) and 95% confidence intervals (CIs) calculated from traditional interaction model with an interaction term. Model 1: Covariates include age (continuous), sex, study, and the first three principal components. Model 2: Includes all covariates in Model 1 + smoking (never/ever), alcohol consumption (nondrinkers; moderate, 1 to 28 g/day; heavy, >28 g/day), BMI (continuous), and red meat intake (study and sex specific quartiles of red meat intake based on controls only).									
	TT			CT			CC		
	Cases	Controls	OR (95% CI)	Cases	Controls	OR (95% CI)	Cases	Controls	OR (95% CI)
Aspirin/NSAID									
Model 1									
Nonregular use	17,861	21,554	1.0 (Ref)	2,410	3,349	1.0 (Ref)	89	120	1.0 (Ref)
Regular use	9,011	14,697	0.73 (0.71–0.76)	1,374	2,073	0.94 (0.85–1.03)	61	68	1.44 (0.82–2.54)
Model 2									
Nonregular use	15,957	19,917	1.0 (Ref)	2,138	3,113	1.0 (Ref)	81	114	1.0 (Ref)
Regular use	8,159	13,423	0.72 (0.70–0.75)	1,243	1,930	0.90 (0.81, 0.99)	54	65	1.33 (0.72, 2.46)
Aspirin only									
Model 1									
Nonregular use	19,535	24,775	1.0 (Ref)	2,689	3,874	1.0 (Ref)	108	142	1.0 (Ref)
Regular use	7,134	11,229	0.76 (0.73–0.79)	1,066	1,497	1.00 (0.90, 1.11)	42	46	1.40 (0.75, 2.59)
Model 2									
Nonregular use	17,517	22,969	1.0 (Ref)	2,399	3,615	1.0 (Ref)	98	136	1.0 (Ref)
Regular use	6,427	10,180	0.76 (0.73–0.79)	958	1,384	0.96 (0.86, 1.07)	37	43	1.28 (0.66, 2.49)

Table 3. Associations between aspirin/NSAID intake and CRC risk stratified by rs350047. Odds ratios (ORs) and 95% confidence intervals (CIs) calculated from traditional interaction model with an interaction term. Model 1: Covariates include age (continuous), sex, study, and the first three principal components. Model 2: Includes all covariates in Model 1 + smoking (never/ever), alcohol consumption (nondrinkers; moderate, 1 to 28 g/day; heavy, >28 g/day), BMI (continuous), and red meat intake (study and sex specific quartiles of red meat intake based on controls only).									
	CC			CT			TT		
	Cases	Controls	OR (95% CI)	Cases	Controls	OR (95% CI)	Cases	Controls	OR (95% CI)
Aspirin/NSAID									
Model 1									
Nonregular use	5,179	6,921	1.0 (Ref)	10,141	12,453	1.0 (Ref)	5040	5649	1.0 (Ref)
Regular use	2,765	4,496	0.81 (0.76–0.87)	5,301	8,425	0.78 (0.74–0.81)	2380	3917	0.67 (0.63–0.72)
Model 2									
Nonregular use	4,630	6,410	1.0 (Ref)	9,057	11,480	1.0 (Ref)	4489	5254	1.0 (Ref)
Regular use	2,504	4,104	0.80 (0.75–0.86)	4,811	7,725	0.76 (0.72–0.80)	2141	3589	0.66 (0.61–0.71)
Aspirin only									
Model 1									
Nonregular use	5,690	7,958	1.0 (Ref)	11,157	14,340	1.0 (Ref)	5485	6493	1.0 (Ref)
Regular use	2,197	3,374	0.85 (0.79–0.91)	4,173	6,389	0.81 (0.77–0.85)	1872	3009	0.69 (0.64–0.74)
Model 2									
Nonregular use	5,107	7,394	1.0 (Ref)	9,999	13,272	1.0 (Ref)	4908	6054	1.0 (Ref)
Regular use	1,978	3,055	0.85 (0.79–0.91)	3,774	5,815	0.81 (0.77–0.85)	1670	2737	0.69 (0.64–0.74)

A *rs72833769* x Aspirin/NSAIDs



B *rs350047* x Aspirin/NSAIDs




Fig. 3. Forest plots of GxE interactions for identified SNPs and aspirin/NSAID and risk of CRC stratified by tumor molecular subtypes. ORs (dots) and 95% CIs (error bars) are plotted for each case stratified by the presence (*BRAF* mutant; CIMP-high; MSI-high; *KRAS* mutant) or absence (*BRAF* wild type; CIMP-low/negative; Non-MSI-high; *KRAS* wild type) of the marker in cases versus controls (referent) for each identified SNP: (A) *rs72833769* and (B) *rs350047*. The P value for heterogeneity (P_{het}) between estimates for cases with the molecular marker present versus absent and the overall association of the interaction for cases with molecular marker data versus controls is also provided.

corroborated when querying eQTLGEN, where essentially the entire region contains significant eQTLs for *PTGER4* in blood cells (table S5B). Similarly, *rs350047* was not an identified eQTL for *PTGER4* expression but was significantly correlated with predicted expression in our data ($P = 2.2 \times 10^{-16}$). In addition to *PTGER4*, several LD SNPs were also significant eQTLs for *DAB2*

in eQTLGEN, approximately 800 kb downstream from the lead SNP (table S5B). Last, we developed genetic models to fit interactions between regular aspirin use and predicted *PHACTR1* (upstream of *rs72833769*) and *DAB2/PTGER4* (near *rs350047*) expression based on the BarcUVa-Seq dataset but did not identify any significant interactions.

DISCUSSION

We report results of the largest genome-wide GxE interaction scan to date for CRC focused on aspirin and NSAIDs. Consistent with previous epidemiological studies, regular use of aspirin/NSAIDs was inversely associated with CRC risk in this analysis. The direct aspirin/NSAIDs associations did not differ substantially when stratified by sex or tumor site. Gene-environment interaction scans identified two previously undescribed interaction SNPs, *rs350047* (5p13.1) and *rs72833769* (6p24.1), and subgroup-specific analyses show different magnitudes of effects of aspirin/NSAIDs on CRC risk at each locus defined by genotype.

Our finding of a significant interaction for *rs350047* (5p13.1) and aspirin's preventive capacity is biologically plausible and potentially functionally relevant. *PTGER4* encodes PTGER4 (EP4), a major receptor for prostaglandin E₂ (PGE₂). PGE₂ is one of the major proinflammatory factors that is produced constitutively in the intestinal epithelium and is elevated during periods of inflammation (18, 19). Aberrant PGE₂ signaling via PTGER4 within the mucosal microenvironment promotes tumorigenesis and affects cellular differentiation processes central to mucosal injury repair (20–23). In the setting of cancer, PGE₂ has been found to induce proliferation of CRC stem cells and CRC liver metastasis in mouse models via EP4-dependent signaling pathways (24, 25). Increased synthesis of PGE₂ [due to increased prostaglandin endoperoxide synthase 2 (PTGS2) expression] is observed in patients with CRC (24, 25). Genetic deletion of *Ptgs2* or downstream PGE₂ receptors results in protection from tumorigenesis in CRC mouse models (26–30). In addition, 15-hydroxyprostaglandin dehydrogenase (*HPGD*; *15-PGDH*), the primary enzyme that catabolizes PGE₂, has been characterized as a tumor suppressor for several human cancers, including colorectal, and is ubiquitously down-regulated in CRC (31–36).

However, in the context of prevention, carefully orchestrated PGE₂-PTGER4 signaling mediated by PTGS2 in the mucosal microenvironment has been demonstrated to have a key role in healthy stem cell function and regenerative programming (20, 21, 37). The leading putative biological mechanism for the chemopreventive effects of aspirin and other nonselective NSAIDs centers on inhibition of PTGS1 and PTGS2, the enzymes that mediate conversion of arachidonic acid to PGE₂ (3). Direct inhibition of PGE₂ synthesis by aspirin has been confirmed in separate clinical trials, where aspirin intervention in patients at risk for CRC significantly reduces the major urinary metabolite of PGE₂ (38–40), 11 α -hydroxy-9,15-dioxo-2,3,4,5-tetranorprostate-1,20-dioic acid (PGE-M), an inflammatory biomarker that may predict individual risk for colorectal neoplasia and may have utility as an efficacy marker for aspirin prevention (11, 39, 41).

Thus, it is biologically plausible that the interaction SNP *rs350047* as an eQTL for *PTGER4* expression may substantially modify the PGE₂-PTGER4 signaling axis, aspirin's effect on this pathway and resultant tumorigenesis, and in turn an individual's risk for CRC. Moreover, the locus 5p13.1 has been previously implicated in investigations of cancer and inflammatory bowel disease. Genome-wide association analyses of CRC risk using data from a broader subset of the same Genetics and Epidemiology of Colorectal Cancer Consortium (GECCO)/Colon Cancer Family Registry (CCFR)/Colorectal Transdisciplinary Study (CORECT) consortia identified two independent risk loci in this region; Schmit *et al.* (14) reported a significant association between indel *rs58791712* (G/GT) and CRC risk, while Huyghe *et al.* (42) reported a separate hit upstream of this variant, (G/A). The association for *rs7708610* was genome-wide significant upon conditioning

models for the previously identified hit at *rs58791712* (using a surrogate SNP), indicating the presence of an independent susceptibility locus (42). The correlation of the SNP *rs350047* showing interaction with *rs58791712* and *rs7708610* in 1000G European population is $R^2 = 0.38$ and 0.042, respectively. Furthermore, several analyses of inflammatory diseases, specifically for, but not limited to, Crohn's disease, also implicated the 5p13.1 region as a risk locus (43–46), although none of the reported markers are in LD with our main finding.

In contrast to *rs350047*, the functional evidence for *rs72833769* (6p24.1) is less clear. However, a significant eQTL with *TBC1D7* might be of interest. This gene is a subunit of the TSC1-TSC2 complex, a key regulator of mammalian target of rapamycin (mTOR) activity, which is a proto-oncogene and downstream effector of the phosphatidylinositol 3-kinase/AKT pathway. As previously described in the context of pancreatic cancer (47), there is potential cross-talk between PGE₂ expression, PTGER4 activation, and the AKT/mTOR pathways. Moreover, in a large, recent study using human normal colon organoids, network analysis of RNA-seq data revealed that both *TBC1D7* and *GFOD1* were present within modules that were both significantly modulated by aspirin treatment in vitro and enriched for PGE₂-related pathways (48). Thus, it appears biologically plausible that aspirin/NSAID-mediated changes in PGE₂ expression can modulate the effectiveness of signal transduction to proliferative signals by these signaling axes.

These results also add to the existing evidence base for SNPs predicting aspirin's chemopreventive benefit for CRC. Nan *et al.* (13) previously reported significant interactions between aspirin use and variants in 12p12.3 (*rs2965667* and *rs10505806*) and 15q25.2 (*rs16973225*) for CRC risk in a subset of studies included in this investigation. Although these SNPs were not identified at genome-wide significance in our main analysis, we did confirm nominally significant interactions for these SNPs. While other prior studies have identified putative SNP-based biomarkers for an interaction of aspirin/NSAID use and CRC risk [summarized in prior reviews (3, 49)], the functional relevance of the SNPs to their preventive mechanism for CRC has been limited, beyond putative indirect linkages with CRC-relevant or prostaglandin-signaling adjacent pathways or enzymes linked to aspirin pharmacokinetics. Similarly, prior studies that have performed targeted genotyping of specific pathways related to aspirin/NSAID chemoprevention (e.g., prostaglandin synthesis) include important a priori signals (50–54), such that the absence of previously studied functional genetic variants from the results here does not preclude their potential utility in precision prevention. Nonetheless, our study is not only the largest study to date to perform a GWIS and, thus, has the greatest power to eliminate potential false negatives, but also by extending our findings to our functional dataset, we have demonstrated that *rs350047* is within an eQTL for *PTGER4*, implicating a mechanistic link for this SNP and the effects of NSAIDs on PGE₂-PTGER4 signaling.

Beyond host genetics, other molecular epidemiology studies have demonstrated that aspirin/NSAID chemopreventive effects may be limited to tumors that would have otherwise developed via pathways sensitive to aspirin prevention. Specifically, these prior studies demonstrate that aspirin may have greater protective effects against *BRAF* wild-type (55) and *PIK3CA* mutant CRCs (56) or those tumors with higher PTGS2 expression (57) or against CRCs arising in individuals with intact *HPGD* expression (58). Our findings that interactions between aspirin/NSAID use and *rs350047* were statistically significant only when comparing *BRAF* mutant, CIMP-high,

or MSI-high cases to controls, whereas interactions between aspirin/NSAID use and *rs72833769* were limited to cases absent of these molecular markers compared to controls, suggests that host genetics may further influence aspirin's ability to differentially prevent tumors arising via separate tumorigenesis pathways (e.g., traditional versus serrated pathways). Combined, our findings support calls for a more nuanced, precision prevention approach to specifically identify subsets of individuals likely to benefit from aspirin/NSAIDs and improve broader, one-size-fits-all recommendations [i.e., only on the basis of age or conditioning on added risk for cardiovascular disease as has been the case for past U.S. Preventive Services Task Force recommendations (59, 60)]. While some markers, such as *rs72833769*, may provide simpler, more qualitative guidance (i.e., go/no-go) for individual stratification, quantitative interaction markers, such as *rs350047*, will be critical to calibrating precision prevention recommendations to maximize net benefit among those more likely to benefit, particularly when they are linked to potential mechanisms of action. These findings can help identify interrelated modes of action that may help clarify differential anticancer effects associated with aspirin/NSAIDs, new and more effective therapeutic or prevention targets, or specific pathways individuals are on toward the development of cancer that may or may not be responsive to these agents. Moreover, quantitative measures may further help explain observed interindividual responses to preventive interventions, even among patients identified as likely to respond by qualitative measures (11). In all, a truly precise precision prevention approach likely requires the incorporation of both qualitative and quantitative genomic interaction markers of risk and response in context of the potential tumorigenesis pathways to which an individual is particularly susceptible to and additional individual risk factors.

Our study has several strengths, including being the largest GWIS of CRC and aspirin/NSAIDs to date, pairing this data with functional datasets that allow for the identification of eQTLs and using efficient and powerful statistical methods to improve power over standard GxE tests. A limitation of our study is that the resolution of data collected on aspirin use alone varied across studies and required us in some cases to group aspirin use, for which there is a clearer chemoprevention benefit established, with other non-aspirin NSAID use. While not ideal, we have performed careful data harmonization across the studies to ensure that we accurately have differentiated between aspirin/NSAID use, which may include aspirin only users along with users of both aspirin and/or other NSAIDs, and those who explicitly recorded aspirin use separately from other NSAID use. In addition, our study focuses on harmonized self-reported regular aspirin/NSAID use presumed to be representative of standard dose recommendations (≤ 325 mg/day) as this was most consistently collected across all studies. Although multiple lines of evidence support the notion that the chemopreventive association with aspirin is strongest 5 to 10 years after commencing continued regular use, we did not have sufficient information to consider the duration of exposure. Despite this measurement error, which is likely to attenuate risk estimates and lead to reduced power, we were able to show strong protective associations for aspirin/NSAIDs and identify two significant interactions. Nonetheless, future studies will need to examine whether the interactions observed for aspirin/NSAID use are specific to use of aspirin or extend to other non-aspirin NSAIDs and in context of other key factors including dose and duration of use. Our analysis is limited to individuals of European ancestry, thereby limiting the direct extension of these findings to different racial and

ethnic populations. Last, we cannot rule out the effects of residual confounding, including from additional CRC risk factors (e.g., inflammatory bowel disease history) not otherwise accounted for in our analysis, or recall bias in our analysis.

In summary, we identified previously undescribed genetic loci that modify the protective effect of regular aspirin/NSAID use on CRC risk. Functional evidence presented in our investigation implicates genes directly involved in CRC-associated signaling pathways, such as PGE₂ synthesis/signaling in the case of *PTGER4*, and downstream pathways involved in tumorigenesis and proposed to be central to aspirin's protective mode of action, suggesting biological plausibility. Validation and additional functional work are necessary to confirm these findings. Furthermore, the likelihood of identifying additional interaction loci in the future can be improved via implementation of tissue-/cell-specific functional annotations, along with multi-ethnic GWIS.

MATERIALS AND METHODS

Study design

We pooled individual level genomic and epidemiological data from three consortia comprising individuals of European ancestry—the GECCO, the CORECT, and the CCFR consortia comprising a total of 52 studies. Study details have been previously published (14, 42, 61) and can be found in table S6. For cohort studies, nested case-control sets were assembled via risk-set sampling, while population-based controls were used for case-control studies. Clinical trials were treated as cohort studies and participants were matched according to trial arm. Cases were defined as CRC or advanced adenomas and were confirmed by medical records, pathological reports, or death certificate information. Controls were matched on age, sex, race, and enrollment date/trial group, when applicable. For the small subset of advanced adenoma cases, matched controls were found to be polyp-free on sigmoidoscopy or colonoscopy at the time of adenoma selection. All participants gave written informed consent, and studies were approved by their respective Institutional Review Boards.

Exposure and aspirin/NSAID use ascertainment

Analyses include individuals with complete exposure and covariate information. We excluded individuals based on discrepancies between reported and genotypic sex, cryptic relatedness, and duplicates. For any individual included in multiple studies, we selected a single record for them, prioritizing the study that genotyped on the more comprehensive platform. Collection of risk factor data and harmonization across contributing studies has been previously described (62). Combined use of any aspirin or non-aspirin NSAIDs at reference time is defined as “aspirin/NSAID use” (yes or no), comprising a final study sample size of 72,667 individuals (30,806 cases and 41,861 controls). When aspirin use was specifically queried separately from NSAID use, we defined regular use of aspirin at reference time as “aspirin only use” (yes or no) to conduct analyses limited to aspirin users ($N = 72,137$; 30,574 cases and 41,563 controls). Multivariate models include age (continuous), sex (male/female), study, smoking status (never/ever), alcohol consumption (nondrinkers; moderate, 1 to 28 g/day; heavy, >28 g/day), BMI (continuous), and red meat intake (study and sex-specific quartiles of red meat intake based on controls only) as specifically denoted. Individuals with missing covariate data were excluded from any model using the covariate.

Genotyping

The genotyping platforms used in each study are summarized in table S6. Briefly, genotyped SNPs were excluded on the basis of call rate (<95 to 98%), lack of Hardy-Weinberg equilibrium ($P < 1 \times 10^{-4}$), and discordant calls between duplicates. All autosomal SNPs of all studies were imputed to the Haplotype Reference Consortium r1.1 (2016) reference panel via the Michigan Imputation Server (63) and converted into a binary format for data management and analyses using R package BinaryDosage. Imputed common SNPs were filtered on the basis of a pooled minor allele frequency (MAF) ≥ 0.01 and imputation accuracy $R^2 > 0.8$. After imputation and quality control analyses, a total of more than 7.2 million SNPs were available for analysis. Principal components analysis for population stratification assessment was performed using PLINK 1.9 on 30,000 randomly sampled imputed SNPs with MAF > 0.05 and imputation accuracy $R^2 > 0.99$. Additional details on genotyping and quality control have been previously published (42).

Statistical analysis

We evaluated associations of CRC risk with regular aspirin/NSAID use and with aspirin only use and CRC risk using random-effects meta-analysis of study specific results to obtain summary ORs and 95% CIs across studies. Association tests were stratified by study design, sex, and tumor subsite. The latter was categorized into the following groups based on the following ICD-9 codes (otherwise excluded): proximal colon (153.0, 153.1, 153.4, 153.6), distal colon (153.2, 153.3, 153.7), and rectum (154.0, 154.1).

Common variant analysis

Genome-wide scans were conducted using GxEScanR, an R package that implements several methods for detecting GxE interactions (<https://CRAN.R-project.org/package=GxEScanR>). Imputed allelic dosages were modeled as continuous variables. Our analysis employs three primary methods: (i) traditional 1-df GxE logistic regression models, (ii) a two-step method using “D versus G” and “E versus G” joint information [the EDGE method (64)] as the step 1 filtering statistic and the 1-df GxE statistic for the step 2 test accounting for LD-based correlation among SNPs in step 2 (65), and (iii) a 3-df joint test that incorporates information from main effects, gene-exposure associations, and GxE interaction statistics in a single model. (66) For the 3-df test, we report only results with GxE P value $< 1 \times 10^{-4}$, since this test captures markers with significant results from any of the three sources of statistics. We adjusted the overall genome-wide significance threshold for each testing procedure to $5 \times 10^{-8}/2.5$ to account for multiple testing with methods that are statistically correlated. We adopt the following notation: E, regular aspirin/NSAID use; G, SNP; D, CRC outcome; and C, set of adjustment covariates. Traditional logistic regression models assessed interactions on a multiplicative scale by including an interaction term in the model $\text{logit}[\text{Pr}(D = 1|G, E, C)] = \beta_0 + \beta_G G + \beta_E E + \beta_{G \times E} G \times E + \beta_C C$, testing $H_0 : \beta_{G \times E} = 0$. The typical focus of a GWIS is on the 1-df test of GxE interaction based on the null hypothesis $H_0 : \beta_{G \times E} = 0$. This test is known to generally have low power, particularly in the context of discovering new interactions in a GWIS. To enhance our power to discover new GxE loci, we used the two-step EDGE method and 3-df joint tests summarized above. Additional details of these approaches are in the Supplementary Materials.

Functional follow-up

We leveraged the BarcUVa-Seq (<https://barcuvaseq.org/>) resource (67) to develop prediction models to test interactions between predicted gene expression and regular aspirin/NSAID use, in a method previously explored in GECCO consortium using GTEx data (68). Weights were generated using elastic net regularized regression models fit on BarcUVa-Seq gene expression (20,693 measured gene expressions) and HRC imputed genotypes. Imputed SNPs were filtered on the basis of MAF > 0.1 and imputation quality ($R^2 > 0.7$); models were adjusted for age, sex, RNA-seq batch, tissue location, principal components, and probabilistic estimation of expression residuals (PEER) factors (69). Weights were then used to predict gene expression in our study sample of 72,667 subjects. In total, 13,393 gene expressions were successfully predicted. Interaction tests were conducted using logistic regression with an interaction term between predicted genetic expression and aspirin/NSAID use and aspirin use alone.

We used LocusZoom v1.3 (70) to generate regional plots for significant findings to inspect and extend the association signal and LD, and position of findings relative to genes in the region. Measures of LD were estimated using European populations from the 1000 Genomes Project. The putative functional role of these SNPs and those in LD ($R^2 > 0.2$) at 500-kb flanking regions was investigated with relation to their potential contribution to gene expression regulation in two ways: first, by their physical location in regions of chromatin accessibility or histone modifications (variant enhancer loci) and, second, through their direct association with expression of nearby genes (eQTLs).

To assess the physical location of the SNPs in regions of chromatin accessibility or histone modifications, we queried overlaps between our findings and regions containing active enhancer elements in tissue from healthy and tumor colon samples in addition to CRC cell lines, obtained from previously analyzed assays for transposase-accessible chromatin with sequencing (ATAC-seq) data, DNase hypersensitivity sequencing, and H3K27ac histone chromatin immunoprecipitation sequencing datasets. (15) We extended this analysis to include additional tissue types by incorporating regulatory annotations of histone modifications from 10 groups of tissues, obtained from several resources (71, 72) and compiled by Finucane *et al.* (16) Furthermore, we queried lead and LD SNPs against functional annotation databases from ENSEMBL using the Variant Effect Predictor tool. (73)

We checked for eQTLs using several resources: (i) GTEx v8, (ii) the colon transcriptome explorer (CoTrEx 2.0; <https://barcuvaseq.org/cotrex/>, accessed May 2021), a resource for transcriptomic data jointly developed by the University of BarcUVa-Seq, which includes eQTL from 445 epithelium-enriched healthy colon biopsies from ascending, transverse, and descending colon, and 3) eQTL results from eQTLGEN, a consortium of 37 cohorts pooling RNA sequencing data from whole blood samples.

Supplementary Materials

This PDF file includes:

Supplementary Materials and Methods
Sources of consortium funding
Figs. S1 to S6
Tables S1 to S6
References

REFERENCES AND NOTES

- Y. Qiao, T. Yang, Y. Gan, W. Li, C. Wang, Y. Gong, Z. Lu, Associations between aspirin use and the risk of cancers: A meta-analysis of observational studies. *BMC Cancer* **18**, 288 (2018).
- S. Friis, A. H. Riis, R. Erichsen, J. A. Baron, H. T. Sorensen, Low-dose aspirin or nonsteroidal anti-inflammatory drug use and colorectal cancer risk: A population-based, case-control study. *Ann. Intern. Med.* **163**, 347–355 (2015).
- D. A. Drew, Y. Cao, A. T. Chan, Aspirin and colorectal cancer: The promise of precision chemoprevention. *Nat. Rev. Cancer* **16**, 173–186 (2016).
- J. Burn, H. Sheth, F. Elliott, L. Reed, F. Macrae, J. P. Mecklin, G. Moslein, F. E. McDonald, L. Bertario, D. G. Evans, A. M. Gerdes, J. W. C. Ho, A. Lindblom, P. J. Morrison, J. Rashbass, R. Ramesar, T. Seppala, H. J. W. Thomas, K. Pylvanainen, G. M. Borthwick, J. C. Mathers, D. T. Bishop, C. Investigators, Cancer prevention with aspirin in hereditary colorectal cancer (Lynch syndrome), 10-year follow-up and registry-based 20-year data in the CAPP2 study: A double-blind, randomised, placebo-controlled trial. *Lancet* **395**, 1855–1863 (2020).
- N. R. Cook, I. M. Lee, S. M. Zhang, M. V. Moorthy, J. E. Buring, Alternate-day, low-dose aspirin and cancer risk: Long-term observational follow-up of a randomized trial. *Ann. Intern. Med.* **159**, 77–85 (2013).
- J. A. Baron, B. F. Cole, R. S. Sandler, R. W. Haile, D. Ahnen, R. Bresalier, G. McKeown-Eyssen, R. W. Summers, R. Rothstein, C. A. Burke, D. C. Snover, T. R. Church, J. I. Allen, M. Beach, G. J. Beck, J. H. Bond, T. Byers, E. R. Greenberg, J. S. Mandel, N. Marcon, L. A. Mott, L. Pearson, F. Saibil, R. U. van Stolk, A randomized trial of aspirin to prevent colorectal adenomas. *N. Engl. J. Med.* **348**, 891–899 (2003).
- R. S. Sandler, S. Halabi, J. A. Baron, S. Budinger, E. Paskett, R. Keresztes, N. Petrelli, J. M. Pipas, D. D. Karp, C. L. Loprinzi, G. Steinbach, R. Schilsky, A randomized trial of aspirin to prevent colorectal adenomas in patients with previous colorectal cancer. *N. Engl. J. Med.* **348**, 883–890 (2003).
- R. F. Logan, M. J. Grainge, V. C. Shepherd, N. C. Armitage, K. R. Muir, ukCAP Trial Group, Aspirin and folic acid for the prevention of recurrent colorectal adenomas. *Gastroenterology* **134**, 29–38 (2008).
- R. Benamouzig, J. Deyra, A. Martin, B. Girard, E. Jullian, B. Piednoir, D. Couturier, T. Coste, J. Little, S. Chaussade, Daily soluble aspirin and prevention of colorectal adenoma recurrence: One-year results of the APACC trial. *Gastroenterology* **125**, 328–336 (2003).
- H. Ishikawa, M. Mutoh, S. Suzuki, S. Tokudome, Y. Saida, T. Abe, S. Okamura, M. Tajika, T. Joh, S. Tanaka, S. E. Kudo, T. Matsuda, M. Iimuro, T. Yukawa, T. Takayama, Y. Sato, K. Lee, S. Kitamura, M. Mizuno, Y. Sano, N. Gondo, K. Sugimoto, M. Kusunoki, C. Goto, N. Matsura, T. Sakai, K. Wakabayashi, The preventive effects of low-dose enteric-coated aspirin tablets on the development of colorectal tumours in Asian patients: A randomised trial. *Gut* **63**, 1755–1759 (2014).
- D. A. Drew, A. T. Chan, Aspirin in the prevention of colorectal neoplasia. *Annu. Rev. Med.* **72**, 415–430 (2021).
- K. McAllister, L. E. Mechanic, C. Amos, H. Aschard, I. A. Blair, N. Chatterjee, D. Conti, W. J. Gauderman, L. Hsu, C. M. Hutter, M. M. Jankowska, J. Kerr, P. Kraft, S. B. Montgomery, B. Mukherjee, G. J. Papanicolaou, C. J. Patel, M. D. Ritchie, B. R. Ritz, D. C. Thomas, P. Wei, J. S. Witte, Current challenges and new opportunities for gene-environment interaction studies of complex diseases. *Am. J. Epidemiol.* **186**, 753–761 (2017).
- H. Nan, C. M. Hutter, Y. Lin, E. J. Jacobs, C. M. Ulrich, E. White, J. A. Baron, S. I. Berndt, H. Brenner, K. Butterbach, B. J. Caan, P. T. Campbell, C. S. Carlson, G. Casey, J. Chang-Claude, S. J. Chanock, M. Cotterchio, D. Duggan, J. C. Figueiredo, C. S. Fuchs, E. L. Giovannucci, J. Gong, R. W. Haile, T. A. Harrison, R. B. Hayes, M. Hoffmeister, J. L. Hopper, T. J. Hudson, M. A. Jenkins, S. Jiao, N. M. Lindor, M. Lemire, L. Le Marchand, P. A. Newcomb, S. Ogino, B. M. Pflugeisen, J. D. Potter, C. Qu, S. A. Rosse, A. Rudolph, R. E. Schoen, F. R. Schumacher, D. Seminara, M. L. Slattery, S. N. Thibodeau, F. Thomas, M. Thornquist, G. S. Warnick, B. W. Zanke, W. J. Gauderman, U. Peters, L. Hsu, A. T. Chan, Association of aspirin and NSAID use with risk of colorectal cancer according to genetic variants. *JAMA* **313**, 1133–1142 (2015).
- S. L. Schmit, C. K. Edlund, F. R. Schumacher, J. Gong, T. A. Harrison, J. R. Huyghe, C. Qu, M. Melas, D. J. Van Den Berg, H. Wang, S. Tring, S. J. Plummer, D. Albanes, M. H. Alonso, C. I. Amos, K. Anton, A. K. Aragaki, V. Arndt, E. L. Barry, S. I. Berndt, S. Bezieau, S. Bien, A. Bloomer, J. Boehm, M. C. Boutrou-Ruault, H. Brenner, S. Brezina, D. D. Buchanan, K. Butterbach, B. J. Caan, P. T. Campbell, C. S. Carlson, J. E. Castella, A. T. Chan, J. Chang-Claude, S. J. Chanock, I. Cheng, Y. W. Cheng, L. S. Chin, J. M. Church, T. Church, G. A. Coetzee, M. Cotterchio, M. C. Correa, K. R. Curtis, D. Duggan, D. F. Easton, D. English, E. J. M. Feskens, R. Fischer, L. M. FitzGerald, B. K. Fortini, L. G. Fritsche, C. S. Fuchs, M. Gago-Dominguez, M. Gala, S. J. Gallinger, W. J. Gauderman, G. G. Giles, E. L. Giovannucci, S. M. Gogarten, C. Gonzalez-Villalpando, E. M. Gonzalez-Villalpando, W. M. Grady, J. K. Greenon, A. Gsur, M. Gunter, C. A. Haiman, J. Hampe, S. Harlid, J. F. Harju, R. B. Hayes, P. Hofer, M. Hoffmeister, J. L. Hopper, S. C. Huang, J. M. Huerta, T. J. Hudson, D. J. Hunter, G. E. Idos, M. Iwasaki, R. D. Jackson, E. J. Jacobs, S. H. Jee, M. A. Jenkins, W. H. Jia, S. Jiao, A. D. Joshi, L. N. Kolonel, S. Kono, C. Kooperberg, V. Krogh, T. Kuehn, S. Kury, A. LaCroix, C. A. Laurie, F. Lejbkowitz, M. Lemire, H. J. Lenz, D. Levine, C. I. Li, L. Li, W. Lieb, Y. Lin, N. M. Lindor, Y. R. Liu, F. Loupakis, Y. Lu, F. Luh, J. Ma, C. Mancao, F. J. Manion, S. D. Markowitz, V. Martin, K. Matsuda, K. Matsuo, K. J. McDonnell, C. E. McNeil, R. Milne, A. J. Molina, B. Mukherjee, N. Murphy, P. A. Newcomb, K. Offit, H. Omichessan, D. Palli, J. P. P. Cotore, J. Perez-Mayoral, P. D. Pharoah, J. D. Potter, C. Qu, L. Raskin, G. Rennert, H. S. Rennert, B. M. Riggs, C. Schafmayer, R. E. Schoen, T. A. Sellers, D. Seminara, G. Severi, W. Shi, D. Shibata, X. O. Shu, E. M. Siegel, M. L. Slattery, M. Southey, Z. K. Stadler, M. C. Stern, S. Stintzing, D. Taverna, S. N. Thibodeau, D. C. Thomas, A. Trichopoulos, S. Tsugane, C. M. Ulrich, F. J. B. van Duijnhoven, B. van Guelpan, J. Vijai, J. Virtamo, S. J. Weinstein, E. White, A. K. Win, A. Wolk, M. Woods, A. H. Wu, K. Wu, Y. B. Xiang, Y. Yen, B. W. Zanke, Y. X. Zeng, B. Zhang, N. Zubair, S. S. Kweon, J. C. Figueiredo, W. Zheng, L. L. Marchand, A. Lindblom, V. Moreno, U. Peters, G. Casey, L. Hsu, D. V. Conti, S. B. Gruber, Novel common genetic susceptibility loci for colorectal cancer. *J. Natl. Cancer Inst.* **111**, 146–157 (2019).
- A. J. Cohen, A. Saikhova, O. Corradin, J. M. Luppino, K. Lovrenert, C. F. Bartels, J. J. Morrow, S. C. Mack, G. Dhillon, L. Beard, L. Myeroff, M. F. Kalady, J. Willis, J. E. Bradner, R. A. Keri, N. A. Berger, S. M. Pruet-Miller, S. D. Markowitz, P. C. Scacheri, Hotspots of aberrant enhancer activity punctuate the colorectal cancer epigenome. *Nat. Commun.* **8**, 14400 (2017).
- H. K. Finucane, B. Bulik-Sullivan, A. Gusev, G. Trynka, Y. Reshef, P. R. Loh, V. Anttila, H. Xu, C. Zang, K. Farh, S. Ripke, F. R. Day, ReproGen Consortium, Schizophrenia Working Group of the Psychiatric Genomics Consortium, The RBC Consortium, S. Purcell, E. Stahl, S. Lindstrom, J. R. Perry, Y. Okada, S. Raychaudhuri, M. J. Daly, N. Patterson, B. M. Neale, A. L. Price, Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nat. Genet.* **47**, 1228–1235 (2015).
- H. K. Finucane, Y. A. Reshef, V. Anttila, K. Slowikowski, A. Gusev, A. Byrnes, S. Gazal, P. R. Loh, C. Lareau, N. Shores, G. Genovese, A. Saunders, E. Macosko, S. Pollack, C. Brainstorm, J. R. B. Perry, J. D. Buenrostro, B. E. Bernstein, S. Raychaudhuri, S. McCarroll, B. M. Neale, A. L. Price, Heritability enrichment of specifically expressed genes identifies disease-relevant tissues and cell types. *Nat. Genet.* **50**, 621–629 (2018).
- T. O. Ishikawa, H. R. Herschman, Conditional knockout mouse for tissue-specific disruption of the cyclooxygenase-2 (Cox-2) gene. *Genesis* **44**, 143–149 (2006).
- N. A. Manieri, M. R. Mack, M. D. Himmelrich, D. L. Worthley, E. M. Hanson, L. Eckmann, T. C. Wang, T. S. Stappenbeck, Mucosally transplanted mesenchymal stem cells stimulate intestinal healing by promoting angiogenesis. *J. Clin. Invest.* **125**, 3606–3618 (2015).
- H. Miyoshi, K. L. VanDussen, N. P. Malvin, S. H. Ryu, Y. Wang, N. M. Sonnek, C. W. Lai, T. S. Stappenbeck, Prostaglandin E2 promotes intestinal repair through an adaptive cellular response of the epithelium. *EMBO J.* **36**, 5–24 (2017).
- M. Roulis, A. Kalamanos, M. Scherthanner, P. Bielecki, J. Zhao, E. Kaffe, L. S. Frommelt, R. Qu, M. S. Knapp, A. Henriques, N. Chalkidi, V. Koliarakis, J. Jiao, J. R. Brewer, M. Bacher, H. N. Blackburn, X. Zhao, R. M. Breyer, Y. Aidinis, D. Jain, B. Su, H. R. Herschman, Y. Kluger, G. Kollias, R. A. Flavell, Paracrine orchestration of intestinal tumorigenesis by a mesenchymal niche. *Nature* **580**, 524–529 (2020).
- U. Jain, C. W. Lai, S. Xiong, V. M. Goodwin, Q. Lu, B. D. Muegge, G. P. Christophi, K. L. VanDussen, B. P. Cummings, E. Young, J. Hambor, T. S. Stappenbeck, Temporal regulation of the bacterial metabolite deoxycholate during colonic repair is critical for crypt regeneration. *Cell Host Microbe* **24**, 353–363.e5 (2018).
- D. C. Montrose, M. Nakanishi, R. C. Murphy, S. Zarin, J. P. McAleer, A. T. Vella, D. W. Rosenberg, The role of PGE2 in intestinal inflammation and tumorigenesis. *Prostaglandins Other Lipid Mediat.* **116–117**, 26–36 (2015).
- K. Bellamkonda, N. K. Chandrashekar, J. Osman, B. C. Selvaranesan, S. Savari, A. Sjolander, The eicosanoids leukotriene D4 and prostaglandin E2 promote the tumorigenicity of colon cancer-initiating cells in a xenograft mouse model. *BMC Cancer* **16**, 425 (2016).
- D. Wang, L. Fu, H. Sun, L. Guo, R. N. DuBois, Prostaglandin E2 promotes colorectal cancer stem cell expansion and metastasis in Mice. *Gastroenterology* **149**, 1884–1895.e4 (2015).
- P. C. Chulada, M. B. Thompson, J. F. Mahler, C. M. Doyle, B. W. Gaul, C. Lee, H. F. Tian, S. G. Morham, O. Smithies, R. Langenbach, Genetic disruption of Ptg-1, as well as Ptg-2, reduces intestinal tumorigenesis in Min mice. *Cancer Res.* **60**, 4705–4708 (1999).
- M. Mutoh, K. Watanabe, T. Kitamura, Y. Shoji, M. Takahashi, T. Kawamori, K. Tani, M. Kobayashi, T. Maruyama, K. Kobayashi, S. Ohuchida, Y. Sugimoto, S. Narumiya, T. Sugimura, K. Wakabayashi, Involvement of prostaglandin E receptor subtype EP(4) in colon carcinogenesis. *Cancer Res.* **62**, 28–32 (2002).
- M. Oshima, J. E. Dinchuk, S. L. Kargman, H. Oshima, B. Hancock, E. Kwong, J. M. Trzaskos, J. F. Evans, M. M. Taketo, Suppression of intestinal polyposis in Apc Δ 716 knockout mice by inhibition of cyclooxygenase 2 (COX-2). *Cell* **87**, 803–809 (1996).
- M. Sonoshita, K. Takaku, N. Sasaki, Y. Sugimoto, F. Ushikubi, S. Narumiya, M. Oshima, M. M. Taketo, Acceleration of intestinal polyposis through prostaglandin receptor EP2 in Apc(Delta 716) knockout mice. *Nat. Med.* **7**, 1048–1051 (2001).
- K. Watanabe, T. Kawamori, S. Nakatsugi, T. Ohta, S. Ohuchida, H. Yamamoto, T. Maruyama, K. Kondo, F. Ushikubi, S. Narumiya, T. Sugimura, K. Wakabayashi, Role of the prostaglandin E receptor subtype EP1 in colon carcinogenesis. *Cancer Res.* **59**, 5093–5096 (1999).
- M. G. Backlund, J. R. Mann, V. R. Holla, F. G. Buchanan, H. H. Tai, E. S. Musiek, G. L. Milne, S. Katkuri, R. N. DuBois, 15-Hydroxyprostaglandin dehydrogenase is down-regulated in colorectal cancer. *J. Biol. Chem.* **280**, 3217–3223 (2005).

32. M. Yan, R. M. Rerko, P. Platzer, D. Dawson, J. Willis, M. Tong, E. Lawrence, J. Lutterbaugh, S. Lu, J. K. Willson, G. Luo, J. Hensold, H. H. Tai, K. Wilson, S. D. Markowitz, 15-Hydroxyprostaglandin dehydrogenase, a COX-2 oncogene antagonist, is a TGF-beta-induced suppressor of human gastrointestinal cancers. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 17468–17473 (2004).
33. S. J. Myung, R. M. Rerko, M. Yan, P. Platzer, K. Guda, A. Dotson, E. Lawrence, A. J. Dannenberg, A. K. Lovgren, G. Luo, T. P. Pretlow, R. A. Newman, J. Willis, D. Dawson, S. D. Markowitz, 15-Hydroxyprostaglandin dehydrogenase is an in vivo suppressor of colon tumorigenesis. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 12098–12102 (2006).
34. M. Yan, S. J. Myung, S. P. Fink, E. Lawrence, J. Lutterbaugh, P. Yang, X. Zhou, D. Liu, R. M. Rerko, J. Willis, D. Dawson, H. H. Tai, J. S. Barnholtz-Sloan, R. A. Newman, M. M. Bertagnoli, S. D. Markowitz, 15-Hydroxyprostaglandin dehydrogenase inactivation as a mechanism of resistance to celecoxib chemoprevention of colon tumors. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 9409–9413 (2009).
35. C. L. Thompson, S. P. Fink, J. D. Lutterbaugh, R. C. Elston, M. L. Veigl, S. D. Markowitz, L. Li, Genetic variation in 15-hydroxyprostaglandin dehydrogenase and colon cancer susceptibility. *PLOS ONE* **8**, e64122 (2013).
36. H. R. Roberts, H. J. Smartt, A. Greenhough, A. E. Moore, A. C. Williams, C. Paraskeva, Colon tumour cells increase PGE(2) by regulating COX-2 and 15-PGDH to promote survival during the microenvironmental stress of glucose deprivation. *Carcinogenesis* **32**, 1741–1747 (2011).
37. Y. R. Na, D. Jung, M. Stakenborg, H. Jang, G. J. Gu, M. R. Jeong, S. Y. Suh, H. J. Kim, Y. H. Kwon, T. S. Sung, S. B. Ryou, K. J. Park, J. P. Im, J. Y. Park, Y. S. Lee, H. Han, B. Park, S. Lee, D. Kim, H. S. Lee, I. Cleynen, G. Matteoli, S. H. Seok, Prostaglandin E(2) receptor PTGER4-expressing macrophages promote intestinal epithelial barrier regeneration upon inflammation. *Gut* **70**, 2249–2260 (2021).
38. V. Fedirko, P. T. Bradshaw, J. C. Figueiredo, R. S. Sandler, E. L. Barry, D. J. Ahnen, G. L. Milne, R. S. Bresalier, J. A. Baron, Urinary metabolites of prostanoids and risk of recurrent colorectal adenomas in the Aspirin/Folate Polyp Prevention Study (AFPPS). *Cancer Prev. Res. (Phila.)* **8**, 1061–1068 (2015).
39. D. A. Drew, M. M. Schuck, M. V. Magicheva-Gupta, K. O. Stewart, K. K. Gilpin, P. Miller, M. P. Parziale, E. N. Pond, O. Takacs-Nagy, D. C. Zerjav, S. M. Chin, J. Mackinnon Krems, D. Meixell, A. D. Joshi, W. Ma, F. P. Colizzo, P. J. Carolan, N. S. Nishioka, K. Staller, J. M. Richter, H. Khalili, M. K. Gala, J. J. Garber, D. C. Chung, J. C. Yarz, L. Zukerberg, G. Petrucci, B. Rocca, C. Patrono, G. L. Milne, M. Wang, A. T. Chan, Effect of low-dose and standard-dose aspirin on pge(2) biosynthesis among individuals with colorectal adenomas: A randomized clinical trial. *Cancer Prev. Res.* **13**, 877–888 (2020).
40. L. L. Garland, J. Guillen-Rodriguez, C. H. Hsu, M. Yozwiak, H. H. Zhang, D. S. Alberts, L. E. Davis, E. Szabo, C. Merenstein, J. Lei, X. Zhang, H. Liu, G. Liu, A. E. Spira, J. E. Beane, M. Wojtowicz, H. S. Chow, Effect of intermittent versus continuous low-dose aspirin on nasal epithelium gene expression in current smokers: A randomized, double-blinded trial. *Cancer Prev. Res.* **12**, 809–820 (2019).
41. N. Bezawada, M. Song, K. Wu, R. S. Mehta, G. L. Milne, S. Ogino, C. S. Fuchs, E. L. Giovannucci, A. T. Chan, Urinary PGE-M levels are associated with risk of colorectal adenomas and chemopreventive response to anti-inflammatory drugs. *Cancer Prev. Res. (Phila.)* **7**, 758–765 (2014).
42. J. R. Huyghe, S. A. Bien, T. A. Harrison, H. M. Kang, S. Chen, S. L. Schmit, D. V. Conti, C. Qu, J. Jeon, C. K. Edlund, P. Greenside, M. Wainberg, F. R. Schumacher, J. D. Smith, D. M. Levine, S. C. Nelson, N. A. Sinnott-Armstrong, D. Albanes, M. H. Alonso, K. Anderson, C. Arnaud-Collell, V. Arndt, C. Bamia, B. L. Banbury, J. A. Baron, S. I. Berndt, S. Bezieau, D. T. Bishop, J. Boehm, H. Boeheim, H. Brenner, S. Brezina, S. Buch, D. D. Buchanan, A. Burnett-Hartman, K. Butterbach, B. J. Caan, P. T. Campbell, C. S. Carlson, S. Castellvi-Bel, A. T. Chan, J. Chang-Claude, S. J. Chanock, M. D. Chirlaque, S. H. Cho, C. M. Connolly, A. J. Cross, K. Cuk, K. R. Curtis, A. de la Chapelle, K. F. Doheny, D. Duggan, D. F. Easton, S. G. Elias, F. Elliott, D. R. English, E. J. M. Feskens, J. C. Figueiredo, R. Fischer, L. M. FitzGerald, D. Forman, M. Gala, S. Gallinger, W. J. Gauderman, G. G. Giles, E. Gillanders, J. Gong, P. J. Goodman, W. M. Grady, J. S. Grove, A. Gsur, M. J. Gunter, R. W. Haile, J. Hampe, H. Hampel, S. Harlid, R. B. Hayes, P. Hofer, M. Hoffmeister, J. L. Hopper, W. H. Hsu, W. Y. Huang, T. J. Hudson, D. J. Hunter, G. Ibanez-Sanz, G. E. Idos, R. Ingersoll, R. D. Jackson, E. J. Jacobs, M. A. Jenkins, A. D. Joshi, C. E. Joshi, T. O. Keku, T. J. Key, H. R. Kim, E. Kobayashi, L. N. Kolonel, C. Kooperberg, T. Kuhn, S. Kury, S. S. Kweon, S. C. Larsson, C. A. Laurie, L. Le Marchand, S. M. Leal, S. C. Lee, F. Lejbkowitz, M. Lemire, C. I. Li, L. Li, W. Lieb, Y. Lin, A. Lindblom, N. M. Lindor, H. Ling, T. L. Louie, S. Mannisto, S. D. Markowitz, V. Martin, G. Masala, C. E. McNeil, M. Melas, R. L. Milne, L. Moreno, N. Murphy, R. Myte, A. Naccarati, P. A. Newcomb, K. Offit, S. Ogino, N. C. Onland-Moret, B. Pardini, P. S. Parfrey, R. Pearlman, V. Perduca, P. D. P. Pharoah, M. Pinchev, E. A. Platz, R. L. Prentice, E. Pugh, L. Raskin, G. Rennert, H. S. Rennert, E. Riboli, M. Rodriguez-Barranco, J. Romm, L. C. Sakoda, C. Schafmayer, R. E. Schoen, D. Seminara, M. Shah, T. Sheldford, M. H. Shin, K. Shulman, S. Sieri, M. L. Slattery, M. C. Southey, Z. K. Stadler, C. Stegmaier, Y. R. Su, C. M. Tang, S. N. Thibodeau, D. C. Thomas, S. S. Thomas, A. E. Toland, A. Trichopoulos, M. G. T. Van Den Berg, F. J. B. van Duijnhoven, B. Van Guelpen, H. van Kranen, J. Vijai, K. Visvanathan, P. Vodicka, L. Vodickova, V. Vymetalkova, K. Weigl, S. J. Weinstein, E. White, A. K. Win, C. R. Wolf, A. Wolk, M. O. Woods, A. H. Wu, S. H. Zaidi, B. W. Zanke, Q. Zhang, W. Zheng, P. C. Scacheri, J. D. Potter, M. C. Bassik, A. Kundaje, G. Casey, V. Moreno, G. R. Abecasis, D. A. Nickerson, S. B. Gruber, L. Hsu, U. Peters, Discovery of common and rare genetic risk variants for colorectal cancer. *Nat. Genet.* **51**, 76–87 (2019).
43. A. Franke, J. Hampe, P. Rosenstiel, C. Becker, F. Wagner, R. Hasler, R. D. Little, K. Huse, A. Ruether, T. Balschun, M. Wittig, A. Elsharawy, G. Mayr, M. Albrecht, N. J. Prescott, C. M. Onnie, H. Fournier, T. Keith, U. Radelof, M. Platzer, C. G. Mathew, M. Stoll, M. Krawczak, P. Nurnberg, S. Schreiber, Systematic association mapping identifies NELL1 as a novel IBD disease gene. *PLOS ONE* **2**, e691 (2007).
44. C. Libioulle, E. Louis, S. Hansoul, C. Sandor, F. Farnir, D. Franchimont, S. Vermeire, O. Dewit, M. de Vos, A. Dixon, B. Demarche, I. Gut, S. Heath, M. Foglio, L. Liang, D. Laukens, M. Mni, D. Zelenika, A. Van Gossom, P. Rutgeerts, J. Belaiche, M. Lathrop, M. Georges, Novel Crohn disease locus identified by genome-wide association maps to a gene desert on 5p13.1 and modulates expression of PTGER4. *PLOS Genet.* **3**, e58 (2007).
45. J. Glas, J. Seiderer, D. Czamara, G. Pasciuto, J. Diegelmann, M. Wetzel, T. Olszak, C. Wolf, B. Muller-Myhsok, T. Balschun, J. P. Achkar, M. I. Kamboh, A. Franke, R. H. Duerr, S. Brand, PTGER4 expression-modulating polymorphisms in the 5p13.1 region predispose to Crohn's disease and affect NF-κB and XBP1 binding sites. *PLOS ONE* **7**, e52873 (2012).
46. J. Z. Liu, S. van Sommeren, H. Huang, S. C. Ng, R. Alberts, A. Takahashi, S. Ripke, J. C. Lee, L. Jostins, T. Shah, S. Abadian, J. H. Cheon, J. Cho, N. E. Dayani, L. Franke, Y. Fuyuno, A. Hart, R. C. Juyal, G. Juyal, W. H. Kim, A. P. Morris, H. Poustchi, W. G. Newman, V. Midha, T. R. Orchard, H. Vahedi, A. Sood, J. Y. Sung, R. Malekzadeh, H. J. Westra, K. Yamazaki, S. K. Yang, The International Multiple Sclerosis Genetics Consortium, The International IBD Genetics Consortium, J. C. Barrett, B. Z. Alizadeh, M. Parkes, T. Bk, M. J. Daly, M. Kubo, C. A. Anderson, R. K. Weersma, Association analyses identify 38 susceptibility loci for inflammatory bowel disease and highlight shared genetic risk across populations. *Nat. Genet.* **47**, 979–986 (2015).
47. H. H. Chang, S. H. Young, J. Sinnett-Smith, C. E. Chou, A. Moro, K. M. Hertz, O. J. Hines, E. Rozengurt, G. Eibl, Prostaglandin E2 activates the mTORC1 pathway through an EP4/cAMP/PKA- and EP1/Ca2+-mediated mechanism in the human pancreatic carcinoma cell line PANC-1. *Am. J. Physiol. Cell Physiol.* **309**, C639–C649 (2015).
48. M. A. M. Devall, D. A. Drew, C. H. Dampier, S. J. Plummer, S. Eaton, J. Bryant, V. Diez-Obrero, J. Mo, D. Kedrin, D. C. Zerjav, O. Takacs-Nagy, L. T. Jennelle, M. W. Ali, O. H. Yilmaz, V. Moreno, S. M. Powell, A. T. Chan, U. Peters, G. Casey, Transcriptome-wide in vitro effects of aspirin on patient-derived normal colon organoids. *Cancer Prev. Res. (Phila.)* **14**, 1089–1100 (2021).
49. C. M. Ulrich, J. Bigler, J. D. Potter, Non-steroidal anti-inflammatory drugs for cancer prevention: Promise, perils and pharmacogenetics. *Nat. Rev. Cancer* **6**, 130–140 (2006).
50. E. M. Poole, L. Hsu, L. Xiao, R. J. Kulmacz, C. S. Carlson, P. S. Rabinovitch, K. W. Makar, J. D. Potter, C. M. Ulrich, Genetic variation in prostaglandin E2 synthesis and signaling, prostaglandin dehydrogenase, and the risk of colorectal adenoma. *Cancer Epidemiol. Biomarkers Prev.* **19**, 547–557 (2010).
51. C. M. Ulrich, J. Bigler, R. Sparks, J. Whitton, J. G. Sibert, E. L. Goode, Y. Yasui, J. D. Potter, Polymorphisms in PTGS1 (=COX-1) and risk of colorectal polyps. *Cancer Epidemiol. Biomarkers Prev.* **13**, 889–893 (2004).
52. C. M. Ulrich, J. Whitton, J. H. Yu, J. Sibert, R. Sparks, J. D. Potter, J. Bigler, PTGS2 (COX-2) -765G > C promoter variant reduces risk of colorectal adenoma among nonusers of nonsteroidal anti-inflammatory drugs. *Cancer Epidemiol. Biomarkers Prev.* **14**, 616–619 (2005).
53. B. L. Seufert, E. M. Poole, J. Whitton, L. Xiao, K. W. Makar, P. T. Campbell, R. J. Kulmacz, J. A. Baron, P. A. Newcomb, M. L. Slattery, J. D. Potter, C. M. Ulrich, IκBκ and NFκB1, NSAID use and risk of colorectal cancer in the Colon Cancer Family Registry. *Carcinogenesis* **34**, 79–85 (2013).
54. K. W. Makar, E. M. Poole, A. J. Resler, B. Seufert, K. Curtin, S. E. Kleinstein, D. Duggan, R. J. Kulmacz, L. Hsu, J. Whitton, C. S. Carlson, C. F. Rimorin, B. J. Caan, J. A. Baron, J. D. Potter, M. L. Slattery, C. M. Ulrich, COX-1 (PTGS1) and COX-2 (PTGS2) polymorphisms, NSAID interactions, and risk of colon and rectal cancers in two independent populations. *Cancer Causes Control* **24**, 2059–2075 (2013).
55. R. Nishihara, P. Lochhead, A. Kuchiba, S. Jung, M. Yamauchi, X. Liao, Y. Imamura, Z. R. Qian, T. Morikawa, M. Wang, D. Spiegelman, E. Cho, E. Giovannucci, C. S. Fuchs, A. T. Chan, S. Ogino, Aspirin use and risk of colorectal cancer according to BRAF mutation status. *JAMA* **309**, 2563–2571 (2013).
56. X. Liao, P. Lochhead, R. Nishihara, T. Morikawa, A. Kuchiba, M. Yamauchi, Y. Imamura, Z. R. Qian, Y. Baba, K. Shima, R. Sun, K. Noshio, J. A. Meyerhardt, E. Giovannucci, C. S. Fuchs, A. T. Chan, S. Ogino, Aspirin use, tumor PIK3CA mutation, and colorectal-cancer survival. *N. Engl. J. Med.* **367**, 1596–1606 (2012).
57. A. T. Chan, S. Ogino, C. S. Fuchs, Aspirin and the risk of colorectal cancer in relation to the expression of COX-2. *N. Engl. J. Med.* **356**, 2131–2142 (2007).
58. S. P. Fink, M. Yamauchi, R. Nishihara, S. Jung, A. Kuchiba, K. Wu, E. Cho, E. Giovannucci, C. S. Fuchs, S. Ogino, S. D. Markowitz, A. T. Chan, Aspirin and the risk of colorectal cancer in relation to the expression of 15-hydroxyprostaglandin dehydrogenase (HPGD). *Sci. Transl. Med.* **6**, 233re2 (2014).

59. K. Bibbins-Domingo, U.S. Preventive Services Task Force, Aspirin use for the primary prevention of cardiovascular disease and colorectal cancer: U.S. preventive services task force recommendation statement. *Ann. Intern. Med.* **164**, 836–845 (2016).
60. J. M. Guirguis-Blake, C. V. Evans, L. A. Perdue, S. I. Bean, C. A. Senger, Aspirin use to prevent cardiovascular disease and colorectal cancer: Updated evidence report and systematic review for the US preventive services task force. *JAMA* **327**, 1585–1597 (2022).
61. F. R. Schumacher, S. L. Schmit, S. Jiao, C. K. Edlund, H. Wang, B. Zhang, L. Hsu, S. C. Huang, C. P. Fischer, J. F. Harju, G. E. Idos, F. Lejbkowitz, F. J. Manion, K. McDonnell, C. E. McNeil, M. Melas, H. S. Rennett, W. Shi, D. C. Thomas, D. J. Van Den Berg, C. M. Hutter, A. K. Aragaki, K. Butterbach, B. J. Caan, C. S. Carlson, S. J. Chanock, K. R. Curtis, C. S. Fuchs, M. Gala, E. L. Giovannucci, S. M. Gogarten, R. B. Hayes, B. Henderson, D. J. Hunter, R. D. Jackson, L. N. Kolonel, C. Kooperberg, S. Kury, A. LaCroix, C. C. Laurie, C. A. Laurie, M. Lemire, D. Levine, J. Ma, K. W. Makar, C. Qu, D. Taverna, C. M. Ulrich, K. Wu, S. Kono, D. W. West, S. I. Berndt, S. Bezieau, H. Brenner, P. T. Campbell, A. T. Chan, J. Chang-Claude, G. A. Coetzee, D. V. Conti, D. Duggan, J. C. Figueiredo, B. K. Fortini, S. J. Gallinger, W. J. Gauderman, G. Giles, R. Green, R. Haile, T. A. Harrison, M. Hoffmeister, J. L. Hopper, T. J. Hudson, E. Jacobs, M. Iwasaki, S. H. Jee, M. Jenkins, W. H. Jia, A. Joshi, L. Li, N. M. Lindor, K. Matsuo, V. Moreno, B. Mukherjee, P. A. Newcomb, J. D. Potter, L. Raskin, G. Rennert, S. Rosse, G. Severi, R. E. Schoen, D. Seminara, X. O. Shu, M. L. Slattery, S. Tsugane, E. White, Y. B. Xiang, B. W. Zanke, W. Zheng, L. Le Marchand, G. Casey, S. B. Gruber, U. Peters, Genome-wide association study of colorectal cancer identifies six new susceptibility loci. *Nat. Commun.* **6**, 7138 (2015).
62. C. M. Hutter, J. Chang-Claude, M. L. Slattery, B. M. Pflugeisen, Y. Lin, D. Duggan, H. Nan, M. Lemire, J. Rangrej, J. C. Figueiredo, S. Jiao, T. A. Harrison, Y. Liu, L. S. Chen, D. L. Stelling, G. S. Warnick, M. Hoffmeister, S. Kury, C. S. Fuchs, E. Giovannucci, A. Hazra, P. Kraft, D. J. Hunter, S. Gallinger, B. W. Zanke, H. Brenner, B. Frank, J. Ma, C. M. Ulrich, E. White, P. A. Newcomb, C. Kooperberg, A. Z. LaCroix, R. L. Prentice, R. D. Jackson, R. E. Schoen, S. J. Chanock, S. I. Berndt, R. B. Hayes, B. J. Caan, J. D. Potter, L. Hsu, S. Bezieau, A. T. Chan, T. J. Hudson, U. Peters, Characterization of gene-environment interactions for colorectal cancer susceptibility loci. *Cancer Res.* **72**, 2036–2044 (2012).
63. S. Das, L. Forer, S. Schönherr, C. Sidore, A. E. Locke, A. Kwong, S. I. Vrieze, E. Y. Chew, S. Levy, M. McGue, D. Schlessinger, D. Stambolian, P. R. Loh, W. G. Iacono, A. Swaroop, L. J. Scott, F. Cucca, F. Kronenberg, M. Boehnke, G. R. Abecasis, C. Fuchsberger, Next-generation genotype imputation service and methods. *Nat. Genet.* **48**, 1284–1287 (2016).
64. W. J. Gauderman, P. Zhang, J. L. Morrison, J. P. Lewinger, Finding novel genes by testing $G \times E$ interactions in a genome-wide association study. *Genet. Epidemiol.* **37**, 603–613 (2013).
65. E. S. Kawaguchi, A. E. Kim, J. P. Lewinger, W. J. Gauderman, Improved two-step testing of genome-wide gene-environment interactions. *Genet. Epidemiol.* **47**, 152–166 (2023).
66. W. J. Gauderman, A. Kim, D. V. Conti, J. Morrison, D. C. Thomas, H. Vora, J. P. Lewinger, A unified model for the analysis of gene-environment interaction. *Am. J. Epidemiol.* **188**, 760–767 (2019).
67. V. Diez-Obrero, C. H. Dampier, F. Moratalla-Navarro, M. Devall, S. J. Plummer, A. Diez-Villanueva, U. Peters, S. Bien, J. R. Huyghe, A. Kundaje, G. Ibanez-Sanz, E. Guino, M. Obon-Santacana, R. Carreras-Torres, G. Casey, V. Moreno, Genetic effects on transcriptome profiles in colon epithelium provide functional insights for genetic risk loci. *Cell. Mol. Gastroenterol. Hepatol.* **12**, 181–197 (2021).
68. X. Wang, Y. R. Su, P. S. Petersen, S. Bien, S. L. Schmit, D. A. Drew, D. Albanes, S. I. Berndt, H. Brenner, P. T. Campbell, G. Casey, J. Chang-Claude, S. J. Gallinger, S. B. Gruber, R. W. Haile, T. A. Harrison, M. Hoffmeister, E. J. Jacobs, M. A. Jenkins, A. D. Joshi, L. Li, Y. Lin, N. M. Lindor, L. L. Marchand, V. Martin, R. Milne, R. MacInnis, V. Moreno, H. Nan, P. A. Newcomb, J. D. Potter, G. Rennert, H. Rennert, M. L. Slattery, S. N. Thibodeau, S. J. Weinstein, M. O. Woods, A. T. Chan, E. White, L. Hsu, U. Peters, Exploratory genome-wide interaction analysis of nonsteroidal anti-inflammatory drugs and predicted gene expression on colorectal cancer risk. *Cancer Epidemiol. Biomarkers Prev.* **29**, 1800–1808 (2020).
69. O. Stegle, L. Parts, R. Durbin, J. Winn, A Bayesian framework to account for complex non-genetic factors in gene expression levels greatly increases power in eQTL studies. *PLoS Comput. Biol.* **6**, e1000770 (2010).
70. R. J. Pruim, R. P. Welch, S. Sanna, T. M. Teslovich, P. S. Chines, T. P. Giedt, M. Boehnke, G. R. Abecasis, C. J. Willer, LocusZoom: Regional visualization of genome-wide association scan results. *Bioinformatics* **26**, 2336–2337 (2010).
71. D. Hnisz, B. J. Abraham, T. I. Lee, A. Lau, V. Saint-Andre, A. A. Sigova, H. A. Hoke, R. A. Young, Super-enhancers in the control of cell identity and disease. *Cell* **155**, 934–947 (2013).
72. G. Trynka, C. Sandor, B. Han, H. Xu, B. E. Stranger, X. S. Liu, S. Raychaudhuri, Chromatin marks identify critical cell types for fine mapping complex trait variants. *Nat. Genet.* **45**, 124–130 (2013).
73. W. McLaren, L. Gil, S. E. Hunt, H. S. Riat, G. R. Ritchie, A. Thormann, P. Flicek, F. Cunningham, The ensemble variant effect predictor. *Genome Biol.* **17**, 122 (2016).
74. P. Kraft, Y. C. Yen, D. O. Stram, J. Morrison, W. J. Gauderman, Exploiting gene-environment interaction to detect genetic associations. *Hum. Hered.* **63**, 111–119 (2007).
75. C. E. Murray, J. P. Lewinger, W. J. Gauderman, Gene-environment interaction in genome-wide association studies. *Am. J. Epidemiol.* **169**, 219–226 (2009).
76. W. J. Gauderman, B. Mukherjee, H. Aschard, L. Hsu, J. P. Lewinger, C. J. Patel, J. S. Witte, C. Amos, C. G. Tai, D. Conti, D. G. Torgerson, S. Lee, N. Chatterjee, Update on the State of the Science for Analytical Methods for Gene-Environment Interactions. *Am. J. Epidemiol.* **186**, 762–770 (2017).
77. C. Kooperberg, M. Leblanc, Increasing the power of identifying gene x gene interactions in genome-wide association studies. *Genet. Epidemiol.* **32**, 255–263 (2008).
78. I. Ionita-Laza, M. B. McQueen, N. M. Laird, C. Lange, Genomewide weighted hypothesis testing in family-based association studies, with an application to a 100K scan. *Am. J. Hum. Genet.* **81**, 607–614 (2007).
79. J. P. Lewinger, E. S. Kawaguchi, W. J. Gauderman, A note on p-value multiple-testing adjustment for two-step genome-wide gene-environment interactions scans. medRxiv, 2023.2006.2027.23291946 [Preprint] (2023). <https://doi.org/10.1101/2023.06.27.23291946>.

Acknowledgments: ASTERISK: We are very grateful to B. Buecher without whom this project would not have existed. We also thank all those who agreed to participate in this study, including the patients and the healthy control persons, as well as all the physicians, technicians, and students. CCFR: The Colon CFR graciously thanks the generous contributions of their study participants, dedication of study staff, and the financial support from the U.S. National Cancer Institute, without which this important registry would not exist. CLUE II: We thank the participants of Clue II and appreciate the continued efforts of the staff at the Johns Hopkins George W. Comstock Center for Public Health Research and Prevention in the conduct of the Clue II Cohort Study. Cancer data were provided by the Maryland Cancer Registry, Center for Cancer Prevention and Control, Maryland Department of Health, with funding from the State of Maryland and the Maryland Cigarette Restitution Fund. The collection and availability of cancer registry data are also supported by the Cooperative Agreement NUS8DP006333, funded by the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services. CPS-II: We express sincere appreciation to all Cancer Prevention Study-II participants and to each member of the study and biospecimen management group. We would like to acknowledge the contribution to this study from central cancer registries supported through the Centers for Disease Control and Prevention's National Program of Cancer Registries and cancer registries supported by the National Cancer Institute's Surveillance Epidemiology and End Results Program. We assume full responsibility for all analyses and interpretation of results. The views expressed here are those of the authors and do not necessarily represent the American Cancer Society or the American Cancer Society–Cancer Action Network. DACHS: We thank all participants and cooperating clinicians, and everyone who provided excellent technical assistance. EDRN: We acknowledge all contributors to the development of the resource at University of Pittsburgh School of Medicine, Department of Gastroenterology, Department of Pathology, Hepatology and Nutrition and Biomedical Informatics. Harvard cohorts: The study protocol was approved by the institutional review boards of the Brigham and Women's Hospital and Harvard T.H. Chan School of Public Health, and those of participating registries as required. We acknowledge Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital as home of the NHS. We would like to acknowledge the contribution to this study from central cancer registries supported through the Centers for Disease Control and Prevention's National Program of Cancer Registries (NPCR) and/or the National Cancer Institute's Surveillance, Epidemiology, and End Results (SEER) Program. Central registries may also be supported by state agencies, universities, and cancer centers. Participating central cancer registries include the following: Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Indiana, Iowa, Kentucky, Louisiana, Massachusetts, Maine, Maryland, Michigan, Mississippi, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Puerto Rico, Rhode Island, Seattle SEER Registry, South Carolina, Tennessee, Texas, Utah, Virginia, West Virginia, and Wyoming. We assume full responsibility for analyses and interpretation of these data. Kentucky: We would like to acknowledge the staff at the Kentucky Cancer Registry. LCCS: We acknowledge the contributions of Jennifer Barrett, Robin Waxman, Gillian Smith and Emma Northwood in conducting this study. NCCCS I & II: We would like to thank the study participants, and the NC Colorectal Cancer Study staff. PLCO: We thank the PLCO Cancer Screening Trial screening center investigators and the staffs from Information Management Services Inc. and Westat Inc. We also thank the study participants for their contributions that made this study possible. Cancer incidence data have been provided by the District of Columbia Cancer Registry, Georgia Cancer Registry, Hawaii Cancer Registry, Minnesota Cancer Surveillance System, Missouri Cancer Registry, Nevada Central Cancer Registry, Pennsylvania Cancer Registry, Texas Cancer Registry, Virginia Cancer Registry, and Wisconsin Cancer Reporting System. All are supported in part by funds from the Center for Disease Control and Prevention; National Program for Central Registries; local states or by the National Cancer Institute; Surveillance, Epidemiology, and End Results program. The results reported here and the conclusions derived are the sole responsibility of the authors. PPS3 and PPS4 Polyp Prevention Study trials: We would like to

thank the study participants, investigators, and staff. **SEARCH:** We thank the SEARCH team. **SELECT:** We thank the research and clinical staff at the sites that participated on SELECT study, without whom the trial would not have been successful. We are also grateful to the 35,533 dedicated men who participated in SELECT. **WHI:** The authors thank the WHI investigators and staff for their dedication, and the study participants for making the program possible. A full listing of WHI investigators can be found at [www-whi-org.s3.us-west-2.amazonaws.com/wp-content/uploads/WHI-Investigator-Long-List.pdf](http://www.whi-org.s3.us-west-2.amazonaws.com/wp-content/uploads/WHI-Investigator-Long-List.pdf). **Disclaimer:** Where authors are identified as personnel of the International Agency for Research on Cancer/World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy or views of the International Agency for Research on Cancer/World Health Organization. **Funding:** Individual authors report the following grants and/or funding support: D.A.D.: K01 DK120742; A.E.K.: R01CA201407; J.M.: R01CA201407, P01CA196569; R01CA273198; J.P.L.: R01CA201407, P01CA196569; R01CA273198; E.K.: R01CA201407, P01CA196569; R01CA273198; Y.F.: P01CA196569; R01CA273198; E.L.B.: R01CA059005 and R01CA098286; S.I.B.: Intramural Research Program, Division of Cancer Epidemiology and Genetics, NCI, NIH; J.C.F.: R01 CA155101; S.B.G.: U19 CA148107, R01 CA263318; M.A.J.: U01 CA167551, U01 CA122839, R01 CA143247, U19 CA148107, R01 CA81488, U19 CA148107; B.M.L.: VCA MCRF18005; J.O.: R03CA270473, R01CA207371, R01CA189184, R01CA254108, and U01CA206110; C.M.U.: R03CA270473, R01CA207371, R01CA189184, R01CA254108, U01CA206110, and P30CA042014; U.P.: R01CA059045, U01CA164930, R01CA244588, R01CA201407, and R01 CA273198; W.J.G.: R01CA201407, P01CA196569; R01CA273198; and Genetics and Epidemiology of Colorectal Cancer Consortium (GECCO): National Cancer Institute, National Institutes of Health, U.S. Department of Health and Human Services (R01 CA059045, U01 CA164930, R01 CA244588, R01 CA201407). Genotyping/Sequencing services were provided by the Center for Inherited Disease Research (CIDR) contract number HHSN268201700006I and HHSN268201200008I. This research was funded in part through the NIH/NCI Cancer Center Support Grant P30 CA015704. Scientific Computing Infrastructure at Fred Hutch was funded by ORIP grant S10OD028685. Additional funding supporting the GECCO Consortium and participating studies is listed in the Supplemental Materials. **Author contributions:** Conceptualization: D.A.D., Y.L., J.P.L., S.A.B., N.D., H.B., P.T.C., J.C.-C., S.B.G., M.J.G., R.N., D.C.T., K.T.T., V.M., U.P., A.T.C., and W.J.G. Methodology: D.A.D., A.E.K., Y.L., J.P.L., E.K., N.D., J.B., H.B., A.B., P.T.C., D.V.C., R.C.-T., S.B.G., M.J.G., R.N., D.C.T., C.M.U., L.H., U.P., and W.J.G. Investigation: D.A.D., A.E.K., Y.L., Y.F., J.B., J.D.P., H.B. A.B.-H., P.T.C., G.C., D.V.C., S.B.G., A.G.,

M.J.G., P.A.N., M.O.-S., S.O., R.K.P., B.P., P.C.S., C.M.U., U.P., and A.T.C. Visualization: D.A.D., A.E.K., Y.L., A.H., and W.J.G. Resources: D.A.D., D.A., A.W., R.L.P., V.A., E.L.B., S.I.B., H.B., A.B.-H., P.T.C., G.C., J.C.-C., J.C.F., S.B.G., A.G., M.H., J.R.H., M.A.J., A.K., L.L.M., L.L., B.L., P.A.N., C.C.N., S.O., A.J.P., E.A.P., R.K.P., P.C.S., R.E.S., M.C.S., C.M.U., C.Y.U., E.W., V.M., A.T.C., and W.J.G. Software: A.E.K., Y.L., J.M., J.P.L., E.K., Y.F., J.B., D.V.C., D.C.T., and W.J.G. Formal analysis: D.A.D., A.E.K., C.Q., J.M., J.P.L., E.K., Y.F., N.Z., V.D.-O., J.B., D.V.C., S.B.G., M.J.G., J.R.H., S.L.S., and W.J.G. Data curation: D.A.D., A.E.K., Y.L., C.Q., V.A., S.B.G., T.A.H., M.H., J.R.H., E.W., V.M., and W.J.G. Validation: D.A.D., A.E.K., H.B., A.B., A.G., R.N., J.O., and W.J.G. Funding acquisition: D.D.B., H.B., P.T.C., D.V.C., S.B.G., M.J.G., M.H., M.A.J., A.K., L.L.M., R.N., P.A.N., E.A.P., D.C.T., C.M.U., E.W., V.M., U.P., A.T.C., and W.J.G. Project administration: D.A.D., S.A.B., S.B.G., L.L., M.H., E.W., U.P., A.T.C., and W.J.G. Supervision: D.A.D., V.M., U.P., A.T.C., and W.J.G. Writing—original draft: D.A.D., A.E.K., and W.J.G. Writing—review and editing: All authors. **Competing interests:** E.K. owns stock in Abbvie (ABBV) and Pfizer (PFE). J.W.B. is an owner and employee of BioRealm LLC. S.A.B. owns stock and is an employee of Adaptive Biotechnologies. S.B.G. has equity in Brogent International LLC not related to this work. K.A.J. has received personal fees from Bristol-Myers Squibb outside the submitted work and reports grants from the National Cancer Institute. A.K. is on the scientific advisory board of PatchBio, TensorBio, SerImmune, and OpenTargets; was a consultant with Illumina; and owns shares in DeepGenomics, ImmunAI, and Freenome. C.M.U. has as cancer center director oversight over research funded by several pharmaceutical companies but has not received funding directly herself. U.P. was a consultant with Abbvie and her husband is holding individual stocks for the following companies: BioNTech SE-ADR, Amazon, CureVac BV, NanoString Technologies, Google/Alphabet Inc. Class C, NVIDIA Corp, and Microsoft Corp. All other authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Summary-level data for the GWAS datasets are available through the GWAS catalog (accession no. GCST012876). For individual-level data, datasets are deposited in dbGaP (accession nos. phs001415.v1.p1, phs001315.v1.p1, phs001078.v1.p1, phs001903.v1.p1, phs001856.v1.p1, and phs001045.v1.p1).

Submitted 14 August 2023

Accepted 26 April 2024

Published 29 May 2024

10.1126/sciadv.adk3121