

Article begins on page three of this document.

<b>Title</b>	Hospital admissions for cardiovascular complications of people with or without diabetes, Victoria, 2004–2016
--------------	--

**Authors:**

	Title	First name	Mid initials	Last name	Postnom (eg, PhD)	Position1	Address1	Position2	Address2	Tel	Email
1	Ms.	Katerina	V	Kiburg	Bachelor of Biomedicine, Master of Public Health	PhD Candidate	1	PhD Candidate	2	+61 3 9231 3574	katerina.kiburg@svha.org.au
2	Assoc. Prof.	Andrew	I	Maclsaac	MBBS, MD, FRACP, FCSANZ	Cardiologist	1		2	0417 390 251	andrew.macisaac@svha.org.au
3	Prof.	Andrew		Wilson	MBBS, FRACP, PhD	Cardiologist	1		2		andrew.wilson@svha.org.au
4	Prof.	Vijaya		Sundararajan	BA (Honours), MD, M.D., MPH, MPH	Researcher	2	Head of Public Health	3		v.sundararajan@latrobe.edu.au
5	Prof.	Richard	J	Maclsaac	BSc(Hons), PhD, MBBS, FRACP	Director of Endocrinology and Professorial Fellow, University of Melbourne	1		2		r.macisaac@unimelb.edu.au

Number of corresponding author:	1
Number of alternative corresponding author:	

**Addresses:**

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/MJA2.51101](https://doi.org/10.1002/MJA2.51101)

	Institution	Postal address	City	State	Post Code	Nation
1	St Vincent's Hospital Melbourne	Level 4, Daly Wing	Melbourne	VIC	3065	
2	The University of Melbourne	35 Victoria Parade	Melbourne	VIC	3102	
3	La Trobe University	Plenty Road & Kingsbury Drive	Melbourne	VIC	3086	
4						
5						

Postal address of first corresponding author (if different from the institutional address given above)	
--	--

Primary Keywords [Office use only]	Endocrine system diseases; Cardiovascular diseases; Nervous system diseases
Secondary keywords [Office use only]	Diabetes complications; Coronary disease; Acute coronary syndrome; Stroke; Heart failure; Diabetes mellitus, type 1; Diabetes mellitus, type 2
Notes:	

**Article details** (press ctrl – 9 to enter details):

Article type	Research letter
Blurb	Admissions of people with diabetes for acute myocardial infarction, stroke, and heart failure declined, but rates remain high
Pullquote	

**Office use**

<i>Ms. Number</i>	mja20.01417.R3
<i>Medical editor</i>	Robyn Godding
<i>Medical editor email</i>	rgodding@mja.com.au
<i>Structural editor</i>	Paul Foley
<i>Structural editor email</i>	pfoley@mja.com.au
<i>Section/Category</i>	Research letter
<i>Strapheading</i>	Research letter
<i>Substrap</i>	

**Elsevier – file data:**

Filename for copyediting	kib_mja20.01417_ms.docx
Accompanying graphics	kib_mja20.01417_gr1.eps; kib_mja20.01417_gr1.jpg
Stock images	None
Appendices	kib_mja2.00000-sup-0001-supinfo <b>Title: Supplementary methods</b>

**Office use – history:**

Event	Date
Original submission received	07/08/2020

Event	Date
Accept	03/03/2021

Proof sent to author	
Proof returned by author	
Published (date format xx/xx/xx)	19/07/21
Issue	2
Vol	215
DOI	10.5694/mja2 0.01417
Journal	The Medical Journal of Australia
Original article DOI (for response)	

Author Manuscript

# Hospital admissions for cardiovascular complications of people with or without diabetes, Victoria, 2004–2016

Intensive metabolic control reduces the incidence and progression of diabetes-related micro- and macrovascular complications.<sup>1,2</sup> Nevertheless, the risk of developing cardiovascular disease is higher for people with diabetes,<sup>3</sup> and cardiovascular disease incidence rates are generally declining more rapidly for people with diabetes than for other people.<sup>4,5</sup>

We analysed hospital discharge data from the Victorian Admitted Episode Dataset<sup>6</sup> for 1 January 1999 – 31 December 2016. We identified incident cases of three cardiovascular disease complications (acute myocardial infarction [AMI], stroke, and heart failure) by International Statistical Classification of Diseases, tenth revision, Australian modification codes. Data for 1999–2003 were examined to ensure that admissions during the observation period (2004–2016) were index admissions for the specific complication, but were not included in our main analysis. Admission rates were separately calculated for people with type 1 or type 2 diabetes (numbers of people with diagnosed diabetes, by year, were obtained from the National Diabetes Services Scheme, which captures 80–90% of diabetes diagnoses<sup>7</sup>) and for people without diabetes (derived from Australian Bureau of Statistics census data<sup>8</sup>).

We analysed changes in admission rates by Joinpoint regression (<https://surveillance.cancer.gov/joinpoint>); points at which changes in the direction or magnitude of linear trends were statistically significant ( $P < 0.05$ ) were determined with permutation tests. Each trend segment was described by an annual percentage change (APC), and the change for the entire study period as the mean APC (further details: online Supporting Information). The study was approved by the St Vincent's Hospital Melbourne Human Research Ethics Committee (HREC/18/SVHM/146).

A total of 382 107 patients were admitted to Victorian hospitals during 2004–2016 with cardiovascular complications: 278 991 without diabetes (73%), 3645 with type 1 diabetes (1%), and 99 471 with type 2 diabetes (26%). AMI admission rates declined during this period for people with type 1 (mean APC,  $-7.7\%$ ; 95% confidence interval [CI],  $-13.4\%$  to  $-1.5\%$ ) or type 2 diabetes (mean APC,  $-11.4\%$ ; 95% CI,  $-13.0\%$  to  $-9.9\%$ ), as well as for people without diabetes (mean APC,  $-5.0\%$ ; 95% CI,  $-6.7\%$  to  $-3.4\%$ ) (Box 1, Box 2).

Stroke admission rates declined significantly during 2004–2016 for people with type 1

diabetes (mean APC, -7.2%; 95% CI, -12.2% to -1.9%); for people with type 2 diabetes, rates declined during 2005–2011 and 2014–2016, but not during 2011–2014 (overall change: -11.9%; 95% CI, -17.0% to -6.5%). For patients without diabetes, the decline during 2005–2014 was significant (mean APC, -4.1%; 95% CI, -5.8% to -2.3%), but not during 2015–2016 (Box 1, Box 2).

Admissions for heart failure declined during 2004–2016 for people with type 1 diabetes (mean APC, -10.3%; 95% CI, -14.1% to -6.4%) or type 2 diabetes (mean APC, -9.2%; 95% CI, -11.0% to -7.3%), and also for people without diabetes (mean APC, -2.8%; 95% CI, -4.1% to -1.5%) (Box 1, Box 2).

As hospital discharge coding data do not include information on metabolic control or medication use, we could not assess whether cardiovascular risk factor modification and use of specific medications were associated with changes in admission rates. We also lacked information on disease duration for patients with hospital-coded diabetes. Further, we have counted admissions of any patients who had presented with complications before 1998 (ie, outside our 5-year clearance period) as incident admissions; these patients would be at very high risk of further admissions, and their inclusion may have inflated the admission rates we report for the observation period of our study.

Few recent studies have assessed outcomes for all three cardiovascular complications in a single investigation. Cardiovascular complication-related admissions to Victorian hospitals declined during 2004–2016 more rapidly for people with diabetes than for those without diabetes. The relatively greater absolute decline in the numbers of admissions of people with diabetes may be related to the fact that they are considered to be at high risk for cardiovascular disease and are therefore treated more aggressively; the scope for reducing risk with multifactorial target-driven interventions is greater in these patients. Nevertheless, admission rates for cardiovascular complications of people with diabetes remain relatively high.

**Acknowledgements:** Some analysed data were provided by the National Diabetes Services Scheme, an Australian Government initiative administered by Diabetes Australia. Katerina Kiburg is supported by an Australian Government Research Training Program Scholarship, St Vincent's Institute of Medical Research and by the Research Endowment Fund, St Vincent's Hospital Melbourne.

**Competing interests:** No relevant disclosures.

Received 7 August 2020, accepted 3 March 2021

## Author details

Katerina V Kiburg<sup>1,2</sup>

Andrew I Maclsaac<sup>1,2</sup>

Andrew Wilson<sup>1,2</sup>

Vijaya Sundararajan<sup>2,3,\*</sup>

Richard J Maclsaac<sup>1,2,\*</sup>

1 St Vincent's Hospital Melbourne, Melbourne, VIC.

2 The University of Melbourne, Melbourne, VIC.

3 La Trobe University, Melbourne, VIC.


\* Equal senior authors.

## References

- 1 Rawshani A, Rawshani A, Franzén S, et al. Risk factors, mortality, and cardiovascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2018; 379: 633-644.
- 2 Gaede P, Pedersen O. Target intervention against multiple-risk markers to reduce cardiovascular disease in patients with type 2 diabetes. *Ann Med* 2004; 36: 355-366.
- 3 Cheng YJ, Imperatore G, Geiss LS, et al. Trends and disparities in cardiovascular mortality among US adults with and without self-reported diabetes, 1988–2015. *Diabetes Care* 2018; 41: 2306-2315.
- 4 Gregg EW, Li Y, Wang J, et al. Changes in diabetes-related complications in the United States, 1990–2010. *N Engl J Med* 2014; 370: 1514-1523.
- 5 Burrows NR, Li Y, Gregg EW, Geiss LS. Declining rates of hospitalization for selected cardiovascular disease conditions among adults aged ≥ 35 years with diagnosed diabetes, US, 1998–2014. *Diabetes Care* 2018; 41: 293-302.
- 6 Victorian Department of Health and Human Services. Victorian Admitted Episodes Dataset. Updated July 2020. <https://www2.health.vic.gov.au/hospitals-and-health-services/data-reporting/health-data-standards-systems/data-collections/vaed> (viewed Aug 2020).
- 7 National Diabetes Service Scheme. Diabetes data snapshots. <https://www.ndss.com.au/about-the-ndss/diabetes-facts-and-figures/diabetes-data-snapshots> (viewed Aug 2020).
- 8 Australian Bureau of Statistics. Historical population. <https://www.abs.gov.au/statistics/people/population/historical-population> (viewed Aug 2020).

**Box 1. Age- and sex-adjusted admission rates for cardiovascular complications (with 95% confidence intervals), Victoria, 2004–2016, by diabetes status of**

Author Manuscript



patients

**Box 2. Annual percentage change (APC) in admissions for cardiovascular complications, Victoria, 2004–2016, by diabetes status**

Cardiovascular complication and diabetes status	Admissions	Change in event rate, 2004–2016*		Change in event rate, by period*
		Overall change (95% CI)	Mean APC (95% CI%)	Mean APC (95% CI)
<b>Acute myocardial infarction</b>				
No diabetes	114 965	–32.7% (–32.9% to –23.5%)	–5.0% (–6.7% to –3.4%)	—
Type 1 diabetes	1272	–8.3% (–9.9% to –7.2%)	–7.7% (–13.4% to –1.5%)	1. 2005–2009: +7.0% (–9.7% to +22.8%) 2. 2009–2016: –15.1% (–21.3% to –8.7%)
Type 2 diabetes	15 278	–222% (–223% to –221%)	–11.4% (–13.0% to –9.9%)	—
<b>Stroke</b>				
No diabetes	52 320	–11.7% (–11.8% to –11.7%)	–1.7% (–4.9% to +1.5%)	1. 2005–2014: –4.1% (–5.8% to –2.3%) 2. 2014–2016: +9.6% (–10.2% to +33.8%)
Type 1 diabetes	504	–80.0% (–99.5% to –70.2%)	–7.2% (–12.2% to –1.9%)	—
Type 2 diabetes	17 440	–212% (–212% to –212%)	–11.9% (–17.0% to –6.5%)	1. 2005–2011: –14.7% (–17.6% to –11.7%) 2. 2011–2014: +5.8% (–19.0% to +38.2%) 3. 2014–2016: –26.1% (–39.8% to –9.2%)
<b>Heart failure</b>				
No diabetes	135 524	–28.7% (–28.9% to –28.6%)	–2.8% (–4.1% to –1.5%)	—
Type 1 diabetes	1393	–122% (–141% to –110%)	–10.3% (–14.1% to –6.4%)	—
Type 2 diabetes	52 831	–207% (–207% to –206%)	–9.2% (–11.0% to –7.3%)	—

CI = confidence interval.

\* Adjusted for age and sex. Event rates for 2004 and 2016 are included in the expanded version of this table in the online [Supporting Information](#).