



Minerva Access is the Institutional Repository of The University of Melbourne

**Author/s:**

Sahle, BW;Ofori-Asenso, R;Renzaho, AMN

**Title:**

ADVERSE CHILDHOOD EXPERIENCES SHOULD BE PRIORITY IN GLOBAL RESPONSE TO COVID-19

**Date:**

2020-10-01

**Citation:**

Sahle, B. W., Ofori-Asenso, R. & Renzaho, A. M. N. (2020). ADVERSE CHILDHOOD EXPERIENCES SHOULD BE PRIORITY IN GLOBAL RESPONSE TO COVID-19. *Journal of Paediatrics and Child Health*, 56 (10), pp.1656-1657. <https://doi.org/10.1111/jpc.15132>.

**Persistent Link:**

<https://hdl.handle.net/11343/276232>

## Adverse childhood experiences should be priority in global response to COVID-19

Berhe W. Sahle <sup>a</sup>, Richard Ofori-Asenso <sup>b,c</sup>, Andre M.N. Renzaho <sup>d,e,f</sup>

<sup>a</sup> Melbourne School of Population and Global Health, University of Melbourne, Parkville, Victoria, Australia.

<sup>b</sup> Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive, Melbourne, Australia.

<sup>c</sup> Copenhagen Centre for Regulatory Science (CORS), University of Copenhagen, Copenhagen, Denmark.

School of Social Sciences, Western Sydney University, Penrith 2751, Australia

Translational Health Research Institute, Western Sydney University, Penrith 2751, Australia

Burnet Institute, Maternal, Child and Adolescent Health Program, Melbourne 3004, Australia

**Correspondence to:** Berhe W. Sahle, Centre for Mental Health Melbourne School of Population and Global Health, The University of Melbourne, 207 Bouverie Street, Carlton, VIC, 3010, Australia, [berhe.sahle@unimelb.edu.au](mailto:berhe.sahle@unimelb.edu.au), +61 3 90357799

**This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/jpc.15132](https://doi.org/10.1111/jpc.15132)**

**Conflict of interest:** The authors declare that they have no competing interests

**Funding:** Berhe W. Sahle acknowledges the support of the NHMRC and Beyond Blue co-funded Centre of Research Excellence in Childhood Adversity and Mental Health (#1153419)

**Abbreviations:** ACEs- Adverse Childhood Experiences

#### **Contributors' Statement Page**

Dr. Berhe W. Sahle conceptualized the study, drafted the initial manuscript, and reviewed and revised the manuscript. Dr Richard and Prof Renzaho have provided critical revision of the manuscript for important intellectual content.

## Adverse childhood experiences should be priority in global response to COVID-19

Berhe W. Sahle <sup>a</sup>, Richard Ofori-Asenso <sup>b,c</sup>, Andre M.N. Renzaho <sup>d,e,f</sup>

<sup>a</sup> Melbourne School of Population and Global Health, University of Melbourne, Parkville, Victoria, Australia.

<sup>b</sup> Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive, Melbourne, Australia.

<sup>c</sup> Copenhagen Centre for Regulatory Science (CORS), University of Copenhagen, Copenhagen, Denmark.

School of Social Sciences, Western Sydney University, Penrith 2751, Australia

Translational Health Research Institute, Western Sydney University, Penrith 2751, Australia

Burnet Institute, Maternal, Child and Adolescent Health Program, Melbourne 3004, Australia

**Correspondence to:** Berhe W. Sahle, Centre for Mental Health Melbourne School of Population and Global Health, The University of Melbourne, 207 Bouverie Street, Carlton, VIC, 3010, Australia, [berhe.sahle@unimelb.edu.au](mailto:berhe.sahle@unimelb.edu.au), +61 3 90357799

**Conflict of interest:** The authors declare that they have no competing interests

**Funding:** Berhe W. Sahle acknowledges the support of the NHMRC and Beyond Blue co-funded Centre of Research Excellence in Childhood Adversity and Mental Health (#1153419)

**Abbreviations:** ACEs- Adverse Childhood Experiences

#### **Contributors' Statement Page**

Dr. Berhe W. Sahle conceptualized the study, drafted the initial manuscript, and reviewed and revised the manuscript. Dr Richard and Prof Renzaho have provided critical revision of the manuscript for important intellectual content.

#### **Adverse childhood experiences should be priority in global response to COVID-19**

We read your editorial<sup>1</sup> and accompanying letter<sup>2</sup> with interest. Children are at low risk of novel coronavirus disease 2019 (COVID-19) infection and tend to experience a milder clinical course and a very low risk of death.<sup>3</sup> Despite the low risk of COVID-19 to children, their risk of exposure to adverse childhood experiences (ACEs), such as childhood maltreatment, household dysfunction, maladaptive parenting, violence and economic adversity, has substantially increased due to the social, financial and psychological consequences of the ongoing COVID-19 pandemic.

Mass unemployment has an adverse impact on the health, financial and social circumstances of workers, families, and communities. Globally, the number of people

reporting depression and anxiety symptoms spiked after the lockdown. Financial hardships and stress from confinement increase the risk of suicide, alcohol abuse, divorce and conflict, domestic violence, mental illness, negative parenting practices and loss of social support networks. The rates of domestic violence and alcohol misuse have increased substantially during the lockdown.<sup>4</sup> ACEs are linked with an increased risk of unhealthy lifestyle behaviours, obesity, suicidality, and chronic diseases, including mental disorders and cardiometabolic outcomes across the life course.<sup>5</sup> About 30% of all mental disorders are attributable to ACEs, including 30% of cases of anxiety, 40% of cases of depression, and 67% of lifetime suicide attempts.<sup>6</sup>

The COVID-19 pandemic has led to a further increase in the burden of ACEs. Even after the global pandemic wears on, it is likely that the health impact of ACEs will continue across the life-course. Unfortunately, at present, there is no systematic approach to the prevention of exposure to ACEs or identification of those exposed to ACEs who are at increased future risk of poor health outcomes. ACEs are diverse and their impact on health and well-being are multifaceted, involving a complex interplay of many biological and behavioural pathways. As a result, preventing and responding to ACEs requires integrated, system-wide, multisectorial approaches to effectively prevent and mitigate the negative impacts of ACEs. Addressing the underlying social determinants of health such as socioeconomic adversity, unemployment, and/or poor and unsafe housing and living conditions could have the largest population level impact on ACEs. Evidence-based parenting and home visiting programs reduce ACEs and global psychopathology. Comprehensive community-wide programs have been shown to reduce child maltreatment and build community-wide resilience. Vulnerability to the impacts of COVID-19 and availability of resources and readiness to respond to the COVID-19 pandemic varies across countries. Addressing access barriers to health and social services, especially among disadvantaged families will be essential to the early detection and responding to ACEs. As such, our response to COVID-19 should consider the socioeconomic, cultural and political context of the affected communities.

## References

1. Isaacs D. Apocalypse Perhaps. *J. Paediatr. Child Health* 2020; **56**:

2. Kilham HA. COVID-19 Apocalypse for children: predictable, preventable? *J. Paediatr. Child Health* 2020; **56**:
3. Ibrahim LF, Tosif S, McNab S, et al. SARS-CoV-2 Testing and Outcomes in the First 30 Days after the First Case of COVID-19 at an Australian Children's Hospital. *Emerg. Med. Austral.* First published: 10 May 2020. <https://doi.org/10.1111/1742-6723.13550>.
4. Bradley NL, DiPasquale AM, Dillabough K, Schneider PS. Health care practitioners' responsibility to address intimate partner violence related to the COVID-19 pandemic. *Can. Med. Assoc. J.* 2020;**192**(22):E609-E610.
5. Deschênes SS, Graham E, Kivimäki M, Schmitz N. Adverse Childhood Experiences and the Risk of Diabetes: Examining the Roles of Depressive Symptoms and Cardiometabolic Dysregulations in the Whitehall II Cohort Study. *Diabetes Care.* 2018;**41**(10):2120-6.
6. Kessler RC, McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky AM, et al. Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. *Brit. J. Psychiatry.* 2010;**197**(5):378-85.