

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Sim, WL;Mutha, V;Ul-Haq, MA;Sasongko, V;Van-Gaal, W

Title:

Clinical characteristics and outcomes of octogenarians presenting with ST elevation myocardial infarction in the Australian population

Date:

2017-05-26

Citation:

Sim, W. L., Mutha, V., Ul-Haq, M. A., Sasongko, V. & Van-Gaal, W. (2017). Clinical characteristics and outcomes of octogenarians presenting with ST elevation myocardial infarction in the Australian population. WORLD JOURNAL OF CARDIOLOGY, 9 (5), pp.437-441. <https://doi.org/10.4330/wjc.v9.i5.437>.

Persistent Link:

<https://hdl.handle.net/11343/256900>

License:

CC BY-NC

World Journal of *Cardiology*

World J Cardiol 2017 May 26; 9(5): 396-469





REVIEW

- 396 Sleep, health behaviors, and behavioral interventions: Reducing the risk of cardiovascular disease in adults
Kaar JL, Luberto CM, Campbell KA, Huffman JC

MINIREVIEWS

- 407 Heart failure after myocardial infarction in the era of primary percutaneous coronary intervention: Mechanisms, incidence and identification of patients at risk
Cahill TJ, Kharbanda RK
- 416 Transcervical access, reversal of flow and mesh-covered stents: New options in the armamentarium of carotid artery stenting
Paraskevas KI, Veith FJ
- 422 Empirical anticoagulation for patients in sinus rhythm at high risk of ischaemic stroke: A review of current literature
Battipaglia I, O'Neill J, Hogarth AJ, Tayebjee MH
- 429 Antitachycardia pacing programming in implantable cardioverter defibrillator: A systematic review
De Maria E, Giacomelli D, Borghi A, Modonesi L, Cappelli S

ORIGINAL ARTICLE

Retrospective Cohort Study

- 437 Clinical characteristics and outcomes of octogenarians presenting with ST elevation myocardial infarction in the Australian population
Sim WL, Mutha V, Ul-Haq MA, Sasongko V, Van-Gaal W

Retrospective Study

- 442 Jailing polymer jacketed guide-wires during bifurcation coronary interventions is associated with procedural myocardial infarction
Chatterjee A, White JS, Hashim T, Leeser MA

Observational Study

- 448 Markers of inflammation and cardiovascular disease in recently diagnosed celiac disease patients
Tetzlaff WF, Meroño T, Menafra M, Martin M, Botta E, Matoso MD, Sorroche P, De Paula JA, Boero LE, Brites F

Prospective Study

- 457** Combined assessment of myocardial damage and electrical disturbance in chronic heart failure
Kadowaki S, Watanabe T, Otaki Y, Narumi T, Honda Y, Takahashi H, Arimoto T, Shishido T, Miyamoto T, Kubota I

CASE REPORT

- 466** Cough induced syncope: A hint to cardiac tamponade diagnosis
Ramirez R, Lasam G

Contents

World Journal of Cardiology
Volume 9 Number 5 May 26, 2017

ABOUT COVER

Editorial Board Member of *World Journal of Cardiology*, Gergely Feher, MD, PhD, Assistant Professor, Department of Neurology, Medical School, University of Pecs, Pecs, Baranya 7623, Hungary

AIM AND SCOPE

World Journal of Cardiology (*World J Cardiol*, *WJC*, online ISSN 1949-8462, DOI: 10.4330) is a peer-reviewed open access journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJC covers topics concerning arrhythmia, heart failure, vascular disease, stroke, hypertension, prevention and epidemiology, dyslipidemia and metabolic disorders, cardiac imaging, pediatrics, nursing, and health promotion. Priority publication will be given to articles concerning diagnosis and treatment of cardiology diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, differential diagnosis, imaging tests, pathological diagnosis, molecular biological diagnosis, immunological diagnosis, genetic diagnosis, functional diagnostics, and physical diagnosis; and comprehensive therapy, drug therapy, surgical therapy, interventional treatment, minimally invasive therapy, and robot-assisted therapy.

We encourage authors to submit their manuscripts to *WJC*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING

World Journal of Cardiology is now indexed in Emerging Sources Citation Index (Web of Science), PubMed, and PubMed Central.

FLYLEAF

I-IV Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Huan-Liang Wu*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Fang-Fang Ji*
Proofing Editorial Office Director: *Xiu-Xia Song*

NAME OF JOURNAL
World Journal of Cardiology

ISSN
ISSN 1949-8462 (online)

LAUNCH DATE
December 31, 2009

FREQUENCY
Monthly

EDITORS-IN-CHIEF
Jian-Jun Li, MD, PhD, Professor, Center for Coronary Artery Disease, Fu Wai Cardiovascular Hospital, Chinese Academy of Medical Science, Beijing 100037, China

Giuseppe De Luca, PhD, Assistant Professor, Department of Cardiology, Piedmont University, Novara 28100, Italy

Nathan D Wong, FACC, FAHA, PhD, Director, Professor, Heart Disease Prevention Program, Division of Cardiology, Department of Medicine, University of California, Irvine, CA 92629, United States

sity of California, Irvine, CA 92629, United States

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/1949-8462/editorialboard.htm>

EDITORIAL OFFICE
Xiu-Xia Song, Director
World Journal of Cardiology
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
May 26, 2017

COPYRIGHT
© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.f6publishing.com>

Retrospective Cohort Study

Clinical characteristics and outcomes of octogenarians presenting with ST elevation myocardial infarction in the Australian population

Wei Liang Sim, Vivek Mutha, Muhammad Asrar Ul-Haq, Victoria Sasongko, William Van-Gaal

Wei Liang Sim, Victoria Sasongko, William Van-Gaal, Department of Cardiology, Northern Hospital, Epping, Victoria 3076, Australia

Vivek Mutha, Department of Cardiology, Monash Health, Melbourne, Victoria 3168, Australia

Muhammad Asrar Ul-Haq, Department of Cardiology, the University of Melbourne, Melbourne, Victoria 3010, Australia

Author contributions: All the authors solely contributed to this paper.

Institutional review board statement: The study was reviewed and approved for publication by our Institutional Reviewer.

Informed consent statement: All study participants or their legal guardian provided informed written consent about personal and medical data collection prior to study enrolment.

Conflict-of-interest statement: All authors have no conflict of interest related to the manuscript.

Data sharing statement: The original anonymous dataset is available on request from the corresponding author at wei_liangs@hotmail.com.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Correspondence to: Dr. Wei Liang Sim, Department of Cardiology, Northern Hospital, 185 Cooper Street, Epping, Victoria 3076, Australia. wei_liangs@hotmail.com
Telephone: +61-04-28909100

Received: August 26, 2016

Peer-review started: August 29, 2016

First decision: October 28, 2016

Revised: February 20, 2017

Accepted: March 21, 2017

Article in press: March 22, 2017

Published online: May 26, 2017

Abstract

AIM

To investigate the characteristics and outcomes of octogenarians who presented with ST-elevation myocardial infarction (STEMI) compared to non-octogenarians and to investigate the outcomes of octogenarians that received primary percutaneous coronary intervention (PCI) compared to those managed conservatively.

METHODS

We performed a single center retrospective case controlled study. All octogenarians who presented with STEMI to a tertiary referring hospital between 2007 and 2012 were included. The subsequent non-octogenarian patient who presented with a STEMI following the octogenarian patient was assigned to the control group in a 1:1 manner. The outcomes measured were peri-procedural cardiac arrest, death on table, cerebrovascular accidents (CVA), in-hospital and 30-d mortality.

RESULTS

A total of 146 patients were analyzed. The octogenarian group had a higher percentage of females, higher overall comorbidities, higher Charlson Comorbidity Index score, worse renal function and were more likely to require residential care and home help. The octogenarian group were also less likely to have PCI attempted and had a longer symptom onset to PCI

time. Mortality rate was high amongst octogenarians who presented with STEMI. However, those managed conservatively had a higher in-hospital and 30-d mortality rate

CONCLUSION

Octogenarians who presented with STEMI that were managed conservatively had a higher mortality rate compared to those who had primary PCI. Therefore, we propose that revascularization may be beneficial to patients in this age group.

Key words: Coronary disease; Acute coronary syndrome; Myocardial infarction; Percutaneous coronary intervention; Aged 80 and over

© The Author(s) 2017. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The octogenarian group represents a complex population with multiple comorbidities. Percutaneous coronary intervention in this group is challenging and is associated with a high rate of failure and complications. This study shows that the mortality rate amongst octogenarians presenting with ST elevation myocardial infarction is high. However, there may be a mortality benefit in those treated with percutaneous coronary intervention, compared to those managed conservatively.

Sim WL, Mutha V, Ul-Haq MA, Sasongko V, Van-Gaal W. Clinical characteristics and outcomes of octogenarians presenting with ST elevation myocardial infarction in the Australian population. *World J Cardiol* 2017; 9(5): 437-441 Available from: URL: <http://www.wjgnet.com/1949-8462/full/v9/i5/437.htm> DOI: <http://dx.doi.org/10.4330/wjc.v9.i5.437>

INTRODUCTION

Advanced age is associated with increased risk of acute coronary syndrome (ACS) and cardiovascular comorbidities^[1]. The octogenarian population (age $\geq$ 80 years) is a fast growing segment of the population worldwide, and represent a high risk group for procedural complications during percutaneous coronary intervention (PCI) particularly in the settings of ST-segment elevation myocardial infarction (STEMI)^[2]. These patients are underrepresented in randomized clinical trials evaluating primary PCI for STEMI and a high mortality has been reported^[3,4]. These patients are typically treated less aggressively than are younger patients, due partly to the increased risk of adverse events and PCI related complications, and partly to a lack of standard management guidelines. Evidence based management of octogenarian patients with STEMI thus remains suboptimal despite the high mortality^[2]. The overseas observational trials have suggested that despite the recommendations being

that age should not influence the decision of reperfusion strategy in STEMI patients, older age remains one of the strong predictors of not receiving it. There is paucity of Australia data on outcomes of octogenarian who present with STEMI. In this context, the aim of our study was to assess the clinical characteristics and outcomes of octogenarians presenting with STEMI, as compared with non-octogenarian patients (age < 80 years), as well as the outcomes of octogenarians who received primary PCI, compared to those that were managed conservatively.

MATERIALS AND METHODS

This study is a single center retrospective case controlled study including all octogenarians who presented with STEMI between 2007 and 2012 in a tertiary Australian hospital. The subsequent non-octogenarian patient who presented with STEMI following an octogenarian STEMI was assigned to the control group in 1:1 manner. Detailed data on baseline and procedural characteristics and patient comorbidities were obtained through electronic medical records, and compared between octogenarian and non-octogenarian STEMI patients. The charlson comorbidity index (CCI) was calculated based on the patient's comorbidities. CCI predicts long-term survival according to a patient's medical condition^[5]. STEMI was defined as persistent angina for 20 min in conjunction with either: (1) an ST-segment elevation at the J point of 0.25 mV in men aged < 40 years or 0.2 mV in men aged > 40 years or 0.15 mV in women in the precordial leads V2 to V3, and 0.1 mV in all other leads; or (2) the presence of a new left bundle branch block^[6]. PCI success was defined as TIMI 2 or 3 flow post intervention. Left ventricular ejection fraction (EF) was derived either from the echocardiogram performed following the presentation or coronary angiogram during the index admission. Outcomes compared between the two groups included peri-procedural cardiac arrest, death on table, cerebrovascular accident (CVA), in-hospital and 30-d mortality. CVA was defined a clinical evidence of neurological deficit leading to a documented diagnosis of transient ischaemic attack or stroke. Subgroups of octogenarian STEMI patients who received PCI vs who did not (conservatively managed) were also compared for baseline and clinical characteristics. In-hospital and 30-d mortality was compared between all subgroups and independent predictors calculated.

All data were analyzed using IBM SPSS v22 and presented as percentages or mean value $\pm$ standard deviation (SD). Independent *t* test was used to compare continuous while χ^2 and Fisher's exact tests were performed for categorical data. Logistic regression and multivariate analysis were performed to identify independent predictors. A two tailed *P* value of < 0.05 was considered statistically significant. The statistical methods of this study were reviewed by our biostatistics expert Dr. Asrar Ul-Haq, MBBS.

Table 1 Baseline characteristics, procedural data, and outcomes of octogenarians as compared to non-octogenarians (controls)

	Octogenarians (<i>n</i> = 73)	Controls (<i>n</i> = 73)	<i>P</i>
Age	85.2 ± 4.1	67.1 ± 5.3	< 0.005
Females	56	29	< 0.005
Residential care	23	2.7	< 0.005
LLC	11	0	< 0.005
HLC	12	2.7	< 0.005
Home help	25	0	< 0.005
Medical comorbidities			
Diabetes	38	19	< 0.05
eGFR	48.7 ± 19.9	68.1 ± 20.3	< 0.005
PVD	30	12	< 0.05
Prior IHD	36	20	0.06
EF	53.6 ± 14.1	50.8 ± 13.1	0.4
Charlsons	3.2 ± 2.3	1.7 ± 2.2	< 0.005
Presentation and procedural characteristics			
Location of MI			
Anterior	51	49	0.8
Inferior	42	47	0.5
Lateral	4.1	4.1	1.0
PCI attempt	47	84	< 0.005
PCI success (TIMI 2-3)	91	99	0.1
Symptoms onset to PCI < 6 h	16	45	< 0.005
Outcomes			
Peri-procedural cardiac arrest	5	3	0.1
Death on table	1.8	0.9	0.2
Stroke	1.4	0	0.3
In-hospital mortality	28	7	< 0.005
30-d mortality	45	12	< 0.005

Data are means ± SD or *n* (%). LLC: Low level care; HLC: High level care; PVD: Peripheral vascular disease; EF: Ejection fraction; PCI: Percutaneous coronary intervention; MI: Myocardial infarction.

RESULTS

Octogenarians vs non-octogenarians

A total of 146 patients were analysed (octogenarians = 73; non-octogenarians = 73). The mean age was 85.2 ± 4.1 years in the octogenarian group and 67.1 ± 5.3 years in the control group (Table 1). The octogenarian group had a higher percentage of females (56% vs 29%, *P* < 0.005), higher overall comorbidities, a higher CCI score (3.2 ± 2.3 vs 1.7 ± 2.2, *P* < 0.001), were more likely to require residential care (23% vs 2.7%, *P* < 0.001) as well as home help (25% vs 0%, *P* < 0.001), and had worse renal function (eGFR 48.7 ± 19.9 vs 68.1 ± 20.3 mL/min per 1.73 m², *P* < 0.001).

Octogenarians were less likely to have PCI attempted compared to the non-octogenarians (47% vs 84%, *P* < 0.001). The rate of symptom onset-to-PCI of < 6 h was significantly lower in octogenarians (16% vs 45%, *P* < 0.001). The rate of PCI success was high in both groups (91% vs 99%, *P* = 0.1). Reasons PCI was not attempted in non-octogenarians include: No culprit found (3), embolic event (1), recent CVA (1), known or new triple vessel disease/complex anatomy (2), other comorbidities (5); and in octogenarians: No culprit found (9), embolic event (2), recent CVA (4), known or

Table 2 Baseline characteristics and outcomes of octogenarians who received percutaneous coronary intervention compared to conservatively managed octogenarians (no-percutaneous coronary intervention)

	PCI (<i>n</i> = 34)	No-PCI (<i>n</i> = 39)	<i>P</i>
Age	84 ± 3.4	86 ± 4.3	< 0.05
Females	44	67	0.06
Residential care	3	41	< 0.005
LLC	3	18	< 0.005
HLC	0	23	< 0.005
Home help	18	35	0.2
Medical comorbidities			
Diabetes	32	44	0.3
eGFR	54 ± 23	44 ± 16	< 0.05
PVD	21	38	0.1
Prior IHD	35	36	1
EF	54 ± 13	53 ± 16	0.2
Charlsons	2.52 ± 2	3.8 ± 2	< 0.05
Presentation			
Location of MI			
Anterior	41	59	0.2
Inferior	50	36	0.2
Lateral	6	3	0.6
Outcomes			
In-hospital mortality	18	37	0.1
30-d mortality	29	59	< 0.05

Data are means ± SD or *n* (%). LLC: Low level care; HLC: High level care; PVD: Peripheral vascular disease; EF: Ejection fraction; PCI: Percutaneous coronary intervention; MI: Myocardial infarction.

new triple vessel disease/complex anatomy (13), other comorbidities (11). Octogenarians had a significantly higher overall in-hospital mortality (28% vs 7%, *P* < 0.005) and 30-d mortality (45% vs 12%, *P* < 0.001).

The independent predictors of 30-d mortality in octogenarians included age (OR 1.20/year of advancing age, *P* < 0.01), place of residence (OR = 4.4, *P* < 0.01 for nursing home), conservative management (No intervention - OR 2.77, *P* < 0.05), and declining renal function (OR = 0.9, *P* < 0.05).

PCI vs conservatively managed octogenarians

The 39 (53%) octogenarians who did not receive PCI were older (86 ± 4.3 years vs 84 ± 3.4 years, *P* < 0.05) and were more likely to be in residential care (41% vs 3%, *P* < 0.001), had higher CCI score (3.8 ± 2 vs 2.52 ± 2, *P* < 0.05) and worse renal function (eGFR 44 ± 16 mL/min vs 54 ± 23 mL/min per 1.73 m²). Type of myocardial infarction was not different as compared to octogenarians who received PCI (Table 2).

Mortality rate was high among octogenarians who presented with STEMI. However, those who were managed conservatively had a higher in-hospital and 30-d mortality (37% vs 18%, *P* = 0.1; and 59% vs 29%, *P* < 0.05 respectively).

Independent predictors of intervention in octogenarians included younger age (OR = 0.86, *P* < 0.05), place of residence (OR = 0.1, *P* < 0.05 for nursing home), lower CCI (OR = 0.7, *P* < 0.05), and renal

function (OR = 1.03, $P < 0.05$).

DISCUSSION

High mortality

Our study demonstrated that mortality rate amongst octogenarians presenting with STEMI is high in Australian population despite the offered treatment, although much worse when treated conservatively. This appears to be associated with higher overall comorbidities, higher CCI score, worse renal function, and need for residential care or home help (which maybe the indirect measure of overall comorbidities and physical state). These findings are consistent with overseas studies looking at similar age groups^[7-9]. Some factors reported to affect the mortality in these studies include heart failure, multiple co-morbidities, cachexia, cognitive state, history of intra-cranial bleeding and pre-hospital physical activity status^[10,11]. Furthermore, it has been shown that the elderly are less likely to receive evidence based medical treatment such as aspirin, clopidogrel, beta-blockers, statins or glycoprotein IIb/IIIa inhibitors^[7,8]. This may be due to concerns with regards to potential side effects in this age group. Previous studies have also shown that the elderly is associated with a higher rate of PCI failure. However, this is not reflected in our study, likely because the candidates for treatment were carefully selected.

Underuse of invasive treatment

Our study suggested that despite having a higher mortality rate, octogenarians are less likely to have PCI attempted as compared to non-octogenarians. Frailty, co-morbidities and time delays have been shown to contribute to the underuse of invasive therapies. In our study, the proportion of those who received PCI in less than 6 h was significantly lower in the octogenarian group. This might reflect difficulty in decision-making with regards to reperfusion strategy. Atypical clinical presentation is also more common in the elderly and could contribute to time delay as well as the higher prevalence of cardiac failure. In addition, female gender was more prevalent in the octogenarian group. Previous studies have shown that female gender is associated with lower use of invasive therapies, especially in the elderly^[12].

Invasive treatment appears beneficial

Another major finding of our study is that despite a relatively poor prognosis, octogenarians who received primary PCI had a significantly lower 30-d mortality. This however maybe related to the selection bias, a limitation of observational study design. Patients managed conservatively in our study were older, more likely to be in residential care, had a higher CCI score and worse renal function, and this represents a group with higher risk profile. The most common reasons PCI was not performed in the elderly were triple vessel disease/complex anatomy, other co-

existing comorbidities and the absence of a clear culprit lesion. There was no significant difference in in-hospital mortality between those managed who received intervention and those managed conservatively. Other factors affecting 30-d mortality included age, place of residence and declining renal function, highlighting the complexity of this patient population.

Limitations

Our study is not randomized, and therefore limited by selection bias. However, it is an "all comers" registry which reflects "real world" data on management and outcomes of elderly patients who presents with STEMI. A randomised trial in this particular group is not viable. Propensity score matching would be the next best option and requires larger studies. Furthermore, we did not evaluate long term mortality and re-infarction rates, which may provide incremental information and a better picture of the utility of invasive management in this group.

In conclusion, this study is the first Australian report on the outcomes of octogenarians who present with ST-elevation myocardial infarction. This group represents a complex population with multiple medical comorbidities and PCI is challenging, associated with high failure and complication rates. Consequently, mortality is high in this group. However, the detrimental prognosis of conservatively managed octogenarians and relative mortality benefit associated with PCI suggests that revascularization therapy may benefit this age group.

COMMENTS

Background

The elderly have an increased risk of acute coronary syndrome. However, there are more likely to be managed conservatively compared to the younger cohort. This is due to concerns regarding procedural complications and success. The elderly is also under-represented in major clinical trials evaluating primary percutaneous coronary intervention (PCI). Recognizing this there have been recent overseas studies evaluating the elderly and primary PCI.

Research frontiers

This study is the first to evaluate the characteristics and outcome of octogenarians who presents with ST-elevation myocardial infarction (STEMI) in an Australian setting. The authors also evaluated the outcomes of octogenarians who received primary PCI compared to those that were managed conservatively.

Innovations and breakthroughs

This paper showed that despite poor prognosis among octogenarians who presents with STEMI, primary PCI may offer some benefit, with significantly lower 30 d mortality in the group that received it.

Applications

Primary PCI should be considered in octogenarians who presents with STEMI. The patient's co-morbidities, quality of life and life expectancy should be taken into account when making this decision.

Terminology

Primary PCI consists of urgent balloon angioplasty (with or without stenting), without the previous administration of fibrinolytic therapy or platelet glycoprotein IIb/IIIa inhibitors, to open the infarct-related artery during an acute myocardial infarction with ST-segment elevation.

Peer-review

This is an interesting and well-written article dealing with care on octogenarians after acute myocardial infarction. The authors have found worse clinical characteristics in octogenarians in comparison with non octogenarians, as well as lower referral for interventional procedures.

REFERENCES

- 1 **Wei JY**, Gersh BJ. Heart disease in the elderly. *Curr Probl Cardiol* 1987; **12**: 1-65 [PMID: 3549164]
- 2 **Levi A**, Kornowski R, Vaduganathan M, Eisen A, Vaknin-Assa H, Abu-Foul S, Lev EI, Brosh D, Bental T, Assali AR. Incidence, predictors, and outcomes of failed primary percutaneous coronary intervention: a 10-year contemporary experience. *Coron Artery Dis* 2014; **25**: 145-151 [PMID: 24281252 DOI: 10.1097/MCA.000000000000065]
- 3 **Lee PY**, Alexander KP, Hammill BG, Pasquali SK, Peterson ED. Representation of elderly persons and women in published randomized trials of acute coronary syndromes. *JAMA* 2001; **286**: 708-713 [PMID: 11495621 DOI: 10.1001/jama.286.6.708]
- 4 **Dodd KS**, Saczynski JS, Zhao Y, Goldberg RJ, Gurwitz JH. Exclusion of older adults and women from recent trials of acute coronary syndromes. *J Am Geriatr Soc* 2011; **59**: 506-511 [PMID: 21361882 DOI: 10.1111/j.1532-5415.2010.03305.x]
- 5 **Charlson ME**, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987; **40**: 373-383 [PMID: 3558716]
- 6 **Thygesen K**, Alpert JS, Jaffe AS, Simoons ML, Chaitman BR, White HD, Thygesen K, Alpert JS, White HD, Jaffe AS, Katus HA, Apple FS, Lindahl B, Morrow DA, Chaitman BA, Clemmensen PM, Johanson P, Hod H, Underwood R, Bax JJ, Bonow RO, Pinto F, Gibbons RJ, Fox KA, Atar D, Newby LK, Galvani M, Hamm CW, Uretsky BF, Steg PG, Wijns W, Bassand JP, Menasché P, Ravkilde J, Ohman EM, Antman EM, Wallentin LC, Armstrong PW, Simoons ML, Januzzi JL, Nieminen MS, Gheorghiade M, Filippatos G, Luepker RV, Fortmann SP, Rosamond WD, Levy D, Wood D, Smith SC, Hu D, Lopez-Sendon JL, Robertson RM, Weaver D, Tendera M, Bove AA, Parkhomenko AN, Vasilieva EJ, Mendis S. Third universal definition of myocardial infarction. *Eur Heart J* 2012; **33**: 2551-2567 [PMID: 22922414 DOI: 10.1093/eurheartj/ehs184]
- 7 **Fach A**, Bünger S, Zabrocki R, Schmucker J, Conradi P, Garstka D, Fiehn E, Hambrecht R, Wienbergen H. Comparison of Outcomes of Patients With ST-Segment Elevation Myocardial Infarction Treated by Primary Percutaneous Coronary Intervention Analyzed by Age Groups (< 75, 75 to 85, and > 85 Years); (Results from the Bremen STEMI Registry). *Am J Cardiol* 2015; **116**: 1802-1809 [PMID: 26602071 DOI: 10.1016/j.amjcard.2015.09.022]
- 8 **Yamanaka F**, Jeong MH, Saito S, Ahn Y, Chae SC, Hur SH, Hong TJ, Kim YJ, Seong IW, Chae JK, Rhew JY, Chae IH, Cho MC, Bae JH, Rha SW, Kim CJ, Choi D, Jang YS, Yoon J, Chung WS, Cho JG, Seung KB, Park SJ. Comparison of clinical outcomes between octogenarians and non-octogenarians with acute myocardial infarction in the drug-eluting stent era: analysis of the Korean Acute Myocardial Infarction Registry. *J Cardiol* 2013; **62**: 210-216 [PMID: 23731919 DOI: 10.1016/j.jjcc.2013.04.003]
- 9 **Vandecasteele EH**, De Buyzere M, Gevaert S, de Meester A, Convens C, Dubois P, Boland J, Sinnave P, De Raedt H, Vranckx P, Coussement P, Evrard P, Beauloye C, Renard M, Claeys MJ. Reperfusion therapy and mortality in octogenarian STEMI patients: results from the Belgian STEMI registry. *Clin Res Cardiol* 2013; **102**: 837-845 [PMID: 23887831 DOI: 10.1007/s00392-013-0600-3]
- 10 **Schwarz AK**, Zahn R, Hochadel M, Kerber S, Hauptmann KE, Glunz HG, Mudra H, Darius H, Zeymer U. Age-related differences in antithrombotic therapy, success rate and in-hospital mortality in patients undergoing percutaneous coronary intervention: results of the quality control registry of the Arbeitsgemeinschaft Leitende Kardiologische Krankenhausärzte (ALKK). *Clin Res Cardiol* 2011; **100**: 773-780 [PMID: 21509588 DOI: 10.1007/s00392-011-0311-6]
- 11 **Miyamoto T**, Obayashi T, Hattori E, Yamauchi Y, Niwa A, Isobe M. Pre-hospital physical activity status affects in-hospital course of elderly patients with acute myocardial infarction. *J Med Dent Sci* 2010; **57**: 119-126 [PMID: 20437772 DOI: 10.11480/jmds.570113]
- 12 **Matsui K**, Fukui T, Hira K, Sobashima A, Okamatsu S, Hayashida N, Tanaka S, Nobuyoshi M. Impact of sex and its interaction with age on the management of and outcome for patients with acute myocardial infarction in 4 Japanese hospitals. *Am Heart J* 2002; **144**: 101-107 [PMID: 12094195 DOI: 10.1067/mhj.2002.123114]

P- Reviewer: Peteiro J, Petretta M, Kettering K, Chang ST

S- Editor: Kong JX **L- Editor:** A **E- Editor:** Wu HL





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

