



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Kim, J;Olaiya, MT;De Silva, DA;Norrving, B;Bosch, J;De Sousa, DA;Christensen, HK;Ranta, A;Donnan, GA;Feigin, V;Martins, S;Schwamm, LH;Werring, DJ;Howard, G;Owolabi, M;Pandian, J;Mikulik, R;Thayabaranathan, T;Cadilhac, DA

Title:

Global stroke statistics 2023: Availability of reperfusion services around the world

Date:

2024-03-01

Citation:

Kim, J., Olaiya, M. T., De Silva, D. A., Norrving, B., Bosch, J., De Sousa, D. A., Christensen, H. K., Ranta, A., Donnan, G. A., Feigin, V., Martins, S., Schwamm, L. H., Werring, D. J., Howard, G., Owolabi, M., Pandian, J., Mikulik, R., Thayabaranathan, T. & Cadilhac, D. A. (2024). Global stroke statistics 2023: Availability of reperfusion services around the world. *International Journal of Stroke*, 19 (3), pp.253-270. <https://doi.org/10.1177/17474930231210448>.

Persistent Link:

<https://hdl.handle.net/11343/350953>

License:

[CC BY-NC](#)

# Global stroke statistics 2023: Availability of reperfusion services around the world

International Journal of Stroke  
2024, Vol. 19(3) 253–270  
© 2023 World Stroke Organization



Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/17474930231210448  
journals.sagepub.com/home/wso



Joosup Kim<sup>1,2,\$</sup> , Muideen T Olaiya<sup>1,\$</sup> , Deidre A De Silva<sup>3</sup> ,  
Bo Norrving<sup>4</sup>, Jackie Bosch<sup>5</sup>, Diana A De Sousa<sup>6</sup> ,  
Hanne K Christensen<sup>7</sup>, Anna Ranta<sup>8</sup> , Geoffrey A Donnan<sup>9</sup> ,  
Valery Feigin<sup>10</sup> , Sheila Martins<sup>11</sup>, Lee H Schwamm<sup>12</sup>,  
David J Werring<sup>13</sup> , George Howard<sup>14</sup>, Mayowa Owolabi<sup>15</sup> ,  
Jeyaraj Pandian<sup>16</sup> , Robert Mikulik<sup>17,18</sup>,  
Tharshanah Thayabaranathan<sup>1,\*</sup> , and  
Dominique A Cadilhac<sup>1,2,\*</sup> 

## Abstract

**Background:** Disparities in the availability of reperfusion services for acute ischemic stroke are considerable globally and require urgent attention. Contemporary data on the availability of reperfusion services in different countries are used to provide the necessary evidence to prioritize where access to acute stroke treatment is needed.

**Aims:** To provide a snapshot of published literature on the provision of reperfusion services globally, including when facilitated by telemedicine or mobile stroke unit services.

**Methods:** We searched PubMed to identify original articles, published up to January 2023 for the most recent, representative, and relevant patient-level data for each country. Keywords included thrombolysis, endovascular thrombectomy and telemedicine. We also screened reference lists of review articles, citation history of articles, and the gray literature. The information is provided as a narrative summary.

**Results:** Of 11,222 potentially eligible articles retrieved, 148 were included for review following de-duplications and full-text review. Data were also obtained from national stroke clinical registry reports, Registry of Stroke Care Quality (RES-Q) and PRE-hospital Stroke Treatment Organization (PRESTO) repositories, and other national sources. Overall, we found evidence of the provision of intravenous thrombolysis services in 70 countries (63% high-income countries (HICs)) and endovascular thrombectomy services in 33 countries (68% HICs), corresponding to far less than half of the countries in the world. Recent data (from 2019 or later) were lacking for 35 of 67 countries with known year of data (52%). We found published data on 74 different stroke telemedicine programs (93% in HICs) and 14 active mobile stroke unit pre-hospital ambulance services (80% in HICs) around the world.

**Conclusion:** Despite remarkable advancements in reperfusion therapies for stroke, it is evident from available patient-level data that their availability remains unevenly distributed globally. Contemporary published data on availability of reperfusion services remain scarce, even in HICs, thereby making it difficult to reliably ascertain current gaps in the provision of this vital acute stroke treatment around the world.

## Keywords

Stroke, worldwide, reperfusion, telemedicine, mobile stroke unit

Received: 10 September 2023; accepted: 9 October 2023

## Introduction

Stroke is a complex and time-sensitive medical emergency. To improve outcomes following stroke, there is a need for

systems of acute care that will optimize timely access to evidence-based therapies,<sup>1</sup> particularly the provision of reperfusion treatments (intravenous thrombolysis or endovascular thrombectomy) for patients with ischemic stroke.

Reperfusion services are provided by interdisciplinary stroke units in hospitals often designated as a comprehensive or primary stroke center.<sup>1</sup> Where stroke centers are unavailable, telemedicine networks can be used to coordinate the provision of reperfusion therapies with specialist investigations and interventions in stroke-capable hospitals often situated in nonurban locations.<sup>1</sup> In addition, with the advent of mobile stroke unit (MSU) pre-hospital ambulance services, reperfusion therapies are now also provided in the pre-hospital setting.

Due to the increasing burden of stroke globally, the World Stroke Organization (WSO) is leading a global call to action to increase access to reperfusion services for acute stroke.<sup>2</sup> Availability of country-level data on the provision of reperfusion services is important for guiding national health care planning and policy, or funding/resource allocation decisions, to improve outcomes after stroke.

We have previously reported stroke statistics (incidence, case fatality, and mortality) by country,<sup>3–6</sup> and described the global access to stroke units, and the use of national clinical quality registries for routine monitoring of the quality of acute stroke care.<sup>3,4</sup> In this new Global Stroke Statistics article, we present an overview of the provision of reperfusion services (i.e. intravenous thrombolysis and endovascular thrombectomy) for treating acute ischemic stroke. Specifically, we provided a repository of the latest published country-specific data on the provision of reperfusion services, either directly or as facilitated by telemedicine or MSU ambulance services.

## Methods

We undertook a comprehensive search of PubMed for peer-reviewed literature on the provision of thrombolytic therapy (intravenous thrombolysis), endovascular thrombectomy, stroke telemedicine services, or MSU pre-hospital ambulance services, for acute stroke care, using the search terms described previously.<sup>3</sup> Original peer-reviewed articles and review articles published in English, as of January 2023, were identified. The most recent report from national clinical stroke registries was also identified, since these reports comprised a standardized overview of country-specific acute stroke care treatments.<sup>6</sup> We included articles in which nationally representative patient-level data on the provision of reperfusion services were reported. Where multiple studies from one country were identified, data were extracted from the most recent and nationally representative source only. For countries in which nationally representative patient-level data were not available, we included patient-level data from regional or single-center studies.

Three authors (JK, MTO, and TT) undertook the (a) screening of articles by title and abstract; (b) initial review of articles to confirm eligibility; (c) development of templates for data extraction; and (d) extraction of relevant data from included articles or reports. Templates used for data extraction were reviewed by a senior author (DAC) and approved by all authors. Co-authors scrutinized the list of identified articles and reports to assess whether relevant data were missing and contributed any additional relevant original peer-reviewed articles or recent registry reports

<sup>1</sup>Stroke and Ageing Research, Department of Medicine, School of Clinical Sciences at Monash Health, Monash University, Clayton, VIC, Australia

<sup>2</sup>Stroke Theme, The Florey Institute of Neuroscience and Mental Health, Heidelberg, VIC, Australia

<sup>3</sup>Department of Neurology, Singapore General Hospital Campus, National Neuroscience Institute, Singapore

<sup>4</sup>Department of Clinical Sciences, Section of Neurology, Lund University, Skåne University Hospital, Lund, Sweden

<sup>5</sup>School of Rehabilitation Science, Population Health Research Institute, McMaster University, Hamilton, ON, Canada

<sup>6</sup>Department of Neurosciences (Neurology), Hospital de Santa Maria, University of Lisbon, Lisbon, Portugal

<sup>7</sup>Department of Neurology, University of Copenhagen and Bispebjerg Hospital, Copenhagen, Denmark

<sup>8</sup>Department of Medicine, University of Otago, Wellington, Wellington, New Zealand

<sup>9</sup>Melbourne Brain Centre, The University of Melbourne, Royal Melbourne Hospital, Parkville, VIC, Australia

<sup>10</sup>National Institute for Stroke and Applied Neurosciences, Auckland University of Technology, Auckland, New Zealand

<sup>11</sup>Neurology Department, Hospital Moinhos de Vento, Porto Alegre, Brazil

<sup>12</sup>Yale School of Medicine, New Haven, CT, USA

<sup>13</sup>Stroke Research Centre, Department of Brain Repair and Rehabilitation, UCL Queen Square Institute of Neurology, London, UK

<sup>14</sup>Department of Biostatistics, School of Public Health, The University of Alabama at Birmingham, Birmingham, AL, USA

<sup>15</sup>Center for Genomic and Precision Medicine, University of Ibadan, Ibadan, Nigeria

<sup>16</sup>Department of Neurology, Christian Medical College and Hospital, Ludhiana, India

<sup>17</sup>Health Management Institute, Brno, Czech Republic

<sup>18</sup>Neurology Department, Bata Hospital, Zlin, Czech Republic


§These two authors are joint first authors

\*These two authors are joint senior authors

### Corresponding author:

Dominique A Cadilhac, Stroke and Ageing Research Group, Clinical Sciences at Monash Health, Victorian Heart Hospital, Monash University, Level 2, Clayton, VIC 3168, Australia.

Email: dominique.cadilhac@monash.edu

**Figure 1.** Screening and selection of published articles and relevant reports.

that were missed in the initial search. Co-authors from Sweden, Denmark, Canada, and Singapore provided unpublished, but verified, latest data from relevant national stroke clinical registries. For the availability of MSUs, we compared published articles with information available in the PRE-hospital Stroke Treatment Organization (PRESTO) website.<sup>7</sup>

Data extracted included (a) the availability of intravenous thrombolysis services in a country; (b) proportions of patients with ischemic stroke provided with intravenous thrombolysis, and proportions of hospitals providing intravenous thrombolysis, in countries where such services were reported to be available; (c) the availability of endovascular thrombectomy services in a country; (d) proportions of patients with ischemic stroke provided with endovascular thrombectomy, and proportions of hospitals providing endovascular thrombectomy, in countries

where such services were reported to be available; (e) availability of stroke telemedicine programs, including the name, location (city and country), and model/composition (number of hospitals, hubs, and spokes) of such programs; and (f) availability of MSU pre-hospital ambulance services, including the location (city and country).

## Results

Overall, 11,222 publications were identified from electronic databases, and four published articles and one report recommended by a co-authors DADS and JB. Data were also obtained from reports from national stroke clinical quality registries of nine countries, the PRESTO, and the Registry of Stroke Care Quality (RES-Q) repositories. Overall, 148 records with data from 70 countries met our inclusion criteria (Figure 1). Data were mostly from the

regions of Western Europe (25.7%), Central Europe (12.9%), Southeast Asia (10.0%), and North Africa and the Middle East (10.0%; Table 1). Data from 26 countries (37.1%) were nationally representative, including 17 from national registry sources and nine from other sources (Table 1). For 35 of 67 countries in which the year of data was known (52.2%), the latest data reported were from 2019 onwards.

### *Provision of acute stroke reperfusion services*

There was evidence of the availability of intravenous thrombolysis services in all 70 countries (Table 1), including 44 high-income countries (HICs; 62.9%), 14 upper-middle-income countries (20.0%), 10 lower-middle-income countries (14.3%), and 2 low-income countries (i.e. Nepal and Democratic Republic of the Congo; 2.9%). We found evidence of the availability of endovascular thrombectomy services in 33 countries (Table 1), including 22 HICs (67.7%), 8 upper-middle-income countries (24.2%), and 3 from lower-middle-income countries (i.e. Egypt, India, and Vietnam; 9.1%). There was no evidence of the availability of endovascular thrombectomy services in any of the low-income countries with published data on intravenous thrombolysis services. Approximately half of the countries with evidence of endovascular thrombectomy services were in Western Europe (10 countries) or Central Europe (5 countries; Figure 2).

We identified 14 countries with nationally representative and recent data (from 2019 onwards) on the proportion of patients with ischemic stroke provided with reperfusion services. This included 12 countries with information on both intravenous thrombolysis and endovascular thrombectomy, and two countries (Scotland and Barbados) with information on intravenous thrombolysis only. Among countries with nationally representative data, the proportion of patients with ischemic stroke treated with thrombolysis ranged from <10% (Singapore, China, France, Barbados), 10–14% (the United Kingdom, Australia, New Zealand, Sweden), and ≥15% (Czech Republic, Scotland, Denmark, Italy, Slovakia, Switzerland; Table 1). This proportion ranged from <5% (Singapore, the United Kingdom, China), 5–9% (Denmark, Czech Republic, New Zealand, Sweden, France, Italy), and >10% (Australia, Slovakia, Switzerland) for the provision of endovascular thrombectomy.

### *Availability of stroke telemedicine services*

We identified 74 stroke telemedicine programs, with 69 programs located in HICs (Table 2). The majority of the stroke telemedicine programs identified were located in the United States (n=39; 52.7%; Figure 3). Other countries with multiple programs identified included Germany

(n=7), Spain (n=4), Australia (n=3), France (n=3), Italy (n=3), and India (n=2). Only one program was identified in Hong Kong, Laos, New Zealand, Norway, Scotland, Singapore, Taiwan, Thailand, and England. Among 68 programs (91.9%) in which the model for the telemedicine program was reported, 56 (82.4%) operated with one central service (hub) providing telemedicine support to multiple spoke hospitals (i.e. “hub and spoke” model). There were three single-center programs, that is Aachen (Germany), Hallingdal (Norway), and Porto Alegre (Brazil), and one program in the East of England described as a horizontal hubless model (n=7 hospitals). We identified only three stroke telemedicine programs in the United States described to have nationwide coverage, including the Veteran’s Affairs National Telestroke Program (n=40 hospitals) TelaDoc Health (n=132 hospitals), and the Telespecialist company (n=171 hospitals). The largest identified stroke telemedicine program outside of the United States was based in India (100 hospitals: 22 hubs and 78 spokes).

### *Availability of mobile stroke unit pre-hospital ambulance services*

We identified 16 MSUs in the literature, 14 of which were already listed in the PRESTO repository of active MSUs. We found one MSU in Homburg (Germany) that appeared to be inactive (<https://www.mobile-stroke-unit.org/>), and another in Tennessee, USA, in implementation phase.<sup>145</sup> In the PRESTO registry, there were 30 active MSU services as of August 2023. Twenty-four of these active MSUs (80%) were located in HICs, 19 of which were located in the United States alone. Other MSU programs present in the American region included those located in Edmonton (Canada), Brasilia (Brazil), and Buenos Aires (Argentina). Four MSUs were present in Europe, all located in Western Europe, including in Berlin (Germany), Hamburg (Germany), Drobak (Norway), and Ipswich (England). Similarly, four MSUs were present in Southeast Asia, located in Assam (India), Coimbatore (India), Bangkok (Thailand), and Sichuan (China). There was one MSU present in Imo (Nigeria) and Melbourne (Australia).

## **Discussion**

In this article, we presented a repository of the latest published country-specific data on the availability of reperfusion services directly within hospitals, and those facilitated via telemedicine or MSU services. We described the worldwide coverage of reperfusion services and highlighted regions where data are limited or not recent.

The information presented likely illustrates disparities in access to reperfusion services for acute stroke around the world. Currently, the WSO is leading a call to action for increasing access to thrombectomy provision as a

**Table 1.** Availability of reperfusion services around the world.

Region	Country	Year	Registry source	Number of hospitals	Nationally representative	Intravenous thrombolysis		Endovascular thrombectomy	
						% AIS patients	% hospitals	Available	% AIS patients
Australasia <sup>8</sup>	Australia	2021	Yes	62	Yes	11.0	—	Yes	10.0
Australasia	New Zealand <sup>a</sup>	2022	Yes	28	Yes	12.8	100	Yes	7.2
Caribbean <sup>9</sup>	Barbados	2020	No	1	Yes	2.5	25	Unknown	—
Central Europe <sup>10</sup>	Bulgaria	2020	No	1	No	—	—	Yes	—
Central Europe <sup>11</sup>	Croatia	2006	No	—	No	—	—	Unknown	—
Central Europe <sup>12</sup>	Czech Republic	2020	Yes	45	Yes	17.4	—	Yes	5.5
Central Europe <sup>13</sup>	Hungary	2015	No	—	Yes	—	—	Unknown	—
Central Europe <sup>14</sup>	Poland	2017	No	181	Yes	12.9	100	Unknown	—
Central Europe <sup>15</sup>	Romania	2017	No	1	No	6.6	—	Unknown	—
Central Europe <sup>16</sup>	Serbia	2020	No	1	No	—	—	Yes	—
Central Europe <sup>17</sup>	Slovakia	2019	Yes	43	Yes	20.1	—	Yes	10.7
Central Europe <sup>18</sup>	Slovenia	2012	No	—	No	—	—	Yes	—
East Asia <sup>19</sup>	China	2020	Yes	280	Yes	8.6	—	Yes	3.4
East Asia <sup>20</sup>	Taiwan	2010	No	285	Yes	0.60	37	Unknown	—
Eastern Europe <sup>21</sup>	Bosnia and Herzegovina	2011	No	—	No	—	—	Unknown	—
Eastern Europe <sup>22</sup>	Estonia	2021	No	1	No	19.5	—	Yes	—
Eastern Europe <sup>23</sup>	Georgia	2019	No	—	No	3.4	—	Unknown	—
Eastern Europe <sup>24</sup>	Latvia	2020	Yes	1	No	17.8	—	Yes	1.2
Eastern Europe <sup>25</sup>	Lithuania	2018	No	—	No	—	—	Unknown	—
High-income Asia Pacific <sup>26</sup>	Japan	2019	Yes	130	Yes	—	—	Yes	—
High-income Asia Pacific	Singapore <sup>a</sup>	2022	Yes	4	Yes	7.1	100	Yes	2.0
High-income Asia Pacific <sup>27</sup>	South Korea	2014	Yes	16	Yes	15.2	—	Yes	4.6
High-income North America <sup>28</sup>	Canada	2019	Yes	136	No	12.2	33	Yes	—

(Continued)

Table 1. (Continued)

Region	Country	Year	Registry source	Number of hospitals	Nationally representative	Intravenous thrombolysis		Endovascular thrombectomy	
						% AIS patients	% hospitals	Available	% AIS patients
High-income North America <sup>29</sup>	USA	2018	Yes	646	Yes	19.1	—	Yes	4.1
Latin America <sup>30</sup>	Argentina	2018	No	1	No	11.0	—	Yes	3.0
Latin America <sup>31</sup>	Brazil	2019	No	1	No	15.0	—	Yes	8.0
Latin America <sup>32</sup>	Chile	2018	No	1	No	—	—	Unknown	—
Latin America <sup>33</sup>	Columbia	—	No	—	No	12.0	—	Unknown	—
Latin America <sup>34</sup>	Mexico	2018	No	1	No	8.9	—	Yes	—
Latin America <sup>35</sup>	Peru	2016	No	1	No	2.0	—	Unknown	—
North Africa and Middle East <sup>36</sup>	Egypt	2018	No	95	No	12.3	—	Yes	1.9
North Africa and Middle East <sup>37</sup>	Iran	—	No	—	No	—	—	Unknown	—
North Africa and Middle East <sup>38</sup>	Morocco	2017	No	1	No	8.4	—	No	—
North Africa and Middle East <sup>39</sup>	Oman	2018	No	1	No	11.9	—	Unknown	—
North Africa and Middle East <sup>40</sup>	Qatar	2010	No	—	No	9.0	—	Unknown	—
North Africa and Middle East <sup>41</sup>	Saudi Arabia	2018	No	1	No	8.6	—	Yes	—
North Africa and Middle East <sup>42</sup>	Turkey	2019	No	1	No	—	—	Yes	—
South Asia <sup>43</sup>	Bangladesh	2020	No	1	No	—	—	Unknown	—
South Asia <sup>44</sup>	India	2020	No	13	No	5.0	—	Yes	5.0
South Asia <sup>45</sup>	Nepal	2018	No	1	No	13.2	—	Unknown	—
South Asia <sup>46</sup>	Pakistan	2016	No	1	No	—	—	Unknown	—
Southeast Asia <sup>47</sup>	Indonesia	2020	No	1	No	2.4	—	Yes	—
Southeast Asia <sup>48</sup>	Malaysia	2016	Yes	15	Yes	—	16.5	Unknown	9
Southeast Asia <sup>49</sup>	Philippines	2016	No	10	No	1.3	—	Unknown	—
Southeast Asia <sup>50</sup>	Sri Lanka	2020	No	1	No	—	—	Unknown	—
Southeast Asia <sup>51</sup>	Thailand	2021	No	—	Yes	7.4	—	Unknown	—


(Continued)

Table 1. (Continued)

Region	Country	Year	Registry source	Number of hospitals	Nationally representative	Intravenous thrombolysis		Endovascular thrombectomy	
						% AIS patients	% hospitals	Available	% AIS patients
Southeast Asia <sup>52</sup>	Vietnam	2021	No	13	Yes	8.0	—	Yes	7.4
Southeast Asia	Brunei Darussalam	2021	No	—	—	10.0	40	Unknown	—
Southeast Asia <sup>53,54</sup>	Myanmar	2019	No	10	No	17.4	—	No	—
Sub-Saharan Africa <sup>55</sup>	Nigeria	2021	No	58	Yes	—	9	No	—
Sub-Saharan Africa <sup>56</sup>	South Africa	2017	No	8	No	1.0	25	Unknown	—
Western Europe <sup>57</sup>	Austria	2018	Yes	38	Yes	21.8	—	Unknown	4.0
Western Europe <sup>58</sup>	Belgium	2017	No	1	No	—	—	Unknown	—
Western Europe	Denmark <sup>a</sup>	2022	No	10	Yes	20.0	—	Yes	6.0
Western Europe <sup>59</sup>	Finland	2015	No	1	No	—	—	Unknown	—
Western Europe <sup>60</sup>	France	2019	No	147	Yes	9.0	—	Yes	7.1
Western Europe <sup>61</sup>	Germany	2017	No	—	Yes	15.9	—	Unknown	5.8
Western Europe <sup>62</sup>	Greece	2020	Yes	7	No	6.9	—	Unknown	—
Western Europe <sup>63</sup>	Ireland	2018	No	—	Unknown	—	—	Yes	—
Western Europe <sup>64</sup>	Israel	2007	No	—	No	—	—	Unknown	—
Western Europe <sup>65</sup>	Italy	2020	No	93	Yes	19.1	—	Yes	9.4
Western Europe <sup>66</sup>	Netherlands	2016	Yes	81	Yes	20.4	93	Yes	4.1
Western Europe <sup>67</sup>	Norway	2016	Yes	75	Yes	—	—	Unknown	—
Western Europe <sup>68,69</sup>	Portugal	2019	No	1	No	6.0	—	Yes	—
Western Europe <sup>70</sup>	Scotland	2021	Yes	—	Yes	18.0	—	Unknown	—
Western Europe <sup>71</sup>	Spain	2020	No	16	No	17.3	—	Yes	22.0
Western Europe	Sweden <sup>a</sup>	2022	Yes	72	Yes	13.9	—	Yes	7.0
Western Europe <sup>72</sup>	Switzerland	2020	Yes	25	Yes	22.1	—	Yes	15.0
Western Europe <sup>73</sup>	United Kingdom	2022	Yes	182	Yes	10.4	—	Yes	2.4

AIS: acute ischemic stroke.

<sup>a</sup>Unpublished and latest registry data verified by co-author.

**Figure 2.** Map showing the availability of intravenous thrombolysis and endovascular thrombectomy services around the world.

game-changing treatment for stroke. The policy objectives of the Stroke Action Plan for Europe also include targets by 2030 for guaranteeing access to reperfusion services to 95% of eligible patients, and achieving intravenous thrombolysis rates above 15% and endovascular therapy above 5% in all European countries.<sup>2</sup> Therefore, the data reported in this review are necessary for underpinning these objectives and targets, commensurate with investments in workforce and system improvements that are required to ensure equitable access to reperfusion services.

Telemedicine services facilitate access to intravenous thrombolysis, especially in more remote areas or smaller hospitals, with this service requiring a robust technical setup, both at the hub and spoke hospitals. Targets to provide all people within a country access to acute stroke reperfusion services may be ambitious. For example in France, a national strategy to have a stroke telemedicine service that began in 2011 is still in progress.<sup>146</sup> However, the required investments are economical compared with other solutions (e.g. MSUs or stroke air ambulances).<sup>147,148</sup> In addition, stroke telemedicine services are used in many regions, but nationwide approaches are rare, and solutions are often regionally based and not standardized. Significant progress has been made in the United States (2019), where it was estimated that 96% of the population had access to a stroke center or stroke telemedicine service.<sup>149</sup> Also, there is evidence from Germany that wider adoption of stroke telemedicine service is feasible and safe, with 38,895

consultations successfully undertaken across 14 telemedicine networks in nine German states.<sup>128</sup>

The availability and extent of MSU services varied widely within each country, and implementation was often limited to specific regions or cities. MSUs were more common in the United States, with several cities successfully integrating MSUs into their emergency response systems. Investments to establish MSUs are considerable, including costs of procuring equipment, training, and integrating into existing health care systems. However, the benefits offset the upfront investment with reductions in time to diagnosis and earlier provision of time-critical treatments, especially in settings requiring extensive time to travel to hospital.<sup>150</sup>

An important finding from this review is that contemporary patient-level data on availability of reperfusion services remain scarce, even in HICs. Based on published data, intravenous thrombolysis and endovascular thrombectomy services were notably absent in low-income countries. Potential barriers to providing reperfusion services in these countries include lack of necessary infrastructure, equipment, and trained personnel, and high costs associated with procuring reperfusion drugs/devices. For countries with no data available on provision of reperfusion services, it is difficult to ascertain current gaps in this important aspect of stroke in these countries. We also found huge variations in access to reperfusion services in HICs, which underlines the need for investment in robust stroke care services and infrastructure globally.

**Table 2.** Availability of stroke telemedicine services around the world.

Region	Country	Location	Telemedicine Program	Year	Hospitals	Spokes	Hubs
Australasia <sup>74</sup>	Australia	New South Wales	John Hunter Hospital	2015	4	3	1
Australasia <sup>75</sup>	Australia	Victoria	Victorian Stroke Telemedicine Program	2016	17	16	1
Australasia <sup>76</sup>	Australia	Wangaratta	Royal Melbourne Hospital	2009	2	1	1
Australasia <sup>77</sup>	New Zealand	Central New Zealand	Central Region Telestroke Network	2021	8	7	1
East Asia <sup>78</sup>	Taiwan	Changhua, Yunlin, and Nantou	Changhua Christian Hospital	2019	7	6	1
East Asia <sup>79</sup>	Hong Kong, China <sup>a</sup>	Hong Kong	Queen Elizabeth Hospital	2012	—	—	—
High-income Asia Pacific <sup>80</sup>	Singapore	Singapore	National Neuroscience Institute	2013	2	1	1
High-income North America <sup>81</sup>	Canada	Alberta	University of Alberta Hospital	2009	8	7	1
High-income North America <sup>82</sup>	Canada	Ontario	Ontario Telestroke Program	2017	—	—	—
High-income North America <sup>83</sup>	USA	Arizona, Florida, Iowa, Minnesota, and Wisconsin	Mayo Clinic Telestroke Network	2020	28	27	1
High-income North America <sup>84</sup>	USA	Arkansas	Arkansas Stroke Assistance through Virtual Emergency Support program	2016	49	48	1
High-income North America <sup>85</sup>	USA	California	Temecula Valley Hospital	2015	2	1	1
High-income North America <sup>86</sup>	USA	Delaware	Thomas Jefferson University Hospital	2013	30	29	1
High-income North America <sup>87</sup>	USA	Florida	Name unknown	2019	2	1	1
High-income North America <sup>88</sup>	USA	Florida and Arizona	Mayo Clinic	2014	15	13	2
High-income North America <sup>89</sup>	USA	Georgia	Georgia Regents Medical Center	2012	18	17	1
High-income North America <sup>90</sup>	USA	Georgia and South Carolina	REACH	—	34	32	2
High-income North America <sup>91</sup>	USA	Georgia	AcuteCare Telemedicine	2012	7	—	—
High-income North America <sup>92</sup>	USA	Hawai'i	Hawai'i Telestroke Program	2017	8	7	1
High-income North America <sup>93</sup>	USA	Illinois	Rush Tele-Stroke Network	—	11	10	1
High-income North America <sup>94</sup>	USA	Maryland	University of Maryland Medical Center	2001	—	—	1
High-income North America <sup>95</sup>	USA	Massachusetts	Massachusetts General Hospital	2008	13	12	1

(Continued)

Table 2. (Continued)

Region	Country	Location	Telemedicine Program	Year	Hospitals	Spokes	Hubs
High-income North America <sup>96</sup>	USA	Massachusetts, Maine, and New Hampshire	Massachusetts General Hospital Telestroke Network	2015	17	16	1
High-income North America <sup>97</sup>	USA	Massachusetts, New Hampshire, and Maine	Partners TeleStroke Consultation Network	2018	43	—	—
High-income North America <sup>98</sup>	USA	New Jersey	Overlook Medical Center	2016	7	6	1
High-income North America <sup>99</sup>	USA	North Eastern region	Name unknown	2019	21	20	1
High-income North America <sup>100</sup>	USA	North Eastern Region	Name unknown	2017	23	22	1
High-income North America <sup>101</sup>	USA	Northern California	Kaiser Permanente Northern California	2016	21	20	1
High-income North America <sup>102</sup>	USA	Pennsylvania	Penn State College of Medicine	2018	16	15	1
High-income North America <sup>71</sup>	USA	Pennsylvania	Name unknown	—	30	29	1
High-income North America <sup>103</sup>	USA	Pennsylvania	UPMC Presbyterian Hospital	—	3	2	1
High-income North America <sup>104</sup>	USA	Pennsylvania and New Jersey	Name unknown	2016	41	40	1
High-income North America <sup>105</sup>	USA	Pennsylvania, New Jersey, and Delaware	Thomas Jefferson University Hospital	2012	29	28	1
High-income North America <sup>106</sup>	USA	Pittsburgh	UPMC telestroke network	2009	13	12	1
High-income North America	USA	South Carolina	Medical University of South Carolina Telestroke Program	2019	27	26	1
High-income North America <sup>107</sup>	USA	South Carolina	Neuro-Direct telestroke network, Greenville Health System	2020	8	7	1
High-income North America <sup>108</sup>	USA	South Carolina	South Carolina Telestroke Network	2016	43	40	3
High-income North America <sup>109</sup>	USA	Southwest Ohio, Northern Kentucky, and Eastern Indiana	University of Cincinnati Stroke Team	2020	31	30	1
High-income North America <sup>110</sup>	USA	Texas	Lone Star stroke consortium telestroke registry	2019	18	17	1
High-income North America <sup>111</sup>	USA	West Virginia	West Virginia University	2018	9	8	1
High-income North America <sup>112</sup>	USA	Western New York Region	REACH	—	11	10	1
High-income North America <sup>113</sup>	USA <sup>a</sup>	Ohio	Name unknown	2017	26	—	—

(Continued)

Table 2. (Continued)

Region	Country	Location	Telemedicine Program	Year	Hospitals	Spokes	Hubs
High-income North America <sup>114</sup>	USA <sup>a</sup>	Sioux Falls, South Dakota	Avera eCare	—	155	—	—
High-income North America <sup>115</sup>	USA <sup>a</sup>	South Carolina	Grand Strand Medical Center	2018	—	—	—
High-income North America <sup>116</sup>	USA <sup>a</sup>		Nationwide, Veterans' Affairs National Telestroke Program	2021	50	—	—
High-income North America <sup>117</sup>	USA <sup>a</sup>		Nationwide, name unknown	2019	132	—	—
High-income North America <sup>118</sup>	USA <sup>a</sup>		Nationwide, Telespecialists	2020	171	—	—
Latin America <sup>119</sup>	Brazil	Rio de Janeiro	Hospital Pró-Cardíaco	2016	6	5	1
Latin America <sup>120</sup>	Brazil <sup>a</sup>	Porto Alegre	HCPA	2015	1	—	—
South Asia <sup>121</sup>	India	East Delhi	Name unknown	2017	7	6	1
South Asia <sup>122</sup>	India	India	Name unknown	2014	100	78	22
Southeast Asia <sup>123</sup>	Laos	Laos	Thailand	2017	3	2	1
Southeast Asia <sup>124</sup>	Thailand <sup>a</sup>	Thailand	Thammasat	2009	—	—	—
Western Europe <sup>125</sup>	France	Burgundy	Burgundy telestroke network	2014	13	11	2
Western Europe <sup>126</sup>	France	East France	University hospital of Nancy	2012	2	1	1
Western Europe <sup>127</sup>	France	Grand-East Region	Virtuall	2017	7	6	1
Western Europe <sup>128</sup>	Germany	Germany	German telemedical stroke networks	2020	186	155	31
Western Europe <sup>129</sup>	Germany	Bavaria	TEMPIS	2021	26	24	2
Western Europe <sup>130</sup>	Germany	North-Western Bavaria	TRANSIT Stroke Network	2019	10	7	3
Western Europe <sup>131</sup>	Germany	Saxony	SOS-NET	2012	15	14	1
Western Europe <sup>132</sup>	Germany	South West Germany	FAST	2020	8	7	1
Western Europe <sup>133</sup>	Germany	Swabia	TESS Project	2002	8	7	1
Western Europe <sup>134</sup>	Germany <sup>a</sup>	Aachen	Name unknown	2015	1	—	—
Western Europe <sup>135</sup>	Germany <sup>a</sup>	Thuringia	SATELIT	2015	—	—	—
Western Europe <sup>136</sup>	Italy	Grosseto	Siena	2018	2	1	1
Western Europe <sup>137</sup>	Italy	Lazio	Name unknown	2020	4	3	1
Western Europe <sup>138</sup>	Italy	Treviso	Treviso Hospital	2013	2	1	1

(Continued)

Table 2. (Continued)

Region	Country	Location	Telemedicine Program	Year	Hospitals	Spokes	Hubs
Western Europe <sup>139</sup>	Norway	Hallingdal	Hallingdal Local Medical Centre	2020	1	—	—
Western Europe <sup>140</sup>	Scotland	Lothian	NHS Lothian	2008	3	—	—
Western Europe <sup>141</sup>	Spain	Catalunya	Name unknown	2008	3	2	1
Western Europe <sup>142</sup>	Spain	Madrid	The Madrid Telestroke Project	2013	2	1	1
Western Europe <sup>143</sup>	Spain <sup>a</sup>	Andalusia	Centro Andaluz de Tele-ictus	2020	—	—	—
Western Europe <sup>144</sup>	United Kingdom <sup>b</sup>	East of England	East of England Stroke Telemedicine Partnership	2019	7	—	—

REACH: Respectful and Equitable Access to Comprehensive Health care; UPMC: University of Pittsburgh Medical Center; HCPA: Hospital de Clinicas de Porto Alegre; SOS-NET: Stroke Eastern Saxony Network; TESS: Telemedicine in Stroke in Swabia; TEMPS: Telemedicine Pilot Project for Integrative Stroke Care; TRANSIT: Transregional Network for Stroke Intervention with Telemedicine; FAST: Stroke Consortium Rhine-Neckar; SATELIT: stroke telemedicine network in Thuringia; NHS: National Health Service.

In some cases, the name of the hub was used as the name of the program; <sup>a</sup>Unknown model of operation; <sup>b</sup>Horizontal hubless model.


National stroke clinical registries are an important source of recent standardized data. To allow for reliable comparison of performance in stroke care (e.g. the provision of intravenous thrombolysis) within and between countries over time, the Action Plan for Stroke in Europe has recently established a tool (Stroke Service Tracker) for reporting stroke data in European countries on an annual basis.<sup>151</sup> This initiative also supports development of reliable stroke registries in European countries. Establishment of such tools in other regions could facilitate collection of data essential for reliably monitoring the provision of reperfusion services, globally.

There are also current initiatives to map the availability of reperfusion services in certain regions. In Europe, although there was an increase in reperfusion therapy rates in many countries between 2016 and 2019 (based on expert opinion), this was halted in 2020.<sup>152</sup> The authors highlighted the ongoing inequalities in acute stroke treatment across Europe.<sup>152</sup> Similarly, there is variability in the standard of acute stroke care reported in Latin America, with reperfusion services reported to be available for a small proportion of the population in several countries.<sup>153,154</sup> In addition, efforts to map reperfusion services are being conducted in Southeast Asia (personal communication with DADS).

Much can be learned from countries identified in this report that represent similar socio-demographic and economic profiles. International collaborations and partnerships between HICs, and lower-middle-income or low-income countries could promote knowledge transfer, training opportunities, and resource sharing. One such initiative is the international stroke telemedicine service. This new model of care appears feasible and has the potential to improve access to specialist care, and timely reperfusion services, in regions (e.g. lower-middle-income and low-income countries) where stroke specialist input is limited.<sup>123,155</sup> These alliances may build local capacity, improve infrastructure, and enhance stroke care networks.

There are some limitations of this review. For nearly half of the countries, the most recent data on the availability of intravenous thrombolysis, endovascular thrombectomy, and stroke telemedicine services were collected prior to 2019. We acknowledge there may be delays in publishing recent data that may have impeded our ability to know with certainty the current gaps in these aspects of stroke care, and our findings may not fully represent the current state of stroke care services in all countries. Indeed, there is evidence from surveys of national organizations that reperfusion services are available in more countries than identified in this review.<sup>152–154</sup> Our findings rely on published patient-level data, and there may be bias in favor of countries with more research infrastructure or motivation to publish data on stroke services. Where possible, data from publications were supplemented by unpublished and verified data available from co-authors or trusted websites (e.g. PRESTO and

**Figure 3.** Map illustrating the number of stroke telemedicine services around the world. The size of the dot in a location reflects the number of services available in that location.



RES-Q repositories) which are routinely updated. Also, some countries may have made significant strides in acute stroke care that have not yet been published or are published in languages other than English. The quality and reliability of data may also vary across different countries and sources. To keep the repository up-to-date and to assist us in updating this review, we encourage readers to submit any relevant unpublished country-specific data on reperfusion services to the corresponding author.

## Conclusion

Despite remarkable advancements in reperfusion therapies for stroke, it is evident from published data that their availability and provision remain unevenly distributed globally. Bridging this treatment gap requires a multifaceted approach, including advocacy on the importance of reperfusion therapies to drive policy changes and secure additional funding for stroke programs, investment in infrastructure, workforce training, awareness campaigns, and collaborative efforts. By providing an updated repository of the latest country-specific data on stroke care services, this review can inform future policy and funding decisions aimed at improving access to evidence-based stroke care worldwide, and can inform organizations, such as the WSO, to advocate for gaps in coverage to be addressed. By prioritizing stroke care as

a global health priority, we can strive toward equitable access to reperfusion therapies and improve stroke outcomes worldwide.

## Acknowledgements

Analyses and interpretations of the data are of the authors alone.

## Author contributions

J.K., M.T.O., T.T., and D.A.C. contributed equally. J.K., M.T.O., and T.T. contributed to the design, undertook literature search, data collection, data analyses, and interpretation, wrote the first draft of the article. D.A.C. contributed to the design, data interpretation, and wrote the first draft of the article, and revised the article. All authors interpreted the data, provided additional data reports as relevant, revised and approved the final version of the article.

## Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: D.A.C. reports Research Fellowship support from the National Health and Medical Research Council (grant no. 1154273).

### ORCID iDs

Joosup Kim  <https://orcid.org/0000-0002-4079-0428>  
 Muideen T Olaiya  <https://orcid.org/0000-0002-4070-0533>  
 Deidre A De Silva  <https://orcid.org/0000-0002-7123-2381>  
 Diana A De Sousa  <https://orcid.org/0000-0002-6702-7924>  
 Anna Ranta  <https://orcid.org/0000-0002-3223-3330>  
 Geoffrey A Donnan  <https://orcid.org/0000-0001-6324-3403>  
 Valery Feigin  <https://orcid.org/0000-0002-6372-1740>  
 David J Werring  <https://orcid.org/0000-0003-2074-1861>  
 Mayowa Owolabi  <https://orcid.org/0000-0003-1146-3070>  
 Jeyaraj Pandian  <https://orcid.org/0000-0003-0028-1968>  
 Tharshanah Thayabaranathan  <https://orcid.org/0000-0003-2504-7772>  
 Dominique A Cadilhac  <https://orcid.org/0000-0001-8162-682X>

### References

- Langhorne P, Audebert HJ, Cadilhac DA, Kim J and Lindsay P. Stroke systems of care in high-income countries: what is optimal? *Lancet* 2020; 396: 1433–1442.
- Norrving B, Barrick J, Davalos A, et al. Action plan for stroke in Europe 2018-2030. *Eur Stroke J* 2018; 3: 309–336.
- Kim J, Thayabaranathan T, Donnan GA, et al. Global stroke statistics 2019. *Int J Stroke* 2020; 15: 819–838.
- Thayabaranathan T, Kim J, Cadilhac DA, et al. Global stroke statistics 2022. *Int J Stroke* 2022; 17: 946–956.
- Thrift AG, Cadilhac DA, Thayabaranathan T, et al. Global stroke statistics. *Int J Stroke* 2014; 9: 6–18.
- Thrift AG, Thayabaranathan T, Howard G, et al. Global stroke statistics. *Int J Stroke* 2017; 12: 13–32.
- The PRE-hospital Stroke Treatment Organization (PRESTO), <https://www.prestosu.org/i4a/pages/index.cfm?pageid=1> (accessed January 2023).
- Cadilhac DA, Dalli LL, Morrison JL, et al. *The Australian stroke clinical registry annual report 2021*. Report no. 14, 22 December 2022, 66 pp. Heidelberg, VIC, Australia: The Florey Institute of Neuroscience and Mental Health.
- Cardiovascular Disease in Barbados. *Annual report 2022*, <http://www.bnr.org.bb/cms/home/37-home/91-bnr-annual-reports>
- Tsalta-Mladenov M and Andonova S. Health-related quality of life after ischemic stroke: impact of sociodemographic and clinical factors. *Neurol Res* 2021; 43: 553–561.
- Vuletić V, Dikanović M, Lezaić Z, Sapina L and Kadojić D. Are we ready for intravenous thrombolysis in acute stroke treatment in our region. *Acta Clin Croat* 2011; 50: 145–148.
- Sedova P, Brown RD, Jr, Bryndziar T, et al. Treat COVID-19, but not only COVID-19: stroke matters as well. *Cerebrovasc Dis* 2022; 51: 52–59.
- Folyovich A, Biczó D, Jarecsny T, et al. Daylight saving time and the incidence of thrombolysis to treat acute ischemic stroke. *Rev Neurol* 2020; 176: 361–365.
- Maluchnik M, Ryglewicz D, Sienkiewicz-Jarosz H, et al. Differences in acute ischaemic stroke care in Poland: analysis of claims database of National Health Fund in 2017. *Neurol Neurochir Pol* 2020; 54: 449–455.
- Uivarosan D, Bungau S, Tit DM, et al. Financial burden of stroke reflected in a pilot center for the implementation of thrombolysis. *Medicina* 2020; 56: 54.
- Ždraljević M, Pekmezović T, Stanarčević P, et al. Influence of thrombocytopenia on the outcome of mechanical thrombectomy in patients with acute ischemic stroke. *J Stroke Cerebrovasc Dis* 2022; 31: 106240.
- Gdovinová Z, Vitková M, Baráková A and Cvpová A. The impact of the COVID-19 outbreak on acute stroke care in Slovakia: data from across the country. *Eur J Neurol* 2021; 28: 3263–3266.
- Jeromel M, Milošević Z, Zaletel M, Žvan B, Švigelj V and Oblak JP. Endovascular therapy for acute stroke is a safe and efficient evolving method: a single-center retrospective analysis. *J Vasc Interv Radiol* 2015; 26: 1025–1030.
- Zhao J, Li H, Kung D, Fisher M, Shen Y and Liu R. Impact of the COVID-19 epidemic on stroke care and potential solutions. *Stroke* 2020; 51: 1996–2001.
- Hsieh CY, Chen CH, Chen YC and Kao Yang Y-H. National survey of thrombolytic therapy for acute ischemic stroke in Taiwan 2003-2010. *J Stroke Cerebrovasc Dis* 2013; 22: e620–e627.
- Smajlović D, Salihović D, Avdić L, Dostović Z, Ibrahimagić OČ and Vidović M. Intravenous thrombolytic therapy for acute ischemic stroke in Tuzla Canton, Bosnia and Herzegovina. *Int J Stroke* 2013; 8: E48.
- Gross-Paju K, Thomson U, Adlas R, et al. Implementation of the Helsinki model at West Tallinn Central Hospital. *Medicina* 2022; 58: 1173.
- Katsarava Z, Akhvediani T, Janelidze T, et al. Establishing stroke services in the Republic of Georgia. *Eur Neurol* 2022; 85: 56–64.
- Karelis G, Micule M, Klavina E, et al. The Riga East University Hospital stroke registry—an analysis of 4915 consecutive patients with acute stroke. *Medicina* 2021; 57: 632.
- Jucevičiūtė N, Mikužis P and Balnytė R. Absolute blood eosinophil count could be a potential biomarker for predicting haemorrhagic transformation after intravenous thrombolysis for acute ischaemic stroke. *BMC Neurol* 2019; 19: 127.
- Toyoda K, Yoshimura S, Nakai M, et al. Twenty-year change in severity and outcome of ischemic and hemorrhagic strokes. *JAMA Neurol* 2022; 79: 61–69.
- Kim JY, Kang K, Kang J, et al. *Stroke statistics in Korea 2018: a report from the epidemiology research council of the Korean Stroke Society*. Korean Stroke Society, [https://www.stroke.or.kr/bbs/skin/notice\\_popup/download.php?code=epidemic1&number=5305](https://www.stroke.or.kr/bbs/skin/notice_popup/download.php?code=epidemic1&number=5305) (2018, accessed April 2023).
- Pardhan A, Sharp S, Yearwood K, et al. *Ontario and Local Health Integration Network (LHIN) 2017/18 stroke report cards and progress reports*. CorHealth, 2019, <https://www.corhealthontario.ca/Ontario-&-LHIN-2017-18-Stroke-Report-Cards-&-Progress-Reports.pdf>
- Asaithambi G, Tong X, Lakshminarayan K, Coleman King SM and George MG. Current trends in the acute treatment of ischemic stroke: analysis from the Paul Coverdell

- National Acute Stroke Program. *J Neurointerv Surg* 2020; 12: 574–578.
30. Alet M, Lucci FR and Ameriso S. Mechanical thrombectomy for reperfusion of acute ischemic stroke in a Stroke Unit in Argentina. *Arq Neuropsiquiatr* 2020; 78: 39–43.
  31. Do Rosário CF, Fernandes Neto WG, Pessotti AL, et al. Epidemiological analysis of stroke patients with emphasis on access to acute-phase therapies. *Arq Neuropsiquiatr* 2022; 80: 117–124.
  32. Brunser AM, Mazzon E, Cavada G, et al. Low doses of alteplase, for ischemic stroke after Enchanted and its determinants, a single center experience. *Arq Neuropsiquiatr* 2020; 78: 681–686.
  33. Gil-Rojas Y and Lasalvia P. Budgetary impact analysis of alteplase—recombinant tissue plasminogen activator (rtPA)—as a thrombolytic treatment for acute ischemic stroke in Colombia. *Expert Rev Pharmacoecon Outcomes Res* 2022; 22: 1095–1102.
  34. Gonzalez-Aquines A, Cordero-Pérez AC, Cristobal-Niño M, Pérez-Vázquez G and Góngora-Rivera F; GECEN Investigators. Contribution of onset-to-alarm time to pre-hospital delay in patients with ischemic stroke. *J Stroke Cerebrovasc Dis* 2019; 28: 104331.
  35. Abanto C, Valencia A, Calle P, et al. Challenges of thrombolysis in a developing country: characteristics and outcomes in Peru. *J Stroke Cerebrovasc Dis* 2020; 29: 104819.
  36. Aref H, Zakaria M, Shokri H, Roushdy T, El Basiouny A and El Nahas N. Changing the landscape of stroke in Egypt. *Cerebrovasc Dis Extra* 2021; 11: 155–159.
  37. Khazaei M, Davoodian A, Taheri M and Ghafouri-Fard S. Former antiplatelet drug administration and consequences of intravenous thrombolysis in acute ischemic stroke. *Hum Antibodies* 2020; 28: 53–56.
  38. Acherqui M, Khattab H, Habtany Y, et al. Assessment of eligibility for thrombolysis in acute ischaemic stroke patients in Morocco. *Pan Afr Med J* 2020; 36: 351.
  39. Al Harthi HA, Al Kashmiri A, Zakaryia LM, et al. Clinical profile of stroke patients presenting to the Emergency Department of a Major Stroke Centre in Oman. *Sultan Qaboos Univ Med J* 2022; 22: 91–97.
  40. Ibrahim F, Deleu D, Akhtar N, et al. Burden of stroke in qatar. *J Stroke Cerebrovasc Dis* 2015; 24(12): 2875–2879.
  41. Al Khathaami AM, Al Bdah B, Tarawneh M, et al. Utilization of intravenous tissue plasminogen activator and reasons for nonuse in acute ischemic stroke in Saudi Arabia. *J Stroke Cerebrovasc Dis* 2020; 29: 104761.
  42. Kömürçü HF, Gözke E, Doğan Ak P, Aslan IK, Salt I and Bi Er ÇİÖ. Changes in neutrophil, lymphocyte, platelet ratios and their relationship with NIHSS after rtPA and/or thrombectomy in ischemic stroke. *J Stroke Cerebrovasc Dis* 2020; 29: 105004.
  43. Hasan ATMH, Das SC, Islam MS, et al. Impact of COVID-19 on hospital admission of acute stroke patients in Bangladesh. *PLoS One* 2021; 16: e0240385.
  44. Mathew T, John SK, Sarma G, et al. COVID-19-related strokes are associated with increased mortality and morbidity: a multicenter comparative study from Bengaluru, South India. *Int J Stroke* 2021; 16: 429–436.
  45. Nepal G, Yadav JK, Basnet B, Shrestha TM, Kharel G and Ojha R. Status of prehospital delay and intravenous thrombolysis in the management of acute ischemic stroke in Nepal. *BMC Neurol* 2019; 19: 155.
  46. Zaman Babar MU, Khan AZ, Hakim H, Gilani J and Wasay M. Utilization and outcomes with low dose tissue plasminogen activator as intravenous thrombolytic therapy for ischaemic stroke at Aga Khan University Hospital, Karachi: a retrospective analysis. *J Pak Med Assoc* 2019; 69: 1705–1710.
  47. Mesiano T, Kurniawan M, Saputri KM, et al. Endovascular treatment in acute ischemic stroke adoption and practice: a single-center Indonesian experience. *Cerebrovasc Dis Extra* 2021; 11: 72–76.
  48. Aziz ZA and Sidek NN. *Annual report of the Malaysian stroke registry 2009-2016*. Kuala Terengganu, Malaysia: Clinical Research Centre, 2017.
  49. Navarro JC, San Jose MC, Collantes E, et al. Stroke thrombolysis in the Philippines. *Neurol Asia* 2018; 23: 115–120.
  50. Herath HMM, Rodrigo C, Alahakoon A, et al. Outcomes of stroke patients undergoing thrombolysis in Sri Lanka; an observational prospective study from a low-middle income country. *BMC Neurol* 2021; 21: 434.
  51. Tiamkao S, Ienghong K, Cheung LW, Celebi I, Suzuki T and Apiratwarakul K. Stroke incidence, rate of thrombolytic therapy, mortality in Thailand from 2009 to 2021. *Open Access Maced J Med Sci* 2022; 9: 110–115.
  52. Mai DT, Dao XC, Luong NK, Nguyen TK, Nguyen HT and Nguyen TN. Current state of stroke care in Vietnam. *Stroke: Vasc Interv Neurol* 2022; 2: e000331.
  53. Ohnmar Kyaw M, Shew ZM, et al. The pattern and burden of neurological disorders: a systemic review of Neurology Department, Yangon General Hospital, Myanmar. *Neurol Asia* 2020; 25: 555–561.
  54. Venketasubramanian N, Khine YM, Ohnmar O, Khin MPPK and Win MT. Burden of stroke in Myanmar. *Cerebrovasc Dis Extra* 2021; 11: 49–51.
  55. Arabambi B, Oshinaike O, Ogun SA, et al. Stroke units in Nigeria: a report from a nationwide organizational cross-sectional survey. *Pan Afr Med J* 2022; 42: 140.
  56. Mandizvidza V, London L and Bryer A. Shortfall in stroke care: a study of ischaemic stroke care practices in a South African metropole. *S Afr Med J* 2021; 111: 215–219.
  57. Marko M, Posekany A, Szabo S, et al. Trends of r-tPA (Recombinant Tissue-Type Plasminogen Activator) treatment and treatment-influencing factors in acute ischemic stroke. *Stroke* 2020; 51: 1240–1247.
  58. Vanhoucke J, Hemelsoet D, Achten E, et al. Impact of a code stroke protocol on the door-to-needle time for IV thrombolysis: a feasibility study. *Acta Clin Belg* 2020; 75: 267–274.
  59. Varjoranta T, Raatiniemi L, Majamaa K, Martikainen M and Liisanantti JH. Prehospital and hospital delays for stroke patients treated with thrombolysis: a retrospective study from mixed rural-urban area in Northern Finland. *Australas Emerg Care* 2019; 22: 76–80.
  60. Olié V, Grave C, Tuppin P, Duloquin G, Béjot Y and Gabet A. Patients hospitalized for ischemic stroke and intracerebral hemorrhage in France: time trends (2008-2019), in-hospital outcomes, age and sex differences. *J Clin Med* 2022; 11: 1669.

61. Weber R, Krogias C, Eyding J, et al. Age and sex differences in ischemic stroke treatment in a nationwide analysis of 1.11 million hospitalized cases. *Stroke* 2019; 50: 3494–3502.
62. Siskou O, Korompoki E, Ntaios G, et al. Access of stroke patients' to optimal healthcare technology in Greece: messages to policy makers. *Stud Health Technol Inform* 2020; 272: 421–424.
63. Griffin E, Murphy S, Sheehan M, et al. Early repatriation post-thrombectomy: a model of care which maximises the capacity of a stroke network to treat patients with large vessel ischaemic stroke. *J Neurointerv Surg* 2020; 12: 1166–1171.
64. Gur AY, Shopin L and Bornstein NM. Lessons learned from 2 years experience in intravenous thrombolysis for acute ischemic stroke in a single tertiary medical center. *Isr Med Assoc J* 2009; 11: 714–716, 718.
65. Manganotti P, Naccarato M, Scali I, et al. Stroke management during the coronavirus disease 2019 (COVID-19) pandemic: experience from three regions of the north east of Italy (Veneto, Friuli-Venezia-Giulia, Trentino-Alto-Adige). *Neurol Sci* 2021; 42: 4599–4606.
66. Kuhrij LS, Wouters MW, Van den Berg-Vos RM, De Leeuw FE and Nederkoorn PJ. The Dutch Acute Stroke Audit: benchmarking acute stroke care in the Netherlands. *Eur Stroke J* 2018; 3: 361–368.
67. Varndal T, Indredavik B, Phan A, Fjærtøft H. Hjerneslag i Norge 2015-16 - behandling og resultater [Stroke in Norway 2015-16 - treatment and outcomes]. *Tidsskr Nor Laegeforen* 2020; 140(2). Doi: 10.4045/tidsskr.19.0246.
68. Sobral S, Taveira I, Seixas R, et al. Late hospital arrival for thrombolysis after stroke in Southern Portugal: who is at risk? *J Stroke Cerebrovasc Dis* 2019; 28: 900–905.
69. Rosa JL, Alves M, Ferreira P, Papoila AL and Nunes AP. Previous disability and benefit of acute phase therapy in functional prognosis of selected patients with ischemic stroke. *J Stroke Cerebrovasc Dis* 2022; 31: 106183.
70. Scottish Stroke Improvement Programme 2022: national report. A management information release for Scotland, 2022, <https://www.publichealthscotland.scot/publications/scottish-stroke-improvement-programme/scottish-stroke-improvement-programme-2022-national-report/>
71. Tejada Meza H, Lambea Gil Á, Saldaña AS, et al. Impact of COVID-19 outbreak on ischemic stroke admissions and in-hospital mortality in North-West Spain. *Int J Stroke* 2020; 15: 755–762.
72. De Marchis GM, Wright PR, Michel P, et al. Association of the COVID-19 outbreak with acute stroke care in Switzerland. *Eur J Neurol* 2022; 29: 724–731.
73. The road to recovery: sentinel Stroke National Audit Programme annual report 2022, <https://www.pslhub.org/learn/patient-safety-in-health-and-care/the-road-to-recovery-sentinel-stroke-national-audit-programme-annual-report-2022-10-november-2022-r8164/> (accessed 10 November 2022).
74. Demeestere J, Sewell C, Rudd J, et al. The establishment of a telestroke service using multimodal CT imaging decision assistance: “Turning on the fog lights.” *J Clin Neurosci* 2017; 37: 1–5.
75. Bladin CF, Kim J, Bagot KL, et al. Improving acute stroke care in regional hospitals: clinical evaluation of the Victorian Stroke Telemedicine program. *Med J Aust* 2020; 212: 371–377.
76. Nagao KJ, Koschel A, Haines HM, Bolitho LE and Yan B. Rural Victorian Telestroke project. *Intern Med J* 2012; 42: 1088–1095.
77. Mahawish K, Gommans J, Kleinig T, Lallu B, Tyson A and Ranta A. Switching to Tenecteplase for stroke thrombolysis: real-world experience and outcomes in a regional stroke network. *Stroke* 2021; 52: e590–e593.
78. Lin CH, Lee KW, Chen TC, et al. Quality and safety of Telemedicine in acute ischemic stroke: early experience in Taiwan. *J Formos Med Assoc* 2022; 121: 314–318.
79. Fong WC, Ismail M, Lo JW, et al. Telephone and teleradiology-guided thrombolysis can achieve similar outcome as thrombolysis by neurologist on-site. *J Stroke Cerebrovasc Dis* 2015; 24: 1223–1228.
80. Mong R, Tiah L, Wong M and Tan C. Improving telestroke treatment times through a quality improvement initiative in a Singapore emergency department. *Singapore Med J* 2019; 60: 69–74.
81. Khan K, Shuaib A, Whittaker T, et al. Telestroke in Northern Alberta: a two year experience with remote hospitals. *Can J Neurol Sci* 2010; 37: 808–813.
82. Porter J, Hall RE, Kapral MK, Fang J, Khan F and Silver FL. Outcomes following telestroke-assisted thrombolysis for stroke in Ontario, Canada. *J Telemed Telecare* 2018; 24: 492–499.
83. Huang JF, Greenway MRF, Nasr DM, et al. Telestroke in the time of COVID-19: the Mayo Clinic experience. *Mayo Clin Proc* 2020; 95: 1704–1708.
84. Nalleballe K, Sharma R, Kovvuru S, et al. Why are acute ischemic stroke patients not receiving thrombolysis in a telestroke network? *J Telemed Telecare* 2020; 26: 317–321.
85. Agrawal K, Raman R, Ernstrom K, et al. Accuracy of stroke diagnosis in telestroke-guided tissue plasminogen activator patients. *J Stroke Cerebrovasc Dis* 2016; 25: 2942–2946.
86. Zanuty M, Chalouhi N, Starke RM, et al. Epidemiology of a large telestroke cohort in the Delaware valley. *Clin Neurol Neurosurg* 2014; 125: 143–147.
87. Cheshire WP, Barrett KM, Eidelman BH, et al. Patient perception of physician empathy in stroke telemedicine. *J Telemed Telecare* 2021; 27: 572–581.
88. Demaerschalk BM, Boyd EL, Barrett KM, et al. Comparison of stroke outcomes of hub and spoke hospital treated patients in Mayo Clinic telestroke program. *J Stroke Cerebrovasc Dis* 2018; 27: 2940–2942.
89. Bruno A, Lanning KM, Gross H, Hess DC, Nichols FT and Switzer JA. Timeliness of intravenous thrombolysis via telestroke in Georgia. *Stroke* 2013; 44: 2620–2622.
90. Switzer JA, Singh R, Mathiassen L, Waller JL, Adams RJ and Hess DC. Telestroke: variations in intravenous thrombolysis by spoke hospitals. *J Stroke Cerebrovasc Dis* 2015; 24: 739–744.
91. Sanders KA, Figiel C, Kiely JM, Gwynn MW and Johnston LH. Expanding access to intravenous tissue-type plasminogen activator treatment with a practice-based telestroke system. *J Stroke Cerebrovasc Dis* 2013; 22: e546–e548.

92. Chaffin HM, Nakagawa K and Koenig MA. Impact of statewide telestroke network on acute stroke treatment in Hawai'i. *Hawaii J Health Soc Welf* 2019; 78: 280–286.
93. Lee VH, Cutting S, Song SY, et al. Participation in a telestroke program improves timeliness of intravenous thrombolysis delivery. *Telemed e-Health* 2016; 23: 60–62.
94. LaMonte MP, Bahouth MN, Hu P, et al. Telemedicine for acute stroke. *Stroke* 2003; 34: 725–728.
95. Pervez MA, Silva G, Masrur S, et al. Remote supervision of IV-tPA for acute ischemic stroke by telemedicine or telephone before transfer to a regional stroke center is feasible and safe. *Stroke* 2010; 41: e18–e24.
96. Moreno A, Schwamm LH, Siddiqui KA, et al. Frequent hub-spoke contact is associated with improved spoke hospital performance: results from the Massachusetts General Hospital telestroke network. *Telemed J E Health* 2018; 24: 678–683.
97. Sharma R, Zachrisson KS, Viswanathan A, et al. Trends in telestroke care delivery: a 15-year experience of an academic hub and its network of spokes. *Circ Cardiovasc Qual Outcomes* 2020; 13: e005903.
98. Belt GH, Felberg RA, Rubin J and Halperin JJ. In-transit telemedicine speeds ischemic stroke treatment. *Stroke* 2016; 47: 2413–2415.
99. Gadhia R, Schwamm LH, Viswanathan A, Whitney C, Moreno A and Zachrisson KS. Evaluation of the experience of spoke hospitals in an academic telestroke network. *Telemed J E Health* 2019; 25: 584–590.
100. Yu AT, Regenhardt RW, Whitney C, et al. CTA protocols in a telestroke network improve efficiency for both spoke and hub hospitals. *AJNR Am J Neuroradiol* 2021; 42: 435–440.
101. Nguyen-Huynh MN, Klingman JG, Avins AL, et al. Novel telestroke program improves thrombolysis for acute stroke across 21 hospitals of an integrated healthcare system. *Stroke* 2018; 49: 133–139.
102. Sinha K and Bettermann K. Clinical outcomes of moderate to severe acute ischemic stroke in a telemedicine network. *J Clin Neurosci* 2019; 69: 160–165.
103. Sobhani F, Desai S, Madill E, et al. Remote longitudinal inpatient acute stroke care via telestroke. *J Stroke Cerebrovasc Dis* 2021; 30: 105749.
104. Sweid A, Atallah E, Saad H, et al. Correlation between pre-admission blood pressure and outcome in a large telestroke cohort. *J Clin Neurosci* 2019; 62: 33–37.
105. Chalouhi N, Dressler JA, Kunkel ES, et al. Intravenous tissue plasminogen activator administration in community hospitals facilitated by telestroke service. *Neurosurgery* 2013; 73: 667–671; discussion 671–672.
106. Zaidi SF, Jumma MA, Urna XN, et al. Telestroke-guided intravenous tissue-type plasminogen activator treatment achieves a similar clinical outcome as thrombolysis at a comprehensive stroke center. *Stroke* 2011; 42: 3291–3293.
107. Poupore N, Edrissi C, Sowah M, et al. Stroke severity among men and women acute ischemic stroke patients in the telestroke network. *Cerebrovasc Dis Extra* 2022; 12: 93–101.
108. Simpson AN, Harvey JB, DiLembo SM, et al. Population health indicators associated with a statewide telestroke program. *Telemed J E Health* 2020; 26: 1126–1133.
109. Hsiao J, Sayles E, Antzoulatos E, et al. Effect of COVID-19 on emergent stroke care: a regional experience. *Stroke* 2020; 51: e2111–e2114.
110. Reddy ST, Savitz SI, Friedman E, et al. Patients transferred within a telestroke network for large-vessel occlusion. *J Telemed Telecare* 2022; 28: 595–602.
111. Adcock AK, Choi J, Alvi M, et al. Expanding acute stroke care in rural America: a model for statewide success. *Telemed J E Health* 2020; 26: 865–871.
112. Ionita CC, Sharma J, Janicke DM, et al. Acute ischemic stroke and thrombolysis location: comparing telemedicine and stroke center treatment outcomes. *Hosp Pract* 2009; 37: 33–39.
113. Lee VH, Howell R, Yadav R, Heaton S, Wiles KL and Lakhani S. Thrombolysis of stroke mimics via telestroke. *Stroke Vasc Neurol* 2022; 7: 267–270.
114. Mohr NM, Young T, Harland KK, et al. Telemedicine is associated with faster diagnostic imaging in stroke patients: a cohort study. *Telemed J E Health* 2019; 25: 93–100.
115. Gutovitz S, Leggett J, Hart L, Leaman SM, James H and Stillinger T. The impact of teleneurologists on acute stroke care at an advanced primary stroke centre. *J Telemed Telecare* 2022; 28: 115–121.
116. Lyerly MJ, Daggy J, LaPradd M, et al. Impact of telestroke implementation on emergency department transfer rate. *Neurology* 2022; 98: e1617–e1625.
117. Zachrisson KS, Sharma R, Wang Y, Mehrotra A and Schwamm LH. National trends in telestroke utilization in a US commercial platform prior to the COVID-19 pandemic. *J Stroke Cerebrovasc Dis* 2021; 30: 106035.
118. Sevilis T, McDonald M, Avila A, et al. Telestroke: maintaining quality acute stroke care during the COVID-19 pandemic. *Telemed J E Health* 2022; 28: 481–485.
119. Carvalho VS, Jr, Picanço MR, Volschan A and Bezerra DC. Impact of simulation training on a telestroke network. *Int J Stroke* 2019; 14: 500–507.
120. Martins SO, Mont'Alverne F, Rebello LC, et al. Thrombectomy for stroke in the public health care system of Brazil. *N Engl J Med* 2020; 382: 2316–2326.
121. Kushwaha S, Talwar P, Chandel N, Anthony A, Maheshwari S and Khurana S. Saving the brain initiative—developing an effective hub-and-spoke model to improve the acute stroke management pathways in urban India. *J Neurol Sci* 2018; 393: 83–87.
122. Srivastava PV, Sudhan P, Khurana D, et al. Telestroke a viable option to improve stroke care in India. *Int J Stroke* 2014; 9: 133–134.
123. Chutinet A, Keosodsay S, Vorasayan P, et al. The first 10 thrombolysis for acute ischemic stroke in Lao People's Democratic Republic under teleconsultation from Thailand. *J Stroke Cerebrovasc Dis* 2019; 28: 104327.
124. Muengtawepong S, Dharmasaroja PA, Maungboon P and Wattanaruangkowit P. Feasibility and safety of remote radiology interpretation with telephone consultation for acute stroke in Thailand. *Neurol India* 2010; 58: 740–742.
125. Legris N, Hervieu-Bègue M, Daubail B, et al. Telemedicine for the acute management of stroke in Burgundy, France: an evaluation of effectiveness and safety. *Eur J Neurol* 2016; 23: 1433–1440.

126. Richard S, Lavandier K, Zioueche Y, Pelletier S, Vezain A and Ducrocq X. Use of telemedicine to manage severe ischaemic strokes in a rural area with an elderly population. *Neurol Sci* 2014; 35: 683–685.
127. Kaminsky AL, Mione G, Omorou Y, et al. Outcome of patients with large vessel occlusion stroke after first admission in telestroke spoke versus comprehensive stroke center. *J Neurointerv Surg* 2020; 12: 753–757.
128. Vollmuth C, Miljukov O, Abu-Mugheisib M, et al. Impact of the coronavirus disease 2019 pandemic on stroke teleconsultations in Germany in the first half of 2020. *Eur J Neurol* 2021; 28: 3267–3278.
129. Hubert GJ, Kraus F, Maegerlein C, et al. The “Flying intervention team”: a novel stroke care concept for rural areas. *Cerebrovasc Dis* 2021; 50: 375–382.
130. Gabriel KMA, Jírů-Hillmann S, Kraft P, et al. Two years’ experience of implementing a comprehensive telemedical stroke network comprising in mainly rural region: the Transregional Network for Stroke Intervention with Telemedicine (TRANSIT-Stroke). *BMC Neurol* 2020; 20: 104.
131. Zerna C, Siepmann T, Barlinn K, et al. Association of time on outcome after intravenous thrombolysis in the elderly in a telestroke network. *J Telemed Telecare* 2016; 22: 18–24.
132. Busetto L, Sert M, Herzog F, et al. “But it’s a nice compromise”—qualitative multi-centre study of barriers and facilitators to acute telestroke cooperation in a regional stroke network. *Eur J Neurol* 2022; 29: 208–216.
133. Wiborg A and Widder B; Telemedicine in Stroke in Swabia Project. Teleneurology to improve stroke care in rural areas. *Stroke* 2003; 34: 2951–2956.
134. Quadflieg LTM, Beckers SK, Bergrath S, et al. Comparing the diagnostic concordance of tele-EMS and on-site-EMS physicians in emergency medical services: a retrospective cohort study. *Sci Rep* 2020; 10: 17982.
135. Klingner C, Tinschert P, Brodoehl S, et al. The effect of endovascular thrombectomy studies on the decision to transfer patients in a telestroke network. *Telemed JE Health* 2020; 26: 388–394.
136. Gallerini S, Marsili L, Groccia V, et al. Appropriateness, safety, and effectiveness of “drip and ship” teleconsultation model in Southeastern Tuscany: a feasibility study. *Neurol Sci* 2020; 41: 2961–2965.
137. Brunetti V, Broccolini A, Caliendo P, et al. Effect of the COVID-19 pandemic and the lockdown measures on the local stroke network. *Neurol Sci* 2021; 42: 1237–1245.
138. Nardetto L, Dario C, Tonello S, et al. A one-to-one telestroke network: the first Italian study of a web-based telemedicine system for thrombolysis delivery and patient monitoring. *Neurol Sci* 2016; 37: 725–730.
139. Kjelle E and Myklebust AM. Telemedicine remote controlled stroke evaluation and treatment, the experience of radiographers, paramedics and junior doctors in a novel rural stroke management team. *BMC Health Serv Res* 2021; 21: 554.
140. Kerr E, Dennis M and Keir S. Thrombolysis: attempting to reduce postcode prescribing in Scotland? *Int J Stroke* 2010; 5: 486–488.
141. Pedragosa A, Alvarez-Sabín J, Rubiera M, et al. Impact of telemedicine on acute management of stroke patients undergoing endovascular procedures. *Cerebrovasc Dis* 2012; 34: 436–442.
142. Martínez-Sánchez P, Miralles A, Sanz de Barros R, et al. The effect of telestroke systems among neighboring hospitals: more and better? The Madrid Telestroke Project. *J Neurol* 2014; 261: 1768–1773.
143. Montaner J, Barragán-Prieto A, Pérez-Sánchez S, et al. Break in the stroke chain of survival due to COVID-19. *Stroke* 2020; 51: 2307–2314.
144. Evans NR, Sibson L, Day DJ, Agarwal S, Shekhar R and Warburton EA. Hyperacute stroke thrombolysis via telemedicine: a multicentre study of performance, safety and clinical efficacy. *BMJ Open* 2022; 12: e057372.
145. Alexandrov AW, Arthur AS, Bryndziar T, et al. High-resolution CT with arch/neck/head CT angiography on a mobile stroke unit. *J Neurointerv Surg* 2022; 14: 623–627.
146. Ohannessian R, Schott AM, Colin C, Nighoghossian N, Medeiros de Bustos E and Moulin T. Acute telestroke in France: a systematic review. *Rev Neurol* 2020; 176: 316–324.
147. Fassbender K, Merzou F, Lesmeister M, et al. Impact of mobile stroke units. *J Neurol Neurosurg Psychiatry* 2021; 92: 815–822.
148. Walter S, Zhao H, Easton D, et al. Air-Mobile Stroke Unit for access to stroke treatment in rural regions. *Int J Stroke* 2018; 13: 568–575.
149. Zachrisson KS, Cash RE, Adeoye O, et al. Estimated population access to acute stroke and telestroke centers in the US, 2019. *JAMA Netw Open* 2022; 5: e2145824.
150. Chen J, Lin X, Cai Y, Huang R, Yang S and Zhang G. A systematic review of mobile stroke unit among acute stroke patients: time metrics, adverse events, functional result and cost-effectiveness. *Front Neurol* 2022; 13: 803162.
151. Stroke Service Tracker (SST), <https://actionplan.eso-stroke.org/stroke-service-tracker-2> (accessed July 2023).
152. Aguiar de Sousa D, Wilkie A, Norrving B, et al. Delivery of acute ischaemic stroke treatments in the European region in 2019 and 2020. *Eur Stroke J* 2023; 8: 618–628.
153. Ouriques Martins SC, Sacks C, Hacke W, et al. Priorities to reduce the burden of stroke in Latin American countries. *Lancet Neurol* 2019; 18: 674–683.
154. Martins SCO, Lavados P, Secchi TL, et al. Fighting against stroke in Latin America: a joint effort of medical professional societies and governments. *Front Neurol* 2021; 12: 743732.
155. Ranta A, Whitehead M, Gunawardana C, et al. International telestroke: the first five cases. *J Stroke Cerebrovasc Dis* 2016; 25: e44–e45.