



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Chen, Y;Zhu, Z;Wang, W;Shang, X;He, M;Li, J

Title:

Association of Myopia With Risk of Incident Metabolic Syndrome: Findings From the UK Biobank Study Cohort of 91,591 Participants

Date:

2022-05-16

Citation:

Chen, Y., Zhu, Z., Wang, W., Shang, X., He, M. & Li, J. (2022). Association of Myopia With Risk of Incident Metabolic Syndrome: Findings From the UK Biobank Study Cohort of 91,591 Participants. *Frontiers in Medicine*, 9, <https://doi.org/10.3389/fmed.2022.872013>.

Persistent Link:

<https://hdl.handle.net/11343/316298>

License:

[CC BY](#)



Association of Myopia With Risk of Incident Metabolic Syndrome: Findings From the UK Biobank Study Cohort of 91,591 Participants

Yanxian Chen¹, Zhuoting Zhu², Wei Wang³, Xianwen Shang^{2,4}, Mingguang He^{1,2,3,4} and Jinying Li^{1*}

¹ Department of Ophthalmology, Peking University Shenzhen Hospital, Shenzhen Peking University-The Hong Kong University of Science and Technology Medical Center, Shenzhen, China, ² Department of Ophthalmology, Guangdong Academy of Medical Sciences, Guangdong Provincial People's Hospital, Guangzhou, China, ³ State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University, Guangzhou, China, ⁴ Centre for Eye Research Australia, Ophthalmology, Department of Surgery, University of Melbourne, Melbourne, VIC, Australia

Purpose: To investigate the association between myopia and risk of metabolic syndrome (MetS) in a prospective cohort from the UK Biobank Study.

Methods: Volunteers (aged 40 years and above) free of baseline MetS and cataract included from the UK Biobank Study, a prospective follow-up cohort. Myopia was defined using unmydriatic autorefractometry, self-report-myopia, and medical records for refractive error at baseline. MetS as well as components of MetS were diagnosed based on health records, blood biochemistry, and questionnaires. Questionnaires determined the status of smoking, drinking, physical activity and dietary supplements, as well as ethnicity and education.

Results: A total of 91,591 participants were available in the analysis, with a mean age of 55.37 ± 8.07 years at baseline and a median follow-up years of 11.16 years. The proportion of myopia was 49.7%, and a total of 937 (1.0%) participants were identified as having incident MetS (0.09/100 person years). Subjects with myopia were more likely to have MetS compared with non-myopic subjects (0.82 vs. 0.21%, Log-rank test $P < 0.001$). Myopes had greater risk of incident MetS (Hazard ratio [HR] = 4.19, 95% confidence interval [CI] 3.57–4.93, $P < 0.001$) adjusting for baseline age, gender, education and ethnicity. After further controlling for lifestyle factors (smoking, drinking, physical activity, and fish oil supplement) or baseline metabolic disorders, the risk of incident MetS were 3.88- and 4.06-fold greater in myopic subjects than those without myopia, respectively ($P < 0.001$ for both models). The severity of myopia was not significantly correlated to incident MetS in multivariate-adjusted models.

Conclusions: An increased risk of incident MetS among the elderly is associated with myopia, but not the degree of myopia. These findings highlighted the need of prevention of MetS among older adults with myopia.

Keywords: myopia, metabolic syndrome, older population, Western lifestyle, cohort study

OPEN ACCESS

Edited by:

Michele Lanza,
University of Campania Luigi
Vanvitelli, Italy

Reviewed by:

Juncheng Dai,
Nanjing Medical University, China
Hong Peng Sun,
Soochow University, China

*Correspondence:

Jinying Li
ljyszbdy@163.com

Specialty section:

This article was submitted to
Ophthalmology,
a section of the journal
Frontiers in Medicine

Received: 09 February 2022

Accepted: 04 April 2022

Published: 16 May 2022

Citation:

Chen Y, Zhu Z, Wang W, Shang X,
He M and Li J (2022) Association of
Myopia With Risk of Incident Metabolic
Syndrome: Findings From the UK
Biobank Study Cohort of 91,591
Participants. *Front. Med.* 9:872013.
doi: 10.3389/fmed.2022.872013

INTRODUCTION

In recent decades, myopia has become a major ocular condition leading to visual impairment with a rapidly increasing prevalence, especially in areas with rapid urbanization, including South Korea (1), Taiwan (2), and mainland China (3). It has been estimated that the global prevalence of myopia will be as high as 50% by 2050 (4). Environmental factors predominantly contribute to the current rise in prevalent myopia, including intensive near work and less time outdoors (5, 6), which are common patterns of activities in modern lifestyle, and may continue to impact behavioral aspects of living (7, 8).

Metabolic syndrome (MetS) is a metabolic disorder involving central obesity, high blood pressure, hyperglycemia and dyslipidemia, closely relating to Western obesogenic diet and lifestyle (9). It has been long postulated that there may be common pathogenetic basis for metabolic disorder and myopia due to the wide acceptance of Western lifestyle over the last few decades (10–12). But the association between myopia and the individual components of MetS have been reported with some inconsistency. For example, in the Handan Eye Study, diabetes mellitus was a risk factor for myopia in the elderly (13), but the Blue Mountains Eye study (14) and the Beaver Dam Eye Study (15) showed no significant association between myopia and diabetes. Elevated systolic blood pressure was observed in highly myopic patients (16) and myopic children (17), while hypertension was not associated with myopia in the Tanjong Pagar Survey (18). There have also been conflicting associations between obesity (or increased body mass index) and myopia (19–21). Heretofore the association between MetS and myopia has not fully identified in existing studies.

In this study, we aimed to investigate the association of myopia and different degree of myopia with incident MetS in a large-scale prospective cohort from the UK Biobank Study.

MATERIALS AND METHODS

Population

The UK Biobank is a large-scale prospective cohort study of health and wellbeing enrolling more than 500,000 participants aged 40–69 years at baseline recruitment between 2006 and 2010 throughout the United Kingdom. Detailed methodology for the UK Biobank Study has been described previously (22). In brief, about 9.2 million subjects aged 40–69 years in 22 assessment centers were invited to participate, and data from 502,505 people were collected (response rate 5.5%). The baseline assessment visit including comprehensive questionnaires, physical and functional measures and collection of biological samples. Ophthalmic assessments were conducted in 6 assessment centers. Non-cycloplegic autorefractometry was performed with the Tomey rc-5000 Auto Refkeratometer (Tomey, Nagoya, Japan) and up to 10 measurements were taken for each eye. A detailed procedure can be seen elsewhere (23). Follow-up visits were performed in 2012–2013, 2014, and 2019.

Ethical approval was obtained from the National Information Governance Board for Health and Social Care and the NHS North West Multicenter Research Ethics

Committee. Written informed consent was obtained from all participants.

Definitions

Myopia was defined using non-cycloplegic autorefractometry and questionnaires data. Subjects with spherical equivalent (sphere + 1/2 cylinder) < -0.50D in either eye were considered to have myopia. For subjects without autorefractometry data, hospital health record, self-reported eye disease and reason for glasses in questionnaires were used to identify status of refractive error. High myopia was defined as an SE < -6.0D, or with self-reported high myopia in questionnaires. Participants who underwent refractive surgery for myopia were also included as myopes. We excluded subjects who had cataract, amblyopia or strabismus diagnosis from hospital record and previous cataract surgery.

In the current study, MetS was defined using the International Diabetes Federation (IDF) criteria (24): (1) Waist circumference > 94 cm in European men and >80 cm in European women [for Chinese and South Asian ethnicity was >90 cm for men and >80 cm for women, adapted from Kendal DM et al. (25)]; (2) Blood pressure > 130/85 mmHg or treatment of previously diagnosed hypertension; (3) Triglyceride > 1.7 mmol/l or specific treatment for lipid abnormality; (4) HDL-C <1.0 mmol/l in men, <1.3 mmol/l in women or specific treatment for lipid abnormality; (5) Fasting plasma glucose > 5.6 mmol/l or previously diagnosed type 2 diabetes. Diagnosis of MetS was considered when the individual met criterion (1) plus any two of the rest disorders. The level of triglyceride, HDL-C and glucose were available in blood biochemistry data. History of treatment for hypertension, diabetes, or lipid abnormality were extracted from questionnaires.

Incident MetS in follow-up period were identified as the proportion of subjects who had no MetS at baseline and who subsequently developed MetS during the follow-up period. Follow-up time was defined as the duration between the date of baseline visit and the date of following status which was the earliest: incident MetS, lost to follow-up, or death.

Potential confounders associating with MetS or myopia were also identified in the analysis. Information of smoking status (previous/current and never), drinking status (previous/current and never) at baseline, educational level (college/university/NVQ and others), ethnicity (white and non-white) were obtained from questionnaires. Level of physical activity was identified as low, moderate and high using International Physical Activity Questionnaire (IPAQ), and further categorized to above moderate/vigorous/walking recommendation and not, according to Metabolic Equivalent Task scores based on IPAQ guidelines (26). Information of dietary supplements were collected from questionnaires and recorded as fish oil supplement and others. Hypertension was defined as a systolic blood pressure >130 mmHg or a diastolic blood pressure >80 mmHg, or a self-reported hypertension, or taking antihypertensive drugs. Diabetes mellitus was defined as a glycosylated hemoglobin level of >6.5%, or a self-reported diabetes mellitus, or taking medication for diabetes mellitus.

Statistical Analysis

We used *t* tests to determine differences in baseline age between participants with myopia and without myopia, and between those with mild/moderate myopia and with high myopia. Pearson's χ^2 test were used for comparison of categorical variables. Survival distribution was compared using Log-rank test, and Cox proportional hazards regression models was used to assess the associations between covariates and incident MetS. All analysis were performed using Stata (version 16.0, StataCorp).

RESULTS

Among the 502,505 participants in the baseline UK Biobank Study, myopia diagnosis was identified in 65,875 subjects: 44,373 with available autorefractometry data, 1,361 with history of refractive surgery for myopia, 2,753 with self-reported myopia and 17,388 with hospital medical record of myopia diagnosis. As showed in **Figure 1**, after excluding subjects with cataract ($n = 41,349$), baseline MetS ($n = 126,147$) and missing data on refraction status (without autorefractometry data, answers to the questions regarding refractive status,

and medical records of refractive error, $n = 241,468$) or baseline data of MetS ($n = 1,950$), a total of 91,591 subjects were available in the analysis, among which 45,487 were myopic (49.7%) and 3,650 were highly myopic (4.0%). Subjects included in our analysis were younger, had more females, more college graduates, and less white people compared with the 410,914 subjects who were excluded from the analysis (**Supplementary Table 1**).

Baseline characteristics between myopes and non-myopes were summarized in **Table 1**. Compared with non-myopic group, myopic group had a significantly younger baseline age (54.32 ± 7.82 years vs. 56.40 ± 8.19 years, $P < 0.001$), more females (55.1 vs. 55.9%, $P = 0.026$), and a higher proportion of subjects with college degree (52.6 vs. 38.9%), more white people (92.0 vs. 90.0%, $P < 0.001$), more central obese (42.1 vs. 35.9%, $P < 0.001$), more hypertensive (48.3 vs. 47.5%, $P = 0.022$), less smokers (37.8 vs. 45.2%, $P < 0.001$), more drinkers (96.4 vs. 95.2%, $P < 0.001$), less physically active (82.2 vs. 84.9%, $P < 0.001$), less fish oil supplement (29.3 vs. 30.6%, $P < 0.001$). The distributions of gender, diabetes mellitus, high triglyceride and low HDL-C showed no significant difference between myopic and non-myopic group.




FIGURE 1 | Flowchart depicting inclusion of participants in this study.

TABLE 1 | Baseline characteristics of participants with and without myopia.

Variable	No myopia (n = 46,104)	Myopia (n = 45,487)	Total (n = 91,591)	P
Baseline age, year, mean (sd)	56.40 (8.19)	54.32 (7.82)	55.37 (8.07)	<0.001
Female, n (%)	25,759 (55.9%)	25,082 (55.1%)	50,841 (55.5%)	0.026
College/university degree, n (%)	17,921 (38.9%)	23,915 (52.6%)	41,836 (45.7%)	<0.001
Ethnicity, n (%)				<0.001
White	41,491 (90.0%)	41,851 (92.0%)	83,342 (91.0%)	
Others	4,613 (10.0%)	3,636 (8.0%)	8,249 (9.0%)	
Central obesity, n (%)	19,393 (42.1%)	16,337 (35.9%)	35,730 (39.0%)	<0.001
Hypertension, n (%)	21,904 (47.5%)	21,954 (48.3%)	43,858 (47.9%)	0.022
Diabetes Mellitus, n (%)	659 (1.4%)	621 (1.4%)	1,280 (1.4%)	0.408
High triglyceride, n (%)	13,420 (29.1%)	13,053 (28.7%)	26,473 (28.9%)	0.169
Low HDL-C, n (%)	2,974 (7.7%)	3,053 (8.0%)	6,027 (7.9%)	0.060
Smoker, n (%)	20,705 (45.2%)	17,123 (37.8%)	37,828 (41.5%)	<0.001
Drinker, n (%)	43,697 (95.2%)	43,754 (96.4%)	87,451 (95.8%)	<0.001
PA above moderate/vigorous/walking recommendation, n (%)	31,868 (84.9%)	31,629 (82.2%)	63,497 (83.5%)	<0.001
Fish oil supplement, n (%)	14,087 (30.6%)	13,189 (29.3%)	27,276 (30.0%)	<0.001

PA, physical activity.

In subjects with high myopia, the proportion of female was higher (58.5 vs. 55.1%, $P < 0.001$), educational level was higher (college graduates 59.6 vs. 49.8%, $P < 0.001$), proportion of central obesity was smaller (34.5 vs. 38.2%, $P < 0.001$), high triglyceride was lower (21.9 vs. 25.2%, $P < 0.001$) and number of smokers was less (33.0 vs. 39.6%, $P < 0.001$) compared with subjects with mild/moderate myopia (**Supplementary Table 2**).

There were 937 participants (1.0%, 0.09/100 person years) having incident MetS during follow-up period. The median (IQR) follow-up post baseline enrollment was 11.16 (0.55) years. **Table 2** demonstrates baseline characteristics of participants with and without incident MetS. Subjects with incident MetS tended to be older (57.18 ± 7.09 years vs. 55.34 ± 8.08 years, $P < 0.001$), have less females (44.3 vs. 55.6%, $P < 0.001$), and more white people (97.0 vs. 90.9%, $P < 0.001$). The proportion of myopes in subjects with incident MetS was significantly higher (79.7 vs. 49.4%, $P < 0.001$) than in those without incident MetS, while the degree of myopia was similar between the two groups. The percentage of baseline central obesity, hypertension, diabetes mellitus, high triglyceride and low HDL-C were significantly higher in subjects with incident MetS than in those without ($P < 0.001$ for all), and the level of physical activity was lower (77.6 vs. 83.6%, $P < 0.001$), but the number of smokers and drinkers did not differ significantly between the two groups.

The incidence of MetS in myopes (0.82%, 0.14/100 person years) were significantly higher than those without baseline myopia (0.21%, 0.03/100 person years, Log-rank test $P < 0.001$), while incident MetS in different degree of myopia was not significantly different (0.13% [0.04/100 person years] in mild/moderate myopia vs. 0.02% [0.05/100 person years] in high myopia, Log-rank test $P = 0.1315$). We performed three different models to assess the contribution of myopia and myopia degree to incident MetS: Model 1 adjusted for baseline age, gender, ethnicity and education; Model 2

adjusted for variables in Model 1 plus smoking, drinking, physical activity level and fish oil supplement; Model 3 adjusted for variables in Model 1 plus baseline central obesity, hypertension, low HDL-C and high triglyceride. The coefficients with 95% confidence intervals (CIs) of each individual covariate in Cox proportional hazards regression model for incident MetS were listed in **Supplementary Table 3**. In summary, baseline age, baseline myopia, baseline central obesity, baseline hypertension, baseline diabetes, baseline high triglyceride and baseline low HDL-C level were independently associated with incident MetS. As seen in **Figure 2**, myopic subjects had a 4.19-fold higher risk of developing MetS (95%CI 3.57–4.93) in Model 1, and a 3.88-fold risk (95%CI 3.25–4.63) in Model 2. After further adjusting for baseline metabolic disorders, myopes still had a 3-times higher risk of future MetS (Hazard ratio 4.06, 95%CI 3.36–4.90) than non-myopes. However, high myopia at baseline was not a significant risk factor of incident MetS when adjusting for age, gender, education, ethnicity, lifestyle factors, or baseline metabolic disorders.

Sensitivity analysis was conducted within different subgroups of participants. As shown in **Supplementary Figure 1**, after excluding participants with college degree or above, or non-white ethnicity, or obesity/hypertension/smoker/low level of physical activity at baseline, the results were comparable to the main analysis, verifying the robustness of our findings.


DISCUSSION

In this large community-based follow-up study, being myopic at baseline was a significant risk factor of incident MetS later in life, and the association remained significant after adjusting for confounders. However, the severity of myopia was not independently associated with future risk of MetS.

TABLE 2 | Baseline characteristics of participants with and without incident metabolic syndrome.

	No incident MetS (n = 90,654)	Incident MetS (n = 937)	P
Baseline age, year, mean (sd)	55.34 (8.08)	57.18 (7.09)	<0.001
Female, n (%)	50,426 (55.6%)	415 (44.3%)	<0.001
College/university degree, n (%)	41,380 (45.7%)	456 (48.7%)	0.065
Ethnicity, n (%)			<0.001
White	82,433 (90.9%)	909 (97.0%)	
Others	8,221 (9.1%)	28 (2.9%)	
Myopia, n (%)	44,740 (49.4%)	747 (79.7%)	<0.001
High myopia, n (%)	3,860 (12.8%)	21 (14.9%)	0.451
Central obesity, n (%)	35,190 (38.8%)	540 (57.6%)	<0.001
Hypertension, n (%)	43,044 (47.5%)	814 (86.9%)	<0.001
Diabetes Mellitus, n (%)	1,228 (1.4%)	52 (5.6%)	<0.001
High triglyceride, n (%)	25,609 (28.3%)	864 (92.2%)	<0.001
Low HDL, n (%)	5,927 (7.8%)	100 (14.1%)	<0.001
Smoker, n (%)	37,436 (41.5%)	392 (41.9%)	0.795
Drinker, n (%)	86,544 (95.8%)	907 (96.8%)	0.114
PA above moderate/vigorous/walking recommendation, n (%)	62,895 (83.6%)	602 (77.6%)	<0.001
Fish oil supplement, n (%)	26,982 (29.9%)	294 (32.7%)	0.066

MetS, metabolic syndrome; PA, physical activity.



To our best knowledge, the current analysis is the first study reporting the association between myopia and incident MetS. In consistent with other studies, incident MetS was strongly

associated with baseline metabolic disorders, including obesity (27, 28), higher systolic blood pressure (28) and low HDL-C (29). Nevertheless, we found that myopia remained to be a

significant predictor for incident MetS after controlling all the baseline metabolic disorders. Diabetes mellitus is a common metabolic abnormality related to myopia, mostly attribute to the influence of glucose level on lens power (30, 31). However, the prevalence of diabetes mellitus was only 1.4% at baseline in the current study. Therefore, we confirmed that myopia is associated with risk of developing MetS as an independent risk factor, not as a result of change in lens power due to metabolic disorders.

The insignificant association between degree of myopia and incident MetS was also observed in the current analysis. The proportion of high myopia in this study was 4.0%, similar to a 2–5% prevalence of high myopia in older population from population-based researches, where high myopia was considered to be more genetic in etiology (32–35). We can speculate that severe myopia is not correlated to incident MetS due to the weak connection with environmental factors. However, this possibility may need to be validated in a cohort consisted of congenital and acquired high myopia due to the lack of differentiation from types of myopia herein.

Several potential mechanisms may explain the association between myopia and the risk of MetS. Firstly, myopia and MetS share common environmental risk factors. Decreased outdoor activity has been well-recognized as a key determinant of myopia development (5, 36) and teenagers and young adults with myopia tend to spend less time on physical activity and more time sedentary or use of screen-based devices (8, 12, 37). Our results also suggested that adults with myopia tend to have a lower level of moderate/vigorous physical activity. Meanwhile, being more physically active significantly reduces risk for metabolic syndrome (38), which is in consistent with the current analysis. It is plausible that physical activity patterns in myopes influences the rate of MetS development later in life. Secondly, there might be interrelation in the pathogenesis between myopia and MetS. Vascular inflammation is commonly seen in pathophysiology of obesity (39, 40), diabetes mellitus (41, 42) as well as in myopia progression (3, 43). In addition, insulin has been found to promote myopia development both *in vitro* (44) and *in vivo* studies (45, 46). Polymorphism of insulin-like growth factor-1 (IGF-1), a downstream production of insulin, also is associated with severe myopia (47).

As the first study investigating the association between myopia and incident MetS, the implications of our findings are far reaching. From a socio-economic perspective, tremendous amounts of resources have been spent on prevention and mitigation of metabolic disorders (48–50). The association between myopia and incident MetS identified in our study indicates the burden on public health will be far more huge when the young people with higher prevalence of myopia grow older. From the perspective regarding service for individual patients as part of a shift toward “precision medicine,” active procedures are warranted in the future to reduce the risk of incident MetS in the elderly with myopia, including accessibility and uptake of regular medical checkup, as well as adapted instructions and exercise therapy for myopes.

Strengths of our study include the large-scale cohort from whom detailed health records identifying MetS and refractive errors were collected, and a detailed assessment of associations between incident MetS and potential risk factors. The main flaw of this study is the lack of axial length data, limiting further investigation of the association of MetS with axial myopia and lens-induced myopia. Future studies with a more comprehensive eye examination might be needed to explore underlying mechanism of the established association in this study. Another potential limitation is that we may not exclude all the confounders especially for the factors related to lifestyle, including sleep duration, time spent on outdoor activities or near work.

CONCLUSIONS

In summary, we found that myopic subjects were at an increased risk of incident MetS controlling for baseline confounders, but the severity of myopia showed little influence on future development of MetS. The findings suggest the significance of prevention of MetS in older adults with myopia.

DATA AVAILABILITY STATEMENT

The data analyzed in this study is subject to the following licenses/restrictions: UK Biobank data are available only for applicants. Requests to access these datasets should be directed to <https://www.ukbiobank.ac.uk>.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the National Information Governance Board for Health and Social Care and the NHS North West Multicenter Research Ethics Committee. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

YC and JL devised the project. MH and XS assisted with project developing. ZZ and WW conducted data processing and analysis. YC wrote the manuscript. JL and MH reviewed and revised the manuscript. All authors contributed to the article and approved the submitted version.

ACKNOWLEDGMENTS

YC received support from Science and Technology Planning Project of Shenzhen Municipality (JCYJ20210324110000002).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmed.2022.872013/full#supplementary-material>

REFERENCES

- Jung SK, Lee JH, Kakizaki H, Jee D. Prevalence of myopia and its association with body stature and educational level in 19-year-old male conscripts in Seoul, South Korea. *Invest Ophthalmol Vis Sci.* (2012) 53:5579–83. doi: 10.1167/iovs.12-10106
- Lin LL, Shih YF, Hsiao CK, Chen CJ. Prevalence of myopia in Taiwanese schoolchildren: 1983 to 2000. *Ann Acad Med Singap.* (2004) 33:27–33.
- Wei CC, Kung YJ, Chen CS, Chang CY, Lin CJ, Tien PT, et al. Allergic conjunctivitis-induced retinal inflammation promotes myopia progression. *EBioMedicine.* (2018) 28:274–86. doi: 10.1016/j.ebiom.2018.01.024
- Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, et al. Global prevalence of myopia and high myopia and temporal trends from 2000 through 2050. *Ophthalmology.* (2016) 123:1036–42. doi: 10.1016/j.ophtha.2016.01.006
- He M, Xiang F, Zeng Y, Mai J, Chen Q, Zhang J, et al. Effect of time spent outdoors at school on the development of myopia among children in China: a randomized clinical trial. *JAMA.* (2015) 314:1142–8. doi: 10.1001/jama.2015.10803
- Huang HM, Chang DS, Wu PC. The association between near work activities and myopia in children—A systematic review and meta-analysis. *PLoS ONE.* (2015) 10:e0140419. doi: 10.1371/journal.pone.0140419
- Baird PN, Saw SM, Lanca C, Guggenheim JA, Smith III EL, Zhou X, et al. Myopia. *Nat Rev Dis Primers.* (2020) 6:99. doi: 10.1038/s41572-020-00231-4
- Hansen MH, Laigaard PP, Olsen EM, Skovgaard AM, Larsen M, Kessel L, et al. Low physical activity and higher use of screen devices are associated with myopia at the age of 16–17 years in the CCC2000 Eye Study. *Acta Ophthalmol.* (2020) 98:315–21. doi: 10.1111/aos.14242
- Yang Q, Cogswell ME, Flanders WD, Hong Y, Zhang Z, Loustalot F, et al. Trends in cardiovascular health metrics and associations with all-cause and CVD mortality among US adults. *JAMA.* (2012) 307:1273–83. doi: 10.1001/jama.2012.339
- Cordain L, Eaton SB, Brand Miller J, Lindeberg S, Jensen C. An evolutionary analysis of the aetiology and pathogenesis of juvenile-onset myopia. *Acta Ophthalmol Scand.* (2002) 80:125–35. doi: 10.1034/j.1600-0420.2002.800203.x
- Galvis V, López-Jaramillo P, Tello A, Castellanos-Castellanos YA, Camacho PA, Cohen DD, et al. Is myopia another clinical manifestation of insulin resistance? *Med Hypotheses.* (2016) 90:32–40. doi: 10.1016/j.mehy.2016.02.006
- Jacobsen N, Jensen H, Goldschmidt E. Does the level of physical activity in university students influence development and progression of myopia?—A 2-year prospective cohort study. *Invest Ophthalmol Vis Sci.* (2008) 49:1322–7. doi: 10.1167/iovs.07-1144
- Liang YB, Wong TY, Sun LP, Tao QS, Wang JJ, Yang XH, et al. Refractive errors in a rural Chinese adult population the Handan eye study. *Ophthalmology.* (2009) 116:2119–27. doi: 10.1016/j.ophtha.2009.04.040
- Guzowski M, Wang JJ, Rochtchina E, Rose KA, Mitchell P. Five-year refractive changes in an older population: the Blue Mountains Eye Study. *Ophthalmology.* (2003) 110:1364–70. doi: 10.1016/S0161-6420(03)00465-2
- Lee KE, Klein BE, Klein R, Wong TY. Changes in refraction over 10 years in an adult population: the Beaver Dam Eye study. *Invest Ophthalmol Vis Sci.* (2002) 43:2566–71.
- Chen SJ, Cheng CY, Li AF, Peng KL, Chou P, Chiou SH, et al. Prevalence and associated risk factors of myopic maculopathy in elderly Chinese: the Shihpai eye study. *Invest Ophthalmol Vis Sci.* (2012) 53:4868–73. doi: 10.1167/iovs.12-9919
- Zhao M, Wang W, Yu H, Ma Y, Zheng L, Zhang L, et al. Elevated blood pressure is associated with higher prevalence of low visual acuity among adolescent males in Northeast China. *Sci Rep.* (2017) 7:15990. doi: 10.1038/s41598-017-14252-9
- Foong AW, Wong TY, Saw SM, Foster PJ. Hypermetropia, axial length, and hypertension: the Tanjong Pagar survey. *Am J Ophthalmol.* (2006) 141:1142–4. doi: 10.1016/j.ajo.2006.01.020
- Harrington SC, Stack J, O'Dwyer V. Risk factors associated with myopia in schoolchildren in Ireland. *Br J Ophthalmol.* (2019) 103:1803–9. doi: 10.1136/bjophthalmol-2018-313325
- Jonas JB, Wang N, Wang YX, You QS, Yang D, Xie X, et al. Subfoveal choroidal thickness and cerebrospinal fluid pressure: the Beijing Eye Study 2011. *Invest Ophthalmol Vis Sci.* (2014) 55:1292–8. doi: 10.1167/iovs.13-13351
- Roy A, Kar M, Mandal D, Ray RS, Kar C. Variation of axial ocular dimensions with age, sex, height, BMI-, and their relation to refractive status. *J Clin Diagn Res.* (2015) 9:AC01–4. doi: 10.7860/JCDR/2015/10555.5445
- Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, et al. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med.* (2015) 12:e1001779. doi: 10.1371/journal.pmed.1001779
- Chua SYL, Thomas D, Allen N, Lotery A, Desai P, Patel P, et al. Cohort profile: design and methods in the eye and vision consortium of UK Biobank. *BMJ Open.* (2019) 9:e025077. doi: 10.1136/bmjopen-2018-025077
- Zimmet P, Magliano D, Matsuzawa Y, Alberti G, Shaw J. The metabolic syndrome: a global public health problem and a new definition. *J Atheroscler Thromb.* (2005) 12:295–300. doi: 10.5551/jat.12.295
- Kendall DM, Sobel BE, Coulston AM, Peters Harmel AL, McLean BK, Peragallo-Dittko V, et al. The insulin resistance syndrome and coronary artery disease. *Coron Artery Dis.* (2003) 14:335–48. doi: 10.1097/01.mca.0000076512.29238.2a
- Cassidy S, Chau JY, Catt M, Bauman A, Trenell MI. Cross-sectional study of diet, physical activity, television viewing and sleep duration in 233,110 adults from the UK Biobank; the behavioural phenotype of cardiovascular disease and type 2 diabetes. *BMJ Open.* (2016) 6:e010038. doi: 10.1136/bmjopen-2015-010038
- De Silva ST, Niriella MA, Ediriweera DS, Kottahachchi D, Kasturiratne A, de Silva AP, et al. Incidence and risk factors for metabolic syndrome among urban, adult Sri Lankans: a prospective, 7-year community cohort, follow-up study. *Diabetol Metab Syndr.* (2019) 11:66. doi: 10.1186/s13098-019-0461-7
- Thuyen TQ, Duong DH, Nga BTT, Ngoc NA, Linh DT, Phuong PT, et al. Incidence and prediction nomogram for metabolic syndrome in a middle-aged Vietnamese population: a 5-year follow-up study. *Endocrine.* (2022) 75:108–18. doi: 10.1007/s12020-021-02836-5
- Bakhshayeshkaram M, Heydari ST, Honarvar B, Keshani P, Roozbeh J, Dabbaghmanesh MH, et al. Incidence of metabolic syndrome and determinants of its progression in Southern Iran: a 5-year longitudinal follow-up study. *J Res Med Sci.* (2020) 25:103. doi: 10.4103/jrms.JRMS_884_19
- Sonmez B, Bozkurt B, Atmaca A, Irkec M, Orhan M, Aslan U. Effect of glycemic control on refractive changes in diabetic patients with hyperglycemia. *Cornea.* (2005) 24:531–7. doi: 10.1097/01.icc.0000151545.00489.12
- Ye L, He J, Zhang X, Xu Y, Chen Q, Yin Y, et al. The associations of lens power with age, axial length and type 2 diabetes mellitus in Chinese adults aged 50 and above. *Eye Vis.* (2020) 7:57. doi: 10.1186/s40662-020-00222-2
- Gao L, Liu W, Liang Y, Zhang F, Wang JJ, Peng Y, et al. Prevalence and characteristics of myopic retinopathy in a rural Chinese adult population: the handan eye study. *Arch Ophthalmol.* (2011) 129:1199–204. doi: 10.1001/archophthalmol.2011.230
- Krishnaiah S, Srinivas M, Khanna RC, Rao GN. Prevalence and risk factors for refractive errors in the South Indian adult population: the Andhra Pradesh Eye disease study. *Clin Ophthalmol.* (2009) 3:17. doi: 10.2147/OPTH.S4139
- Vongphanit J, Mitchell P, Wang JJ. Prevalence and progression of myopic retinopathy in an older population. *Ophthalmology.* (2002) 109:704–11. doi: 10.1016/S0161-6420(01)01024-7
- Asakuma T, Yasuda M, Ninomiya T, Noda Y, Arakawa S, Hashimoto S, et al. Prevalence and risk factors for myopic retinopathy in a Japanese population: the Hisayama Study. *Ophthalmology.* (2012) 119:1760–5. doi: 10.1016/j.ophtha.2012.02.034
- Rose KA, Morgan IG, Ip J, Kifley A, Huynh S, Smith W, et al. Outdoor activity reduces the prevalence of myopia in children. *Ophthalmology.* (2008) 115:1279–85. doi: 10.1016/j.ophtha.2007.12.019
- Deere K, Williams C, Leary S, Mattocks C, Ness A, Blair S, et al. Myopia and later physical activity in adolescence: a prospective study. *Br J Sports Med.* (2009) 43:542–4. doi: 10.1136/bjsm.2008.049288
- Myers J, Kokkinos P, Nyelin E. Physical activity, cardiorespiratory fitness, and the metabolic syndrome. *Nutrients.* (2019) 11:1652. doi: 10.3390/nu11071652
- Koenen M, Hill MA, Cohen P, Sowers JR. Obesity, adipose tissue and vascular dysfunction. *Circ Res.* (2021) 128:951–68. doi: 10.1161/CIRCRESAHA.121.318093
- Van de Voorde J, Boydens C, Pauwels B, Decaluwé K. Perivascular adipose tissue, inflammation and vascular dysfunction in obesity. *Curr*

- Vasc Pharmacol.* (2014) 12:403–11. doi: 10.2174/1570161112666140423220628
41. Barbu E, Popescu MR, Popescu AC, Balanescu SM. Inflammation as a precursor of atherothrombosis, diabetes and early vascular aging. *Int J Mol Sci.* (2022) 23:963. doi: 10.3390/ijms23020963
 42. Hartge MM, Unger T, Kintscher U. The endothelium and vascular inflammation in diabetes. *Diab Vasc Dis Res.* (2007) 4:84–8. doi: 10.3132/dvdr.2007.025
 43. Tien PT, Lin CH, Chen CS, Chang CY, Ku H, Gan D, et al. Diacerein inhibits myopia progression through lowering inflammation in retinal pigment epithelial cell. *Mediators Inflamm.* (2021) 2021:6660640. doi: 10.1155/2021/6660640
 44. Sheng C, Zhu X, Wallman J. In vitro effects of insulin and RPE on choroidal and scleral components of eye growth in chicks. *Exp Eye Res.* (2013) 116:439–48. doi: 10.1016/j.exer.2013.08.005
 45. Feldkaemper MP, Neacsu I, Schaeffel F. Insulin acts as a powerful stimulator of axial myopia in chicks. *Invest Ophthalmol Vis Sci.* (2009) 50:13–23. doi: 10.1167/iovs.08-1702
 46. Zhu X, Wallman J. Opposite effects of glucagon and insulin on compensation for spectacle lenses in chicks. *Invest Ophthalmol Vis Sci.* (2009) 50:24–36. doi: 10.1167/iovs.08-1708
 47. Xue M, Ke Y, Ren X, Zhou L, Liu J, Zhang X, et al. Proteomic analysis of aqueous humor in patients with pathologic myopia. *J Proteomics.* (2021) 234:104088. doi: 10.1016/j.jprot.2020.104088
 48. Kostova D, Spencer G, Moran AE, Cobb LK, Husain MJ, Datta BK, et al. The cost-effectiveness of hypertension management in low-income and middle-income countries: a review. *BMJ Glob Health.* (2020) 5:e002213. doi: 10.1136/bmjgh-2019-002213
 49. Ramzan S, Timmins P, Hasan SS, Babar ZU. Cost analysis of type 2 diabetes mellitus treatment in economically developed countries. *Expert Rev Pharmacoecon Outcomes Res.* (2019) 19:5–14. doi: 10.1080/14737167.2018.1513790
 50. Tremmel M, Gerdtham UG, Nilsson PM, Saha S. Economic burden of obesity: a systematic literature review. *Int J Environ Res Public Health.* (2017) 14:435. doi: 10.3390/ijerph14040435

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Chen, Zhu, Wang, Shang, He and Li. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.