



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Mirams, M;Ayodele, BA;Tatarczuch, L;Henson, FM;Pagel, CN;Mackie, EJ

Title:

Identification of novel osteochondrosis - Associated genes

Date:

2016-03-01

Citation:

Mirams, M., Ayodele, B. A., Tatarczuch, L., Henson, F. M., Pagel, C. N. & Mackie, E. J. (2016). Identification of novel osteochondrosis - Associated genes. *Journal of Orthopaedic Research*, 34 (3), pp.404-411. <https://doi.org/10.1002/jor.23033>.

Persistent Link:

<https://hdl.handle.net/11343/123470>

This is the accepted version of the following article: Mirams, M., Ayodele, B. A., Tatarczuch, L., Henson, F. M., Pagel, C. N. and Mackie, E. J. (2016), Identification of novel osteochondrosis— Associated genes. *J. Orthop. Res.*, 34: 404–411. doi:10.1002/jor.23033, which has been published in final form at <http://onlinelibrary.wiley.com/doi/10.1002/jor.23033/abstract>



Identification of novel osteochondrosis-associated genes

Journal:	<i>Journal of Orthopaedic Research</i>
Manuscript ID:	JOR-15-0234.R1
Wiley - Manuscript type:	Research Article (Non-Member)
Date Submitted by the Author:	n/a
Complete List of Authors:	Mirams, Michiko; University of Melbourne, Veterinary and Agricultural Sciences Ayodele, Babatunde; University of Melbourne, Veterinary and Agricultural Sciences Tatarczuch, Liliana; University of Melbourne, Veterinary and Agricultural Sciences Henson, Frances; University of Cambridge, Veterinary Medicine Pagel, Charles; University of Melbourne, Veterinary and Agricultural Sciences Mackie, Eleanor; University of Melbourne, Veterinary and Agricultural Sciences
Keywords:	osteochondrosis, chondrocyte, ATP6V0D2, thymosin β 4

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **Revised Manuscript JOR-15-0234**

2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

3 **IDENTIFICATION OF NOVEL OSTEOCHONDROSIS-ASSOCIATED GENES**

5 Michiko Mirams^{1#}, Babatunde A Ayodele¹, Liliana Tatarczuch¹, Frances M Henson², Charles
6 N Pagel¹, Eleanor J Mackie¹

8 ¹ Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Parkville,
9 VIC 3010, Australia

10 ² Department of Veterinary Medicine, University of Cambridge, Cambridge, UK

11 [#] Current address: Department of Microbiology and Immunology, The University of
12 Melbourne

16 Corresponding author:

17 Professor Eleanor Mackie
18 Faculty of Veterinary and Agricultural Sciences,
19 University of Melbourne, Parkville,
20 Victoria 3010, Australia.
21 Tel: +61 3 8344 7360
22 Email: ejmackie@unimelb.edu.au

25 Running title: Novel cartilage genes in osteochondrosis

27 Author contributions: MM – research design, acquisition, analysis and interpretation of data,
28 drafting manuscript; BA – acquisition and interpretation of data, drafting manuscript; LT –
29 acquisition and interpretation of data, critical revision of manuscript; FH – acquisition of
30 data, critical revision of manuscript; CP – research design, interpretation of data, critical
31 revision of manuscript; EM – research design, interpretation of data, critical revision of
32 manuscript. All authors have read and approved the final submitted manuscript.

ABSTRACT

During the early stages of articular osteochondrosis, cartilage is retained in subchondral bone, but the pathophysiology of this condition of growing humans and domestic animals is poorly understood. A subtractive hybridisation study was undertaken to compare gene expression between the cartilage of early experimentally induced equine osteochondrosis lesions and control cartilage. Of the many putative differentially expressed genes identified, eight were confirmed by quantitative PCR analysis as differentially expressed, in addition to those already known to be associated with early lesions. Genes encoding vacuolar H⁺-ATPase V₀ subunit d₂ (ATP6V0D2), cathepsin K, integrin-binding sialoprotein, integrin αV, low density lipoprotein receptor-related protein 4, lumican, osteopontin and thymosin β4 (TMSB4) were expressed at higher levels in lesions than in control cartilage. These genes included 34 genes not previously identified in cartilage. Some genes identified as associated with early lesions are known chondrocyte hypertrophy-associated genes, and in transmission electron microscopy studies normal hypertrophic chondrocytes were observed in lesions. Differential expression of ATP6V0D2 and TMSB4 in the cartilage of early naturally occurring osteochondrosis lesions was confirmed by immunohistochemistry. These results identify novel osteochondrosis-associated genes and provide evidence that articular osteochondrosis does not necessarily result from failure of chondrocytes to undergo hypertrophy.

Keywords: osteochondrosis, chondrocyte, ATP6V0D2, thymosin β4

55 INTRODUCTION

56 Most bones grow through the process of endochondral ossification, in which
57 chondrocytes play a central role. Growth cartilage is comprised of chondrocytes arranged in
58 zones that correspond to the stages of an organised program of sequential biological events.
59 A zone of resting chondrocytes blends into a zone of proliferative chondrocytes and then a
60 zone of hypertrophic chondrocytes, which ultimately undergo physiological death. Within
61 the zone of hypertrophy, the cartilage matrix surrounding individual chondrocytes is partially
62 degraded, leaving behind cartilage remnants that form vertical struts onto which bone matrix
63 is deposited by invading osteoblasts (reviewed by Mackie et al., 2011¹). The processes of
64 chondrocyte proliferation, differentiation and hypertrophy are tightly regulated by a variety
65 of growth factors, hormones, transcription factors and components of the cartilage matrix.
66 Many of these factors have been characterised, but it is likely that more remain to be
67 identified. The process of physiological death undergone by chondrocytes is less well
68 characterised and there has been a debate in the literature as to whether it occurs through
69 apoptosis or a non-apoptotic mode of death^{2, 3}. Moreover, the chondrocytes in growth
70 cartilage express factors capable of regulating the behaviour of cells in the invading
71 ossification front, including vascular endothelial cells, osteoclasts and osteoblasts¹.

72 Osteochondrosis is a developmental orthopaedic disorder in which endochondral
73 ossification is disrupted; it affects growing animals of a number of species, including
74 humans⁴. Disregulated endochondral ossification in the articular-epiphyseal growth cartilage
75 results in focal retention of cartilage in subchondral bone, which can lead to cartilage fissures
76 and synovitis. The pathophysiology of osteochondrosis is poorly understood, and possible
77 causes proposed in the literature include ischaemic necrosis of growth cartilage, trauma,
78 altered cartilage extracellular matrix composition and failure of chondrocytes in growth
79 cartilage to undergo hypertrophy⁵⁻⁹. It has been proposed that common mechanisms may
80 occur across species, and there is strong evidence for ischaemic necrosis as the aetiology of

1
2 81 osteochondrosis in pigs and horses⁴. A number of studies in horses have observed an
3
4 82 association between the presence of necrotic blood vessels in cartilage canals and
5
6 83 osteochondrosis lesions, and experimental surgical transection of blood vessels within
7
8 84 cartilage canals causes necrosis of the surrounding cartilage and an associated focal delay in
9
10 85 endochondral ossification¹⁰⁻¹². It appears, however, that there may be more than one
11
12 86 mechanism in horses since many early lesions in this species show no evidence of ischaemic
13
14 87 necrosis^{4; 10; 13-15}. The histology of early lesions in humans has not been described.

17 88 In order to learn about the tissue changes that initiate osteochondrosis, it is important
18
19 89 to study early lesions, before the secondary changes associated with clinical disease are
20
21 90 observed. A number of recent studies in horses have compared gene or protein expression in
22
23 91 the cartilage of early osteochondrosis lesions with that of normal articular cartilage, with the
24
25 92 aim of learning more about the pathophysiology of this condition. The studies undertaken so
26
27 93 far have focussed on specific molecules known to be involved in chondrocyte function. The
28
29 94 distribution of transforming growth factor- β 1 and collagen type VI in cartilage is altered in
30
31 95 early lesions^{14; 16}. Transcripts shown to be more highly expressed in early lesions compared
32
33 96 to normal cartilage include collagen types I and X (*COL1A1* and *COL10A1*), matrix
34
35 97 metalloproteinases-3 and -13 (*MMP3* and *MMP13*), Runx2 (*RUNX2*), Indian hedgehog
36
37 98 (*IHH*) and platelet-derived growth factor A (*PDGFA*)^{13; 17}. Since many of these genes are
38
39 99 up-regulated with chondrocyte hypertrophy, these observations have led to the conclusion
40
41
42 100 that failure of chondrocytes to undergo hypertrophy is unlikely to be a primary cause of
43
44 101 equine osteochondrosis^{1; 13}. The aim of the current project was to take an unbiased approach
45
46 102 (subtractive hybridization) to identification of genes selectively expressed in early equine
47
48
49 103 osteochondrosis lesions.
50
51
52

53 104

55 105

57 106

1
2 107 **MATERIALS AND METHODS**

3
4 108 **Tissue samples**

5
6 109 The use of animal tissues met the requirements of the University of Melbourne
7
8 110 Animal Ethics Committee. Articular osteochondrosis lesions were collected (in a previously
9
10 111 published study¹³) from horses (aged 6-11 months) in which osteochondrosis was induced
11
12 112 through a high energy diet fed from the age of 4-7 months, following a method that had been
13
14 113 shown to induce osteochondrosis¹⁸. Lesions were identified grossly in osteochondrosis
15
16 114 predilection sites as foci of retained cartilage in subchondral bone; none of the lesions
17
18 115 showed signs associated with clinical osteochondrosis (such as cartilage fractures or
19
20 116 synovitis). Samples of lesions were excised with some underlying bone, and similar samples
21
22 117 of normal articular cartilage were obtained from the same joints as the lesions. Samples were
23
24 118 divided into two parts. One part was used for RNA extraction from cartilage excised from
25
26 119 the specimens, as described¹³. The other part was fixed in 5% gluteraldehyde/4%
27
28 120 paraformaldehyde and embedded in Spurr's resin as described¹³, then processed for
29
30 121 transmission electron microscopy (TEM) analysis as described². Lesions and control
31
32 122 cartilage from three animals were examined. Specimens of normal articular-epiphyseal
33
34 123 growth cartilage were harvested from four equine foetuses (estimated gestational age 9-10
35
36 124 months) obtained from a local knackery, and processed for TEM as above.

37
38 125 Naturally occurring early articular osteochondrosis lesions with some adjacent normal
39
40 126 articular cartilage (7 specimens) were collected during a previous study; 6 samples were from
41
42 127 the lateral trochlear ridge of the femur of horses aged 4-12 months and 1 from the distal
43
44 128 intermediate ridge of the tibia of a 5-month-old horse¹⁴. In this study, lesions were identified
45
46 129 as for the feeding study above, and no specimens from joints showing secondary pathological
47
48 130 changes were included. Samples were fixed and embedded in paraffin as described¹⁴.

49
50 131

51
52 132

133 **Subtractive Hybridisation**

134 Subtractive hybridisation experiments were carried out using mRNA pooled from 6
135 osteochondrosis lesions (from shoulder joints and articular processes of cervical vertebrae)
136 and from normal articular cartilage collected from 4 horses. Subtractive hybridisation is a
137 labour-intensive method, which cannot feasibly be undertaken on multiple independent
138 samples per group. Samples from different animals and anatomical locations were pooled to
139 increase the chances of identification of genes regulated in all lesions (rather than genes
140 unique to an individual animal or anatomical site). All lesions were considered to be at a
141 similar stage of development since they were induced by feeding the high energy diet. For
142 each lesion, a matching specimen of normal cartilage was obtained from the same joint; we
143 had previously determined that this was the best source of control cartilage for such
144 experiments because there is considerable variability between horses in gene expression in
145 articular cartilage from a specific anatomical site, and considerable variability within
146 individual horses between anatomical sites¹³. The mRNA was purified using magnetic beads
147 coated with oligo dTs and amplified using the MessageAmp™ II aRNA Amplification Kit
148 (Life Technologies, Carlsbad, CA, USA). Subtractive hybridisation was carried out using the
149 PCR Select cDNA Subtraction Kit (Clontech Laboratories Inc., Mountain View, CA, USA).
150 Briefly, the two mRNA populations (lesion and control) were converted to cDNA and
151 digested. One was labelled then the two were hybridised with the unlabelled population in
152 excess. Hybridised sequences were removed. The non-hybridised sequences represented
153 differentially expressed sequences in the labelled population. Two rounds of hybridisation
154 were carried out to identify differentially expressed sequences in the osteochondrosis and the
155 control cDNA.

156 Cloning of subtractive hybridisation products representing candidate differentially
157 expressed genes was carried out using pBluescript II vector and XL-2 Ultracompetent E. coli
158 (Stratagene, La Jolla, CA, USA), with clones containing inserts detected by blue-white

159 colony selection. Using M13 forward and reverse sequencing primers, PCR was carried out
160 on inserts from 650 control and 1327 lesion clones. Of these amplified products, 106 from
161 control cartilage and 291 from lesions were sequenced (AGRF, Melbourne, VIC, Australia).
162 The identity of sequenced products was determined using BLAST. Of the identified genes, 30
163 were represented 2 or more times and approximately one third of the sequences from the
164 lesions aligned to *COL1A2*.

165

166 **Quantification of mRNA by quantitative PCR**

167 Oligonucleotide primers for quantitative PCR (qPCR) were designed for the gene
168 products identified in the subtractive hybridisation study that had not previously been
169 associated with osteochondrosis (Supplementary Table 1). QPCR was used to confirm
170 differential expression in osteochondrosis lesions compared to normal cartilage using RNA
171 from lesions and controls collected from 5 horses. The qPCR was performed using a
172 Stratagene MX3000 machine using Sybr Green chemistry (Life Technologies, Carlsbad, CA,
173 USA). Ribosomal protein S23 (RPS23) was used as a housekeeping gene.

174

175 **Statistical analysis**

176 Results for qPCR are presented as mean normalised expression (MNE)¹⁹ ± standard
177 deviation (SD) of genes of interest in lesions and in control cartilage samples (normalised for
178 RPS23). Results were analysed for significant differences between lesion and control
179 cartilage by one-tailed Wilcoxon matched pairs signed rank test.

180

181 **Immunohistochemistry**

182 Paraffin-embedded sections of naturally occurring osteochondrosis lesions were used
183 for detection of expression of vacuolar H⁺ ATPase V₀ subunit d₂ (ATP6V0D2) and thymosin
184 β4 (TMSB4) by immunohistochemistry. Sections were de-paraffinised and hydrated,

185

186

187

188

189

185 incubated at 95°C in Tris-EDTA solution (10 mM Tris base, 1 mM EDTA, 0.05% Tween 20,
186 pH 9.0) for 30 min for antigen retrieval, washed twice in phosphate-buffered saline
187 containing 0.05% Tween 20 (PBS/Tween), then incubated in 3% hydrogen peroxide in water
188 for 30 min. Sections were incubated for 30 min in 2.5% foetal calf serum in PBS, then in
189 rabbit anti-human ATP6V0D2 (1:50 in PBS containing 0.5% v/v Triton X100 and 1% w/v
190 bovine serum albumin; AbCam, Cambridge, UK), rabbit anti-human TMSB4 (1:1000 in
191 PBS/Tween; AbCam) or normal rabbit serum diluted appropriately. Primary antibodies were
192 detected using biotinylated goat anti-rabbit Ig (2 µg/ml in PBS/Tween, 30 min; Vector
193 Laboratories, Burlingame, CA, USA) followed by peroxidase-labelled avidin-biotin
194 complexes (ABC kit, Vector Laboratories) used according to the manufacturer's instructions,
195 then diaminobenzidine tetrahydrochloride (DakoCytomation, Glostrup, Denmark). Sections
196 were mounted with DPX. Adjacent sections were stained with haematoxylin and eosin using
197 standard methods.

198

199 RESULTS

200 Gene expression analysis

201 Subtractive hybridisation experiments were carried out to identify genes that were
202 differentially expressed in cartilage from osteochondrosis lesions compared to normal
203 cartilage. Sixty-six genes were identified as putative differentially expressed genes, either
204 more highly expressed in lesions or more highly expressed in normal cartilage
205 (Supplementary Table 2). These included four genes previously found to be differentially
206 expressed in osteochondrosis, i.e. *MMP13*, *RUNX2*, *COL1A1* and *COL10A1*, and two
207 previously found not to be differentially expressed, i.e. *COL2A1* and *VEGFA* (not listed in
208 the table)¹³. Thirty-four of the genes have not previously been described in chondrocytes
209 (Supplementary Tables 2, 3).

1
2 210 Quantitative PCR was undertaken in individual lesions isolated from horses that
3
4 211 received the high energy diet to determine whether the genes identified in the subtractive
5
6 212 hybridisation study were indeed differentially expressed. Results of qPCR experiments
7
8 213 confirmed that eight of the genes identified were significantly more highly expressed in
9
10 214 lesions compared to normal cartilage (Figure 1); these genes were vacuolar H⁺ ATPase
11
12 215 subunit V₀ d₂ (*ATP6V0D2*), cathepsin K (*CTSK*), integrin-binding sialoprotein (*IBSP*),
13
14 216 integrin αV (*ITGAV*), low density lipoprotein receptor-related protein 4 (*LRP4*), lumican
15
16
17 217 (*LUM*), osteopontin (*SPP1*) and thymosin β4 (*TMSB4*). These eight genes have not
18
19 218 previously been reported to have any involvement in osteochondrosis, and have diverse
20
21 219 functions. Some genes were identified by subtractive hybridisation as being putatively more
22
23 220 highly expressed in normal cartilage specimens than in osteochondrosis lesions, but the
24
25 221 differences were not found to be significant by qPCR. All genes that were not differentially
26
27 222 expressed between lesions and normal cartilage were however confirmed as being
28
29 223 reproducibly expressed in cartilage samples, with Ct values of less than 40.
30
31
32

224

225 **Evidence for the presence of hypertrophic chondrocytes in lesions**

36
37 226 Some of the differentially expressed genes identified here and in previous studies are
38
39 227 known to be up-regulated with chondrocyte hypertrophy^{13;17}. Moreover, we have previously
40
41 228 demonstrated in light microscopy studies that the mean area of chondrocytes in sections of
42
43 229 experimentally induced lesions is greater than that of chondrocytes in control cartilage¹³, thus
44
45 230 TEM studies of lesions were undertaken to determine whether they contained typical
46
47 231 hypertrophic chondrocytes. At the time experimentally induced lesions were harvested, the
48
49 232 horses were sufficiently old for normal articular-epiphyseal growth cartilage to have been
50
51 233 almost completely removed by the invading ossification front, thus the normal cartilage
52
53 234 specimens from these horses consisted of articular cartilage containing non-hypertrophic
54
55 235 chondrocytes (Figure 2A)¹³. The chondrocytes in the lesions showed variable morphology,
56
57
58
59
60

1
2 236 but many had morphology very similar to that of hypertrophic chondrocytes in foetal
3
4 237 articular-epiphyseal growth cartilage (Figure 2B, C). Each of these cells showed multiple
5
6 238 processes extending into the surrounding lacuna, the edge of which was at some distance
7
8 239 from the main body of the cell, in contrast to the chondrocytes in normal cartilage, which
9
10 240 filled their lacunae.

11
12
13 241

14 15 242 **Immunohistochemical analysis of ATP6V0D2 and TMSB4**

16
17 243 Two of the differentially expressed genes (*ATP6V0D2* and *TMSB4*) were selected for
18
19 244 immunohistochemical investigation of protein expression in naturally occurring
20
21 245 osteochondrosis lesions and adjacent normal cartilage; Figure 3 shows staining of a
22
23 246 representative lesion. In normal cartilage, chondrocytes showed weak specific staining for
24
25 247 *ATP6V0D2* (Figure 3B). In five of seven lesions examined, however, intense staining was
26
27 248 detected in chondrocytes in many regions within the retained cartilage (Figure 3D). In the
28
29 249 two remaining lesions, only weak staining was observed in the chondrocytes of the retained
30
31 250 cartilage. Weak to moderate immunostaining for *TMSB4* was detected in normal cartilage;
32
33 251 staining was observed both in chondrocytes and in extracellular matrix (Figure 3C). Staining
34
35 252 of chondrocytes was generally stronger in lesions than in normal cartilage in all seven of the
36
37 253 specimens examined with this antibody; the strongest staining in lesions was observed in
38
39 254 chondrocytes arranged in clusters (Figure 3E). The profiles of cell-associated staining for
40
41 255 *TMSB4* in lesions were larger than those in normal cartilage, and were suggestive of staining
42
43 256 of pericellular matrix within large lacunae (Figure 3C, E); this was in contrast to staining of
44
45 257 *ATP6V0D2*, which appeared to be restricted to the cytoplasm (Figure 3D).

46
47
48 258

49
50
51 259

52
53
54 260

55
56
57 261

58
59
60

262 **DISCUSSION**

263 One approach to investigating the pathophysiology of osteochondrosis is to study
264 differences in gene expression between lesions and normal tissue. A number of published
265 studies taking this approach to investigate the disease in horses have used samples from
266 lesions of a sufficiently advanced stage to be identifiable clinically, often collected at the time
267 of surgery to correct the defect^{6; 20; 21}. In such studies, many of the differences in gene
268 expression are likely to be part of a secondary inflammatory response, rather than associated
269 with the primary change that leads to cartilage retention. In the current study in horses, we
270 have used an experimental model of osteochondrosis in order to obtain lesions at an early
271 stage. Previous studies in this model as well as in early naturally occurring equine lesions
272 have investigated the association of known cartilage genes and/or proteins with
273 osteochondrosis^{13; 14; 16; 17}. In contrast, in the current study, we have taken an unbiased
274 approach to identifying genes that are differentially expressed in early lesions; the subtractive
275 hybridisation method has allowed for the possibility of detecting osteochondrosis-associated
276 genes that were not previously known to be expressed in cartilage. Of the 66 putative
277 differentially expressed genes, four had previously been shown to be more highly expressed
278 in early lesions than in normal cartilage¹³. Since the subtractive hybridisation was
279 undertaken with specimens pooled either from lesions or normal cartilage, qPCR studies were
280 undertaken to determine whether putative differences were statistically significant. Through
281 this process, eight genes not previously identified as early osteochondrosis-associated genes
282 were confirmed as being more highly expressed in lesions than in normal cartilage, including
283 one gene that had not previously been identified in cartilage, i.e. *ATP6V0D2*. The
284 differentially expressed genes cover a variety of functions, but the majority encode
285 extracellular matrix proteins or proteins with known or putative functions in extracellular
286 matrix turnover or interactions with cells.

1
2 287 Three of the genes found to be more highly expressed in the osteochondrosis lesions
3
4 288 than in controls encode extracellular matrix proteins in bone and/or cartilage. Lumican is a
5
6 289 small proteoglycan known to be expressed in articular cartilage; the level of expression in
7
8 290 this location in humans is higher in adults than in infants²². IBSP and osteopontin (encoded
9
10 291 by *SPPI*) are both important components of bone extracellular matrix, but they are also both
11
12 292 expressed selectively by hypertrophic chondrocytes in the growth plate²³. Elevated
13
14 293 expression of these genes is in agreement with previous observations of selective expression
15
16 294 in osteochondrosis lesions of hypertrophy-associated genes including *MMP13* and *COL10A1*
17
18 295^{13; 17}. The TEM studies presented here demonstrate that the morphology of many
19
20 296 chondrocytes in lesions is similar to that of hypertrophic chondrocytes in articular-epiphyseal
21
22 297 growth cartilage rather than chondrocytes in normal cartilage from the same joints as the
23
24 298 lesions. We and others have previously hypothesised that osteochondrosis results from a
25
26 299 failure of chondrocytes in growth cartilage to undergo hypertrophy^{5-8; 15}, but the gene
27
28 300 expression and TEM observations presented here indicate that generalised suppression of
29
30 301 hypertrophy cannot account for the presence of the lesions.

31
32
33
34
35 302 TMSB4 is an actin sequestering protein with functions relating to cell migration,
36
37 303 angiogenesis, tumour cell invasion and wound healing²⁴. Levels of its transcript were higher
38
39 304 in osteochondrosis lesions than in normal cartilage, and immunohistochemical staining was
40
41 305 more intense in chondrocytes in lesions than in normal cartilage. While TMSB4 has a clear
42
43 306 intracellular role in actin binding, many cellular responses are observed when it is applied
44
45 307 extracellularly. TMSB4 expression has been described in cultured articular cartilage explants
46
47 308 and shown to be up-regulated in response to mechanical loading; treatment of primary
48
49 309 chondrocyte cultures with TMSB4 stimulated expression and activation of MMP-9²⁵. The
50
51 310 elevated expression in osteochondrosis lesions may be related to altered cytoskeletal
52
53 311 organisation in hypertrophic chondrocytes, and such changes may in turn influence
54
55 312 expression of other genes such as MMPs. It is interesting to note, however, that in the
56
57
58
59
60

1
2 313 current study specific staining for TMSB4 was observed in the cartilage extracellular matrix
3
4 314 (as well as in the chondrocyte cytoplasm), suggesting that it may play roles in cartilage
5
6 315 independently of its ability to bind actin.
7

8 316 Integrin αV , a subunit of a number of receptors for extracellular matrix proteins, is
9
10 317 known to be expressed by chondrocytes²⁶ and mediates mechanical loading-induced
11
12 318 proteoglycan synthesis by chondrocytes²⁷. Expression of integrin αV is greater in
13
14 319 chondrocytes from osteoarthritic cartilage compared to those from normal cartilage²⁸. The
15
16 320 higher level of expression of integrin αV in osteochondrosis lesions is likely to mediate
17
18 321 altered interactions between chondrocytes and their surrounding matrix, and may influence
19
20 322 cytoskeletal organisation and hypertrophy.
21
22
23

24 323 Cathepsin K is expressed by osteoclasts and assists in osteoclastic bone resorption
25
26 324 through its proteolytic activity²⁹. Selective expression of cathepsin K has previously been
27
28 325 detected in hypertrophic chondrocytes in mouse growth plate³⁰. The elevated expression of
29
30 326 this hypertrophy-associated matrix-degrading protease is in keeping with our earlier finding
31
32 327 of elevated *MMP13* expression in osteochondrosis lesions.
33
34

35 328 LRP4 is a member of the LRP family of receptors; in osteoblasts, LRP4 interacts with
36
37 329 the Wnt antagonist sclerostin to enhance its inhibitory activity³¹. It has recently been
38
39 330 demonstrated that *Lrp4* expression by mouse chondrocytes is up-regulated with hypertrophy,
40
41 331 and that *Lrp4* overexpression stimulates expression of extracellular matrix genes in cultured
42
43 332 chondrocytes³². Sclerostin is also expressed by hypertrophic chondrocytes and inhibits IL-
44
45 333 1α -stimulated aggrecan degradation in cartilage explants^{33; 34}. It is possible that the enhanced
46
47 334 expression of *LRP4* in osteochondrosis lesions (observed in the current study) contributes to
48
49 335 the retention of cartilage in subchondral bone both through increased expression of cartilage
50
51 336 matrix and inhibition of matrix degradation. Such roles for *LRP4* may help to explain how
52
53 337 cartilage is retained in subchondral bone despite the concurrently elevated levels of matrix-
54
55 338 degrading enzymes such as cathepsin K and MMP-13.
56
57
58
59
60

1
2 339 ATP6V0D2, like cathepsin K, is expressed by osteoclasts; it is required for osteoclast
3
4 340 differentiation as well as the bone-resorptive function of mature osteoclasts³⁵. This protein is
5
6 341 a subunit of the vacuolar H⁺ ATPase, which is responsible for acidification of the
7
8 342 microenvironment between osteoclasts and the bone surface. Evidence for the presence of
9
10 343 this enzyme in chondrocytes has been described³⁶. The vacuolar H⁺ ATPase is comprised of
11
12 344 at least 14 subunits, of which several exist as multiple isoforms expressed differentially
13
14 345 between tissues³⁵. The ATP6V0D2 subunit has not previously been identified in cartilage. It
15
16 346 seemed possible that *ATP6V0D2* may have been detected in the hybridisation and PCR
17
18 347 studies due to the presence of osteoclasts in cartilage canals in the lesions, however
19
20 348 immunohistochemistry was used to confirm that it is expressed by chondrocytes. Moreover,
21
22 349 the immunohistochemical results demonstrated that up-regulation of ATP6V0D2 occurred in
23
24 350 most of the naturally occurring osteochondrosis lesions examined (not just the experimentally
25
26 351 induced lesions used for the subtractive hybridisation and qPCR studies). Vacuolar H⁺
27
28 352 ATPase activity is required for interleukin-1-stimulated proteoglycan degradation by
29
30 353 chondrocytes³⁶, thus *ATP6V0D2* is another example of a gene that is highly expressed in
31
32 354 osteochondrosis lesions that might be expected to contribute to matrix degradation.

33
34
35
36
37 355 Other genes that were identified by subtractive hybridisation but were not confirmed
38
39 356 as being differentially expressed in osteochondrosis lesions by qPCR included 34 genes that
40
41 357 have not been described as being expressed in cartilage (Supplementary Table 3). Many of
42
43 358 these are genes encoding proteins with known functions in transcription as well as protein
44
45 359 translation and secretion, which is in keeping with the high secretory activity of
46
47 360 chondrocytes. Further investigations will be needed to clarify specific roles of the novel
48
49 361 genes in regulation of chondrocyte behaviour and/or cartilage function.

50
51
52 362 A limitation of the study was that insufficient material was available from some of the
53
54 363 horses used in the experimental osteochondrosis study (on which the current study was
55
56 364 based) to undertake qPCR for all of the 66 putative differentially expressed genes. It is
57
58
59
60

1
2 365 possible that additional genes may have proved to be differentially expressed if more
3
4 366 specimens had been available.

5
6 367 A recent study by Olstad *et al*¹⁰ describing a detailed examination of early
7
8 368 osteochondrosis lesions of the distal intermediate ridge of the equine tibia recommended that
9
10 369 future studies on the pathogenesis of osteochondrosis at that site should be ‘focused on foals
11
12 370 younger than 4.5 months old’. This recommendation was based on the conclusion that the
13
14 371 lesions are caused by necrosis of cartilage canal blood vessels, and the fact that such vessels
15
16 372 are absent after 4.5 months. As noted in the introduction, while there is strong evidence that
17
18 373 many equine osteochondrosis lesions are caused by necrosis related to failure of cartilage
19
20 374 canal blood vessels, in some studies lesions have been described in which no such association
21
22 375 has been observed. Since there is a possibility that some lesions are caused by a different
23
24 376 mechanism, it is appropriate to investigate lesions from sites related to growth plates that are
25
26 377 still active, with the proviso that no gross signs of secondary changes are present. This is the
27
28 378 approach taken for the specimens used in the immunohistochemical studies presented here;
29
30 379 the specimens (distal ends of femur and tibia) were taken several months before the expected
31
32 380 closure of the associated growth plates³⁷. The specimens from the feeding trial were from
33
34 381 sites (glenoid of the scapula and cervical vertebrae) for which specific growth plate closure
35
36 382 data are not available, however these bones continue to grow significantly in the second year
37
38 383 of life³⁸. Horses reach full skeletal maturity in the third year of life.

39
40
41
42
43 384 The results presented here provide further evidence that the fundamental defect
44
45 385 leading to cartilage retention in subchondral bone at the initiation of equine osteochondrosis
46
47 386 is not failure of chondrocytes to undergo hypertrophy or to express matrix-degrading
48
49 387 enzymes. Many chondrocytes in lesions are morphologically indistinguishable from
50
51 388 hypertrophic chondrocytes in normal articular-epiphyseal growth cartilage, and of all the
52
53 389 genes shown here to be more highly expressed in lesions, four are known hypertrophy-
54
55 390 associated genes (*CTSK*, *IBSP*, *LRP4* and *SSP1*). However it is possible that these cells
56
57
58
59
60

1
2 391 exhibit a different gene expression profile from normal hypertrophic chondrocytes and that
3
4 392 they or other chondrocytes in the lesions express a combination of genes that outweigh the
5
6 393 effects of those expected to lead to replacement of cartilage by bone. Indeed, two of the
7
8 394 remaining genes are potentially involved in synthesis or maintenance of cartilage
9
10 395 extracellular matrix, including a matrix protein (*LUM*) and a putative mediator of matrix
11
12 396 synthesis (*ITGAV*). Future studies on the roles of these genes in cartilage may shed light on
13
14
15 397 the events associated with the initiation of osteochondrosis.
16
17
18
19
20
21
22

23 400 **ACKNOWLEDGMENTS**

24 401 This work was supported by the Australian Research Council and Racing Victoria Pty
25
26 402 Ltd (Linkage Grant no. LP0348867), and the Rural Industries Research and Development
27
28 403 Corporation (Grant no. PRJ-000216). B. Ayodele's contribution was supported by a Dr Sue
29
30 404 Newton Travelling Scholarship awarded by the Faculty of Veterinary and Agricultural
31
32
33 405 Sciences.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

408 REFERENCES

- 409 1. Mackie EJ, Tatarczuch L, Mirams M. 2011. The skeleton: a multi-functional
410 complex organ: the growth plate chondrocyte and endochondral ossification. *The*
411 *Journal of endocrinology* 211:109-121.
- 412 2. Ahmed YA, Tatarczuch L, Pagel CN, et al. 2007. Physiological death of
413 hypertrophic chondrocytes. *Osteoarthritis and cartilage / OARS, Osteoarthritis*
414 *Research Society* 15:575-586.
- 415 3. Roach HI, Clarke NM. 2000. Physiological cell death of chondrocytes in vivo is not
416 confined to apoptosis. New observations on the mammalian growth plate. *The*
417 *Journal of bone and joint surgery British volume* 82:601-613.
- 418 4. McCoy AM, Toth F, Dolvik NI, et al. 2013. Articular osteochondrosis: a
419 comparison of naturally-occurring human and animal disease. *Osteoarthritis and*
420 *cartilage / OARS, Osteoarthritis Research Society* 21:1638-1647.
- 421 5. Jeffcott LB, Henson FM. 1998. Studies on growth cartilage in the horse and their
422 application to aetiopathogenesis of dyschondroplasia (osteochondrosis).
423 *Veterinary journal* 156:177-192.
- 424 6. Semevolos SA, Strassheim ML, Haupt JL, et al. 2005. Expression patterns of
425 hedgehog signaling peptides in naturally acquired equine osteochondrosis.
426 *Journal of orthopaedic research : official publication of the Orthopaedic Research*
427 *Society* 23:1152-1159.
- 428 7. Ekman S, Heinegard D. 1992. Immunohistochemical localization of matrix
429 proteins in the femoral joint cartilage of growing commercial pigs. *Veterinary*
430 *pathology* 29:514-520.
- 431 8. Farnum CE, Wilsman NJ. 1986. Ultrastructural histochemical evaluation of
432 growth plate cartilage matrix from healthy and osteochondritic swine. *American*
433 *journal of veterinary research* 47:1105-1115.
- 434 9. Laverty S, Girard C. 2013. Pathogenesis of epiphyseal osteochondrosis.
435 *Veterinary journal* 197:3-12.
- 436 10. Olstad K, Ytrehus B, Ekman S, et al. 2007. Early lesions of osteochondrosis in the
437 distal tibia of foals. *Journal of orthopaedic research : official publication of the*
438 *Orthopaedic Research Society* 25:1094-1105.
- 439 11. Carlson CS, Cullins LD, Meuten DJ. 1995. Osteochondrosis of the articular-
440 epiphyseal cartilage complex in young horses: evidence for a defect in cartilage
441 canal blood supply. *Veterinary pathology* 32:641-647.
- 442 12. Olstad K, Hendrickson EH, Carlson CS, et al. 2013. Transection of vessels in
443 epiphyseal cartilage canals leads to osteochondrosis and osteochondrosis
444 dissecans in the femoro-patellar joint of foals; a potential model of juvenile
445 osteochondritis dissecans. *Osteoarthritis and cartilage / OARS, Osteoarthritis*
446 *Research Society* 21:730-738.
- 447 13. Mirams M, Tatarczuch L, Ahmed YA, et al. 2009. Altered gene expression in early
448 osteochondrosis lesions. *Journal of orthopaedic research : official publication of*
449 *the Orthopaedic Research Society* 27:452-457.
- 450 14. Henson FM, Davies ME, Jeffcott LB. 1997. Equine dyschondroplasia
451 (osteochondrosis)--histological findings and type VI collagen localization.
452 *Veterinary journal* 154:53-62.
- 453 15. Shingleton WD, Mackie EJ, Cawston TE, et al. 1997. Cartilage canals in equine
454 articular/epiphyseal growth cartilage and a possible association with
455 dyschondroplasia. *Equine veterinary journal* 29:360-364.

- 1
2 456 16. Henson FM, Schofield PN, Jeffcott LB. 1997. Expression of transforming growth
3 457 factor-beta 1 in normal and dyschondroplastic articular growth cartilage of the
4 458 young horse. *Equine veterinary journal* 29:434-439.
- 5 459 17. Riddick TL, Duesterdieck-Zellmer K, Semevolos SA. 2012. Gene and protein
6 460 expression of cartilage canal and osteochondral junction chondrocytes and full-
7 461 thickness cartilage in early equine osteochondrosis. *Veterinary journal* 194:319-
8 462 325.
- 9 463 18. Savage CJ, McCarthy RN, Jeffcott LB. 1993. Effects of dietary energy and protein
10 464 on induction of dyschondroplasia in foals. *Equine Vet J Suppl* 16:74-79.
- 11 465 19. Simon P. 2003. Q-Gene: processing quantitative real-time RT-PCR data.
12 466 *Bioinformatics* 19:1439-1440.
- 13 467 20. Semevolos SA, Nixon AJ, Brower-Toland BD. 2001. Changes in molecular
14 468 expression of aggrecan and collagen types I, II, and X, insulin-like growth factor-I,
15 469 and transforming growth factor-beta1 in articular cartilage obtained from horses
16 470 with naturally acquired osteochondrosis. *American journal of veterinary*
17 471 *research* 62:1088-1094.
- 18 472 21. Garvican ER, Vaughan-Thomas A, Redmond C, et al. 2008. Chondrocytes
19 473 harvested from osteochondritis dissecans cartilage are able to undergo limited in
20 474 vitro chondrogenesis despite having perturbations of cell phenotype in vivo.
21 475 *Journal of orthopaedic research : official publication of the Orthopaedic Research*
22 476 *Society* 26:1133-1140.
- 23 477 22. Grover J, Chen XN, Korenberg JR, et al. 1995. The human lumican gene.
24 478 Organization, chromosomal location, and expression in articular cartilage. *The*
25 479 *Journal of biological chemistry* 270:21942-21949.
- 26 480 23. Komori T. 2010. Regulation of bone development and extracellular matrix
27 481 protein genes by RUNX2. *Cell and tissue research* 339:189-195.
- 28 482 24. Sribenja S, Wongkham S, Wongkham C, et al. 2013. Roles and mechanisms of
29 483 beta-thymosins in cell migration and cancer metastasis: an update. *Cancer*
30 484 *investigation* 31:103-110.
- 31 485 25. Blain EJ, Mason DJ, Duance VC. 2002. The effect of thymosin beta4 on articular
32 486 cartilage chondrocyte matrix metalloproteinase expression. *Biochemical Society*
33 487 *transactions* 30:879-882.
- 34 488 26. Hausler G, Helmreich M, Marlovits S, et al. 2002. Integrins and extracellular
35 489 matrix proteins in the human childhood and adolescent growth plate. *Calcified*
36 490 *tissue international* 71:212-218.
- 37 491 27. Chai DH, Arner EC, Griggs DW, et al. 2010. Alphas and beta1 integrins regulate
38 492 dynamic compression-induced proteoglycan synthesis in 3D gel culture by
39 493 distinct complementary pathways. *Osteoarthritis and cartilage / OARS,*
40 494 *Osteoarthritis Research Society* 18:249-256.
- 41 495 28. Kostopoulou F, Gkretsi V, Malizos KN, et al. 2012. Central role of SREBP-2 in the
42 496 pathogenesis of osteoarthritis. *PloS one* 7:e35753.
- 43 497 29. Novinec M, Lenarcic B. 2013. Cathepsin K: a unique collagenolytic cysteine
44 498 peptidase. *Biological chemistry* 394:1163-1179.
- 45 499 30. Rantakokko J, Aro HT, Savontaus M, et al. 1996. Mouse cathepsin K: cDNA cloning
46 500 and predominant expression of the gene in osteoclasts, and in some
47 501 hypertrophying chondrocytes during mouse development. *FEBS letters* 393:307-
48 502 313.
- 49 503 31. Leupin O, Piters E, Halleux C, et al. 2011. Bone overgrowth-associated mutations
50 504 in the LRP4 gene impair sclerostin facilitator function. *The Journal of biological*
51 505 *chemistry* 286:19489-19500.

- 1
2 506 32. Asai N, Ohkawara B, Ito M, et al. 2014. LRP4 induces extracellular matrix
3 507 productions and facilitates chondrocyte differentiation. *Biochemical and*
4 508 *biophysical research communications* 451:302-307.
5 509 33. van Bezooijen RL, Bronckers AL, Gortzak RA, et al. 2009. Sclerostin in
6 510 mineralized matrices and van Buchem disease. *Journal of dental research*
7 511 88:569-574.
8 512 34. Chan BY, Fuller ES, Russell AK, et al. 2011. Increased chondrocyte sclerostin may
9 513 protect against cartilage degradation in osteoarthritis. *Osteoarthritis and*
10 514 *cartilage / OARS, Osteoarthritis Research Society* 19:874-885.
11 515 35. Qin A, Cheng TS, Pavlos NJ, et al. 2012. V-ATPases in osteoclasts: structure,
12 516 function and potential inhibitors of bone resorption. *The international journal of*
13 517 *biochemistry & cell biology* 44:1422-1435.
14 518 36. Yocum SA, Lopresti-Morrow LL, Gabel CA, et al. 1995. Bafilomycin A1 inhibits IL-
15 519 1-stimulated proteoglycan degradation by chondrocytes without affecting
16 520 stromelysin synthesis. *Archives of biochemistry and biophysics* 316:827-835.
17 521 37. Strand E, Braathen LC, Hellsten MC, et al. 2007. Radiographic closure time of
18 522 appendicular growth plates in the Icelandic horse. *Acta Vet Scand* 49:19.
19 523 38. Anderson TM, McIlwraith CW. 2004. Longitudinal development of equine
20 524 conformation from weanling to age 3 years in the Thoroughbred. *Equine*
21 525 *veterinary journal* 36:563-570.
22 526
23
24
25
26 527
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

528 **FIGURE LEGENDS**

529 **Figure 1: Genes differentially expressed in osteochondrosis lesions.**

530 QPCR results showing MNE (\pm SD) of genes in osteochondrosis samples compared with
531 control samples. Significant differences between lesions and control cartilage are indicated
532 as $*= p<0.05$; $n = 5$ horses.

533
534 **Figure 2: Morphology of chondrocytes in equine osteochondrosis lesions.**

535 TEM images of representative chondrocytes from: the deep zone of normal articular cartilage
536 (A), an osteochondrosis lesion from the same shoulder joint as the cartilage shown in A (B),
537 and normal foetal articular-epiphyseal growth cartilage from the head of the humerus of a 9-
538 month equine foetus (C). A light microscopy image of the specimen shown in C is presented
539 in Figure S-1. The images presented in A and B are all from a single animal, but are
540 representative of cells observed in specimens from 3 animals. Arrows indicate the border of
541 the chondrocyte lacuna; n – nucleus. All images have the same magnification; bar = 5 μ m.

542
543 **Figure 3: ATP6V0D2 and TMSB4 expression in osteochondrosis lesions.**

544 Immunohistochemistry was carried out on paraffin sections of equine osteochondrosis lesions
545 with adjacent normal articular cartilage. A: Haematoxylin and eosin staining of a specimen
546 taken from the lateral trochlear ridge of the distal end of the femur of a six month old foal,
547 indicating the regions shown in images B-G. Normal articular cartilage overlying
548 subchondral bone can be seen to the left of the image, and the lesion can be seen to the right,
549 with cartilage retained in subchondral bone (arrows indicate bone-cartilage interface).
550 Normal region (B, C, F, G) and lesion (D, E) stained with anti-ATP6V0D2 (B, D), anti-
551 TMSB4 (C, E) or normal rabbit serum (NRS; F, G). Arrowheads (inset in D, E) indicate the
552 border of the chondrocyte lacuna. Bar in A = 500 μ m. The magnification for parts B – G is
553 the same; bar = 200 μ m for main image and 25 μ m for insets.

554 **SUPPLEMENTARY TABLE 1: Primer sequences.** Forward and reverse sequences of
 555 primers used for qPCR.
 556

Gene	Forward	Reverse
<i>ANKRA2</i>	GCTGCTGACTGTGGAGTTG	TTGGGTCAGCTCCATTTTCT
<i>ANXA1</i>	TAAGGGTGACCGATCTGAGG	TTCTCCTTTCTCCTGCTTCG
<i>ATP6V0D2</i>	AGAGTGGCTGAGCACTACGG	CACATCCTCCAAGGTCTTTCC
<i>BAK1</i>	GGCCCTGCATGTCTACCAG	GCCATTTCCTCAAGTCCAGAG
<i>BCL2</i>	TGGATGACCGAGTACCTGAAC	CCTTCAGAGACAGCCAGGAG
<i>CALM1</i>	GCACCATTGACTTCCCAGAA	GTAGTTCTGCCGCACTGATG
<i>CDK11A</i>	CATACAACAACCTCCGCAAA	CTTGAGACCGTCTCTGCAT
<i>CHTOP</i>	TCTGTCCAGGCAGCATTAAA	GGGCAAGCCTCTCTGGATTA
<i>CLIC1</i>	ACTCTGGCTGACTGCAACCT	GAACCTTCCCAGCATAGG
<i>CTSK</i>	AAGCCTGACCTCCTTCCAGT	CCGCATGGTTCAGATTATCA
<i>CYTB</i>	CCGAAAATCTCACCCACTAA	GAGGGAGCCGAAGTTTCATC
<i>DDX5</i>	CGACCTTATCTCTGTGCTTCG	AATATCTGTCCCAGCGGTCA
<i>EEF1G</i>	AACGAGTTACGCTGGCTGAC	GCTGGTTAATGCAGGTGAGG
<i>FBN1</i>	AGGAAACGGAGAAGCACAAA	AACATGGCGGTCTTCTCAAC
<i>FOXA3</i>	CTGGCCGAGTGGAGCTACTA	CAGAGGGTTCAGGGTTCATGT
<i>FTL</i>	ATCAAGAAGATGGGCGACAA	CTTGAGGGTGTAGCCTTTCAA
<i>GJA1</i>	GTCACCTGGCGAGCGTACTT	GCAGGATTCGGAAAATGAAA
<i>GNB1</i>	GGGATAACCTCCGTGTCCTT	AGGCAGCTGACACGGTTG
<i>GOLF3</i>	GCCTCCAGAGACTGTCCAGA	CCCTTTTCCACCAGGTTTTT
<i>HIVEP2</i>	TTGTCCATGGGGCAGTATTT	TGGAGCCTGGATGTTTCATAA
<i>HSP70</i>	ATCGAGCGCATGGTTCTG	ATCTTGCCCTTCAAGCCTTC
<i>IBSP</i>	GCCACAATATTCTCTTTACAAGCA	CCATCTCCATTTTCTTCAGAGG
<i>ITGAV</i>	GCAGTTCTAGCAGGGTTGTTG	CCATTTTCATGAGGTTGAAGC
<i>KCTD10</i>	GAAGCAACAACAATACTCATATACCA	TGAAGCGCAGAGACAGCTTA
<i>KDELR2</i>	TCACAACCTCGTTACCTGGATCTT	ACACTGTGGCATAGGAGCAG
<i>LDHA</i>	AATCTTAGGCGAGTGCATCC	ACGGCACACTAAGGAAGACG
<i>LECT1</i>	CAGAGCATCTCCTCCGAACT	CAAGAAGCTGTTGTCTTCCACA
<i>LRP4</i>	ATTGAAGCAATCCCAACACC	AGGTCATCCCCTCAGCATC
<i>LUM</i>	CCTCTCTGCTTGAGCTGGAT	GAGGTCCCAGGATCTTACAGAA
<i>MFAP2</i>	TTCCAGTTCAGTCTCAGCA	CTCCGTCTCTGCATTTCTCTG
<i>MFN1</i>	CAACACAATGCAAAGCTCCT	CACTGCTTGAATGTAGAACTGC
<i>MRRF</i>	ACGGAAGGTTTCGTACCAATG	ACCTGTCCAGCTCTGCAACT
<i>MT-ND2</i>	TCCCACACATGAAAACAAAACA	GGGAGGATGATGCTGCTATT
<i>NACA</i>	GCAGCTGCTGAGAAATTCAA	ACTTCCACACCCGTTTCATC
<i>NHLRC3</i>	TCATGGTATATTTGCAGCCAGT	TGGGGTACCCAAGACTTGAA
<i>NUDT16</i>	GGACCCAGGGCTGCTCTT	GTCTGCCTGTTACCAA
<i>OGN</i>	CACCATTAACCCCAAGAAA	GGACAGCATCAATGTCCACTT
<i>PIP4K2A</i>	GCCAAAGAACTGCCAACTTT	TGAGCTTTAACTGGGCAAGAA
<i>PLOD2</i>	GCACCCGTTACTACTGAAGGT	CGTAGAAGCGTCTGTGATGAG
<i>PP1CB</i>	TGGCGAGTTTGATAATGCTG	TTAGCTGTCCGAGGTGGAGT
<i>PSAP</i>	GTGGAATCCTGGTGGAGGT	AAAGAGGCTTATGGGCTGTG
<i>PTAR1</i>	GGCATATTTTCTACCTTCAGCA	GGGAATAGCCTTGCTTGCTA
<i>PURB</i>	AcGTGGGgTGCAACAAGTat	ATCCGCATACCGaCAAAAaG

<i>RAP1B</i>	GTCGTGTCGGCTGCTTTA	GGTTTAACAAAGTTGGAATGCTG
<i>RERE</i>	CCGACTGCAGATGTTCAATG	GTCAACCAGCGGGTGAAC
<i>RNF130</i>	CCTGCCCTATGTGCAAACCTT	TCGGTAACTGCTTGGGTTT
<i>RPN2</i>	CAGGGCAGTGAATGTATTGG	GATGCTCCATCGACTGTCT
<i>RPS19</i>	CTGGTTCTACACGCGAGCTG	TTCTCTGACGTCCCCATAG
<i>RPS7</i>	CACCTGGATAAAGCACAGCA	ATTAACATCCTTGCCCGTGA
<i>SERPINA1</i>	CAGACCTCTCCGGGATCACT	GTGCCTTCTCGTCAATGGT
<i>SF3B1</i>	TATGGCCCAATGTGTTTGGAG	GGGCTGGGTGAAACAAAC
<i>SPP1</i>	AATCCAGGAAGTTTCGCAGA	ATGTCCTTGCTTCCACAGG
<i>SRSF3</i>	AAGGAGTCCTCCACCTCGAC	AGGATCGAGACGGCTTATGA
<i>SSR4</i>	GCAGGCACCTATGAGGTCAG	AGGGGCGGGATGATAGAAAT
<i>SSRP1</i>	TTGGGATGAACTGGAGGAAG	CTCTTCCGGCTCATACTCG
<i>TMSB4</i>	ACAAACCCGATATGGCTGAG	CCTGCTTGCTTCTCCTGTTC
<i>TPI1</i>	TGGCTTAAGTCCAACGTCTC	AGAGAAGCACCACCCACAAG
<i>TPM2</i>	CAAACCTGCTGGAGGAGAAGC	CTTCCAGGTCATCGATGGTT
<i>WSB2</i>	CCAATTGCGTTTGCTCCTAT	TCGGCATAAGTGCTTCAGTG
<i>WTAP</i>	CCCAGCGATCAACTTGTTTT	CATTCGACACTTCGCCATTA

557

Peer Review

558 **SUPPLEMENTARY TABLE 2: Putative differentially expressed genes identified by**
 559 **subtractive hybridisation of osteochondrosis lesions and normal cartilage.** Articular
 560 osteochondrosis lesions were obtained from shoulder joints and articular processes of cervical
 561 vertebrae.

562

563

Protein and gene	qPCR ¹	Cartilage ²	Protein and gene	qPCR ¹	Cartilage ²
Ankyrin repeat, family a, 2 (<i>ANKRA2</i>)	No	No	Mitofusin 1 (<i>MFN1</i>)	No	No
Annexin A1 (<i>ANXA1</i>)	No	Yes	NADH dehydrogenase subunit 2, mitochondrial (<i>MT-ND2</i>)	No	No
Arginine-glutamic acid dipeptide repeats (<i>RERE</i>)	No	Yes	Nascent polypeptide-associated complex alpha subunit (<i>NACA</i>)	No	No
B cell CLL/lymphoma 2 (<i>BCL2</i>)	No	Yes	Non-Hodgkins Lymphoma repeat containing 3 (<i>NHLRC3</i>)	No	No
BCL2-antagonist/killer 1 (<i>BAK1</i>)	No	Yes	Nudix (nucleoside diphosphate linked moiety X)-type motif 16 (<i>NUDT16</i>)	No	No
Calmodulin 1 (<i>CALM1</i>)	No	Yes	Osteoglycin (<i>OGN</i>)	No	Yes
Cathepsin K (<i>CTSK</i>)	Yes	Yes	Osteopontin (<i>SPPI</i>)	Yes	Yes
Chloride intracellular channel protein 1 (<i>CLIC1</i>)	No	No	Phosphatidylinositol-4-phosphate 5-kinase type II alpha (<i>PIP4K2A</i>)	No	No
Chondromodulin I (<i>LECT1</i>)	No	Yes	Potassium channel tetramerisation domain containing 10 (<i>KCTD10</i>)	No	No
Chromatin target of PRMT (<i>CHTOP</i>)	No	No	Procollagen-lysine, 2-oxoglutarate 5-dioxygenase 2 (<i>PLOD2</i>)	No	Yes
Cyclin-dependent kinase 11A (<i>CDK11A</i>)	No	No	Prosaposin (<i>PSAP</i>)	No	No
Cytochrome b, mitochondrial (<i>CYTB</i>)	No	Yes	Protein phosphatase 1, catalytic subunit, beta isoform (<i>PP1CB</i>)	No	Yes
DEAD (Asp-Glu-Ala-Asp) box helicase 5 (<i>DDX5</i>)	No	No	Protein prenyl transferase alpha subunit-containing 1 (<i>PTARI</i>)	No	No
Eukaryotic translation elongation factor 1 gamma (<i>EEF1G</i>)	No	No	Purine-rich element binding protein B (<i>PURB</i>)	No	No
Ferritin, light polypeptide (<i>FTL</i>)	No	Yes	RAS oncogene family member (<i>RAP1B</i>)	No	No
Fibrillin 1 (<i>FBN1</i>)	No	Yes	Ribophorin II (<i>RPN2</i>)	No	No
Forkhead box A3 (<i>FOXA3</i>)	No	Yes	Ribosomal protein S7 (<i>RPS7</i>)	No	No
Gap junction protein, alpha 1 (<i>GJA1</i>)	No	Yes	Ribosomal protein S19 (<i>RPS19</i>)	No	No
Golgi phosphoprotein 3 (<i>GOLPH3</i>)	No	No	Ring finger protein 130 (<i>RNF130</i>)	No	No
Guanine nucleotide binding protein beta polypeptide 1 (<i>GNB1</i>)	No	No	Serpin peptidase inhibitor, clade A member 1 (<i>SERPINA1</i>)	No	Yes
Heat shock protein 70 (<i>HSP70</i>)	No	Yes	Signal sequence receptor, delta (<i>SSR4</i>)	No	No
Human immunodeficiency virus type I enhancer binding protein 2 (<i>HIVEP2</i>)	No	No	Splicing factor 3b, subunit 1 (<i>SF3B1</i>)	No	No
Integrin alpha V (<i>ITGAV</i>)	Yes	Yes	Splicing factor, arginine/serine-rich 3 (<i>SRSF3</i>)	No	No
Integrin-binding sialoprotein, bone sialoprotein (<i>IBSP</i>)	Yes	Yes	Structure specific recognition protein 1 (<i>SSRP1</i>)	No	No
KDEL (Lys-Asp-Glu-Leu) endoplasmic reticulum protein retention receptor 2 (<i>KDELR2</i>)	No	No	Tropomyosin 2 (<i>TPM2</i>)	No	No
Lactate dehydrogenase A, (<i>LDHA</i>)	No	Yes	Thymosin beta 4, X-linked (<i>TMSB4</i>)	Yes	Yes
Low density lipoprotein receptor-related protein 4 (<i>LRP4</i>)	Yes	Yes	Triose phosphate isomerase 1 (<i>TPI1</i>)	No	Yes

Lumican (<i>LUM</i>)	Yes	Yes	Vacuolar H ⁺ ATPase subunit V0 D2 (<i>ATP6V0D2</i>)	Yes	No
Microfibrillar associated protein 2 (<i>MFAP2</i>)	No	Yes	WD repeat and soxs box-containing 2 (<i>WSB2</i>)	No	No
Mitochondrial ribosome recycling factor (<i>MRRF</i>)	No	No	Wilms tumour 1-associating protein, isoform 1 (<i>WTAP</i>)	No	No

564
565
566

1 – Genes confirmed in qPCR analysis as being more highly expressed in lesions than in normal cartilage
2 – Genes already known to be expressed in cartilage or by chondrocytes (on the basis of a PubMed search)

For Peer Review

567 **SUPPLEMENTARY TABLE 3: Novel cartilage-associated genes.** Genes not previously
568 identified in cartilage, grouped into categories according to known functions.

Category	Gene
Chromatin-associated factors	Ankyrin repeat, family a, 2 (<i>ANKRA2</i>)
	Chromatin target of PRMT (<i>CHTOP</i>)
	Splicing factor, arginine/serine-rich 3 (<i>SRSF3</i>)
	Splicing factor 3b, subunit 1 (<i>SF3B1</i>)
	Structure specific recognition protein 1 (<i>SSRP1</i>)
Ion channels	Chloride intracellular channel protein 1 (<i>CLIC1</i>)
	Vacuolar H ⁺ ATPase subunit V0 D2 (<i>ATP6V0D2</i>)
Mitochondrial factors	Mitochondrial ribosome recycling factor (<i>MRRF</i>)
	Mitofusin I (<i>MFN1</i>)
	NADH dehydrogenase subunit 2, mitochondrial (<i>MT-ND2</i>)
Poorly characterised	Non-Hodgkins Lymphoma repeat containing 3 (<i>NHLRC3</i>)
	Protein prenyl transferase alpha subunit-containing 1 (<i>PTARI</i>)
	Ring finger protein 130 (<i>RNF130</i>)
	WD repeat and socs box-containing 2 (<i>WSB2</i>)
Protein secretion and trafficking	KDEL (Lys-Asp-Glu-Leu) endoplasmic reticulum protein retention receptor 2 (<i>KDELRL2</i>)
	Nascent polypeptide-associated complex alpha subunit (<i>NACA</i>)
	Signal sequence receptor, delta (<i>SSR4</i>)
Regulation of transcription	DEAD (Asp-Glu-Ala-Asp) box helicase 5 (<i>DDX5</i>)
	Human immunodeficiency virus type I enhancer binding protein 2 (<i>HIVEP2</i>)
	Potassium channel tetramerisation domain containing 10 (<i>KCTD10</i>)
	Purine-rich element binding protein B (<i>PURB</i>)
Ribosomal proteins	Ribosomal protein S7 (<i>RPS7</i>)
	Ribosomal protein S19 (<i>RPS19</i>)
Intracellular signalling	Guanine nucleotide binding protein beta polypeptide 1 (<i>GNB1</i>)
	Phosphatidylinositol-4-phosphate 5-kinase type II alpha (<i>PIP4K2A</i>)
Translation and post-translational modification	Eukaryotic translation elongation factor 1 gamma (<i>EEF1G</i>)
	Golgi phosphoprotein 3 (<i>GOLPH3</i>)
	Ribophorin II (<i>RPN2</i>)
Others	Cyclin-dependent kinase 11A (<i>CDK11A</i>)
	Nudix (nucleoside diphosphate linked moiety X)-type motif 16 (<i>NUDT16</i>)
	Prosaposin (<i>PSAP</i>)
	RAS oncogene family member (<i>RAP1B</i>)
	Tropomyosin 2 (<i>TPM2</i>)
	Wilms tumour 1-associating protein, isoform 1 (<i>WTAP</i>)

569

570

1
2 571 **SUPPLEMENTARY FIGURE LEGEND**

3 572

4 573

5 574

6 575

7 576

8 577

9 578

Figure S-1: Light microscopy image of the normal foetal articular-epiphyseal growth cartilage specimen shown in Figure 2C, showing the presence of the different chondrocyte zones. Semi-thin section of Spurr's resin-embedded tissue, stained with methylene blue. P – zone of proliferative chondrocytes; H – zone of hypertrophic chondrocytes; O – ossification front.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review




Figure 1: Genes differentially expressed in osteochondrosis lesions.

QPCR results showing MNE (\pm SD) of genes in osteochondrosis samples compared with control samples. Significant differences between lesions and control cartilage are indicated as * = $p < 0.05$; $n = 5$ horses.

57x40mm (600 x 600 DPI)




Figure 2: Morphology of chondrocytes in equine osteochondrosis lesions.

TEM images of representative chondrocytes from: the deep zone of normal articular cartilage (A), an osteochondrosis lesion from the same shoulder joint as the cartilage shown in A (B), and normal foetal articular-epiphyseal growth cartilage from the head of the humerus of a 9-month equine foetus (C). The images presented in A and B are all from a single animal, but are representative of cells observed in specimens from 3 animals. Arrows indicate the border of the chondrocyte lacuna; n - nucleus. All images have the same magnification; bar = 5 μ m.

144x125mm (300 x 300 DPI)





Figure 3: ATP6V0D2 and TMSB4 expression in osteochondrosis lesions. Immunohistochemistry was carried out on paraffin sections of equine osteochondrosis lesions with adjacent normal articular cartilage. A: Haematoxylin and eosin staining of a specimen taken from the lateral trochlear ridge of the distal end of the femur of a six month old foal, indicating the regions shown in images B-G. Normal articular cartilage overlying subchondral bone can be seen to the left of the image, and the lesion can be seen to the right, with cartilage retained in subchondral bone (arrows indicate bone-cartilage interface). Normal region (B, C, F, G) and lesion (D, E) stained with anti-ATP6V0D2 (B, D), anti-TMSB4 (C, E) or normal rabbit serum (NRS; F, G). Arrowheads (inset in D, E) indicate the border of the chondrocyte lacuna. Bar in A = 500 μ m. The magnification for parts B - G is the same; bar = 200 μ m for main image and 25 μ m for insets. 199x305mm (300 x 300 DPI)



Supplementary Figure 1: Light microscopy image of the normal foetal articular-epiphyseal growth cartilage specimen shown in Figure 2C, showing the presence of the different chondrocyte zones. Semi-thin section of Spurr's resin-embedded tissue, stained with methylene blue. P – zone of proliferative chondrocytes; H – zone of hypertrophic chondrocytes; O – ossification front.